

DAFTAR PUSTAKA

- Amanda, D. A. (2020) 'Rasio Neutrofil-Limfosit pada Covid-19; Sebuah tinjauan literatur', *Wellness And Healthy Magazine*, 2(2), pp. 219–223. doi: 10.30604/well.0202.8200100.
- Andrea Kuczynski et al (2017) 'Ischemic-Stroke-Pathogenesis'. on www.thecalgaryguide.com.
- Anrather, J. and Iadecola, C. (2016) 'Inflammation and Stroke: An Overview', *Neurotherapeutics*. *Neurotherapeutics*, 13(4), pp. 661–670. doi: 10.1007/s13311-016-0483-x.
- Bandera, E. *et al.* (2006) 'Cerebral blood flow threshold of ischemic penumbra and infarct core in acute ischemic stroke: A systematic review', *Stroke*, 37(5), pp. 1334–1339. doi: 10.1161/01.STR.0000217418.29609.22.
- Benítez Brito, N. *et al.* (2016) 'Relationship between mid-upper arm circumference and body mass index in inpatients', *PLoS ONE*, 11(8), pp. 1–10. doi: 10.1371/journal.pone.0160480.
- Boehme, A. K., Esenwa, C. and Elkind, M. S. V (2017) 'Stroke Risk Factors , Genetics , and Prevention', pp. 472–496. doi: 10.1161/CIRCRESAHA.116.308398.
- Burgos, R. *et al.* (2018) 'ESPEN guideline clinical nutrition in neurology', *Clinical Nutrition*, 37(1), pp. 354–396. doi: 10.1016/j.clnu.2017.09.003.
- Care, N. *et al.* (2016) 'Clinical Value of Neutrophil to Lymphocyte and Platelet to Lymphocyte Ratio After Aneurysmal Subarachnoid Hemorrhage'. doi: 10.1007/s12028-016-0332-0.
- Corrigan, M. L. *et al.* (2011) 'Nutrition in the stroke patient', *Nutrition in Clinical Practice*, 26(3), pp. 242–252. doi: 10.1177/0884533611405795.
- Das, P. *et al.* (2018) 'The validity of mid-upper arm circumference as an indicator of low BMI in population screening for undernutrition: A study among adult slum dwellers in eastern India', *Public Health Nutrition*, 21(14), pp. 2575–2583. doi: 10.1017/S1368980018001301.
- Dhian Ayudhia Kasim, dkk (2016) 'Asupan Makanan , Status Gizi Dan Lama Hari Rawat Inap Pada Pasien Penyakit Dalam Di Rumah Sakit Advent Manado', *Gizido*, 8(2), pp. 22–34.
- Dian Saraswati, R. (2021) 'Transisi Epidemiologi Stroke sebagai Penyebab Kematian pada Semua Kelompok Usia di Indonesia', *Seminar Nasional Riset Kedokteran (SENSORIK II*, 2(1), p. 2021. Available at: <https://conference.upnvj.ac.id/index.php/sensorik/article/view/1001>.

Esenwa, C. C. and Elkind, M. S. (2016) 'Inflammatory risk factors, biomarkers and associated therapy in ischaemic stroke', *Nature Reviews Neurology*. Nature Publishing Group, 12(10), pp. 594–604. doi: 10.1038/nrneuro.2016.125.

Ghani, L., Mihardja, L. K. and Delima, D. (2016) 'Faktor Risiko Dominan Penderita Stroke di Indonesia', *Buletin Penelitian Kesehatan*, 44(1), pp. 49–58. doi: 10.22435/bpk.v44i1.4949.49-58.

Gomes, F., Emery, P. W. and Weekes, C. E. (2016) 'Risk of Malnutrition Is an Independent Predictor of Mortality, Length of Hospital Stay, and Hospitalization Costs in Stroke Patients', *Journal of Stroke and Cerebrovascular Diseases*. Elsevier Inc., 25(4), pp. 799–806. doi: 10.1016/j.jstrokecerebrovasdis.2015.12.017.

Harris, S. *et al.* (2018) 'TOAST Subtypes of Ischemic Stroke and Its Risk Factors: A Hospital-Based Study at Cipto Mangunkusumo Hospital, Indonesia', *Stroke Research and Treatment*. doi: 10.1155/2018/9589831.

Holil M. Par'i, S.K.M., M.Kes, et al (2017) *Penilaian Status Gizi*. Oktober 20. Edited by N. Thamaria. Jakarta: Kementerian Kesehatan Republik Indonesia.

Hou, Y. *et al.* (2019) 'Association between mid-upper arm circumference and cardiometabolic risk in Chinese population: A cross-sectional study', *BMJ Open*, 9(9), pp. 1–8. doi: 10.1136/bmjopen-2019-028904.

Hua, J. and Zhou, W. (2020) 'Elevated neutrophil-to-lymphocyte and platelet-to-lymphocyte ratios predict post-stroke depression with acute ischemic stroke', *EXPERIMENTAL AND THERAPEUTIC MEDICINE*. Jiangsu, China, 19, pp. 2497–2504. doi: 10.3892/etm.2020.8514.

Hunaifi, I. and Cahyawati, T. D. (2020) 'Korelasi Antara Rasio Neutrofil Limfosit Dengan Volume Infark Serebri Pada Penderita Stroke Iskemik Akut', *Majalah Kedokteran Neurosains Perhimpunan Dokter Spesialis Saraf Indonesia*, 36(4). doi: 10.52386/neurona.v36i4.87.

'Indikator_Kinerja_Rumah_Sakit_Menurut_Depkes. RI' (2015).

Indonesia, P. D. S. G. K. (2019) *PEDOMAN NASIONAL PELAYANAN KEDOKTERAN (PNPK) Tatalaksana Malnutrisi Dewasa, Kemenkes RI*.

Irdelia, R. R., Tri Joko, A. and Bebasari, E. (2014) 'Profil Faktor Risiko yang dapat dimodifikasi pada Kasus Stroke Berulang di RSUD Arifin Achmad Provinsi Riau', *Jom FK*, 1, pp. 1–15.

Irisawa, T. *et al.* (2013) 'An association between systolic blood pressure and stroke among patients with impaired consciousness in out-of-hospital emergency settings', *BMC Emergency Medicine*, 13(1), pp. 2–5. doi: 10.1186/1471-227X-13-24.

Jickling, G. C. *et al.* (2015) 'Targeting neutrophils in ischemic stroke : translational

insights from experimental studies', (March), pp. 888–901. doi: 10.1038/jcbfm.2015.45.

Katakami, N. (2018) 'Mechanism of development of atherosclerosis and cardiovascular disease in diabetes mellitus', *Journal of Atherosclerosis and Thrombosis*, 25(1), pp. 27–39. doi: 10.5551/jat.RV17014.

Kaya, T. *et al.* (2019) 'Association between neutrophil-to-lymphocyte ratio and nutritional status in geriatric patients', *Journal of Clinical Laboratory Analysis*, 33(1), pp. 1–6. doi: 10.1002/jcla.22636.

Kemenkes RI (2018) 'Hasil Utama Riskesdas 2018'.

Kim, S. M. *et al.* (2013) 'Determinants of the Length of Stay in Stroke Patients', *Osong Public Health and Research Perspectives*. Elsevier Korea LLC, 4(6), pp. 329–341. doi: 10.1016/j.phrp.2013.10.008.

Knops, M. *et al.* (2013) 'Investigation of changes in body composition, metabolic profile and skeletal muscle functional capacity in ischemic stroke patients: The rationale and design of the Body Size in Stroke Study (BoSSS)', *Journal of Cachexia, Sarcopenia and Muscle*, 4(3), pp. 199–207. doi: 10.1007/s13539-013-0103-0.

Lubis, I. K. and Susilawati, S. (2018) 'Analisis Length Of Stay (Los) Berdasarkan Faktor Prediktor Pada Pasien DM Tipe II di RS PKU Muhammadiyah Yogyakarta', *Jurnal Kesehatan Vokasional*, 2(2), p. 161. doi: 10.22146/jkesvo.30330.

Luo, L. *et al.* (2019) 'Predicting intracerebral hemorrhage patients' length-of-stay probability distribution based on demographic, clinical, admission diagnosis, and surgery information', *Journal of Healthcare Engineering*, 2019. doi: 10.1155/2019/4571636.

Manurung, M., Diani, N. and Agianto (2015) 'Analisis Faktor Risiko Stroke Pada Pasien Stroke', *Jurnal Dunia Keperawatan*, 3(1), pp. 3761–3770.

Maulina, M. (2015) 'Gambaran Karakteristik Dan Status Gizi Berdasarkan Lingkar Lengan Atas (Lila) Pada Pasien Penyakit Jantung Koroner (Pjk) Di Rumah Sakit Umum Cut Meutia', *Lentera*, 15(13), pp. 29–36.

Mir, M. A., Al-Baradie, R. S. and Alhussainawi, M. D. (2015) *Pathophysiology of Strokes*.

Nastiti, D. (2011) 'Gambaran Faktor Risiko Kejadian Stroke Pada Pasien Stroke Rawat Inap Di Rumah Sakit Krakatau Medika Tahun 2011 Pasien Stroke Rawat Inap Di Rumah Sakit Krakatau Medika Tahun 2011', *Universitas Indonesia*.

Of, I. J. (2016) 'CLINICAL PATHOLOGY AND Majalah Patologi Klinik Indonesia dan Laboratorium Medik CLINICAL PATHOLOGY AND Majalah Patologi Klinik Indonesia dan Laboratorium Medik', 23(1).

Parmar, P. (2018) 'Stroke: Classification and diagnosis', *Clinical Pharmacist*, 10(1). doi: 10.1211/CP.2018.20204150.

Probowati, W., Bayupurnama, P. and Ratnasari, N. (2017) 'Correlation Between Neutrophil To Lymphocyte Ratio With Child Turcotte Pugh In Liver Cirrhosis Patients', *Acta Interna: The Journal of Internal Medicine*, 6(1), pp. 28–35. doi: 10.22146/actainterna.27003.

Ropper, A. H., Samuels, M. A. and Klein, J. P. (2014) *Principles of Neurology Adams and Victor's 10th ed, Archives of Otolaryngology - Head and Neck Surgery*.

Sabbouh, T. and Torbey, M. T. (2018) 'Malnutrition in Stroke Patients: Risk Factors, Assessment, and Management', *Neurocritical Care*. Springer US, 29(3), pp. 374–384. doi: 10.1007/s12028-017-0436-1.

Saxena, A. and Prasad, R. N. (2016) 'Factors Predicting Length of Hospital Stay in Acute Stroke Patients Admitted in a Rural Tertiary Care Hospital', *Journal of Gerontology & Geriatric Research*, 01(s5), pp. 2–6. doi: 10.4172/2167-7182.s5-003.

Şengeze, N. and Giray, S. (2020) 'The relationship between mean platelet volume, platelet count, platelet lymphocyte ratio, and recanalization success in first-pass thrombectomy of middle cerebral artery occlusions', *Turk Noroloji Dergisi*, 26(2), pp. 133–137. doi: 10.4274/tnd.2020.87405.

Severity, S. (2019) 'HUBUNGAN LUARAN SUBJECTIVE GLOBAL ASSESSMENT (SGA) DENGAN DERAJAT KEPARAHAN STROKE'.

Sharma, D. and Gandhi, N. (2021) 'Role of platelet to lymphocyte ratio (plr) and its correlation with nihss (national institute of health stroke scale) for prediction of severity in patients of acute ischemic stroke', *Journal of Association of Physicians of India*, 69(1), pp. 56–60.

Sid Gilman (2007) *Neurobiology of Disease*. London: Elsevier.

Simats, A., García-Berrocso, T. and Montaner, J. (2016) 'Neuroinflammatory biomarkers: From stroke diagnosis and prognosis to therapy', *Biochimica et Biophysica Acta - Molecular Basis of Disease*. Elsevier B.V., 1862(3), pp. 411–424. doi: 10.1016/j.bbadis.2015.10.025.

Smajlović, D. (2015) 'Strokes in young adults: Epidemiology and prevention', *Vascular Health and Risk Management*, 11, pp. 157–164. doi: 10.2147/VHRM.S53203.

Song, S. *et al.* (2019) 'Clinical Significance of Baseline Neutrophil-to-Lymphocyte Ratio in Patients With Ischemic Stroke or Hemorrhagic Stroke : An Updated', 10(October). doi: 10.3389/fneur.2019.01032.

Song, S. Y. *et al.* (2019) 'Clinical Significance of Baseline Neutrophil-to-

Lymphocyte Ratio in Patients With Ischemic Stroke or Hemorrhagic Stroke: An Updated Meta-Analysis', *Frontiers in Neurology*, 10(October). doi: 10.3389/fneur.2019.01032.

Subagio, H. W., Dianggra, P. S. and Himawan, M. R. A. (2018) 'Lama Pencapaian Target Energi dalam Masa Perawatan Pasien Stroke', *Jnh (Journal of Nutrition and Health)*, 6(2), p. 57. doi: 10.14710/jnh.6.2.2018.57-64.

Tao, C. *et al.* (2017) 'Clinical Value of Neutrophil to Lymphocyte and Platelet to Lymphocyte Ratio After Aneurysmal Subarachnoid Hemorrhage', *Neurocritical Care*, 26(3), pp. 393–401. doi: 10.1007/s12028-016-0332-0.

Tedja, V. R. (2012) 'Hubungan Antara Faktor Individu, Sosial Demografi, dan Administrasi dengan Lama Hari Rawat Pasien Rawat Inap Rumah Sakit Pantai Indah Kapuk Tahun 2011', pp. 1–157.

Toprak, E. *et al.* (2017) 'Platelet-to-lymphocyte ratio: A new inflammatory marker for the diagnosis of preterm premature rupture of membranes', *Journal of the Turkish German Gynecology Association*. doi: 10.4274/jtgga.2017.0028.

Usrin, I., Mutiara, E. and Yusad, Y. (2013) 'Pengaruh Hipertensi Terhadap Kejadian Stroke Iskemik dan Stroke Hemoragik di Ruang Neurologi di Rumah Sakit Stroke Nasional (RSSN) Bukittinggi Tahun 2011', *Neliti*, 2, pp. 1–9. Available at: fmipa.umri.ac.id/wp-content/uploads/2016/09/Rendy-Pengaruh-Hipertensi-pada-Stroke-iskemik.pdf.

Wang, J. and Doré, S. (2007) 'Inflammation after intracerebral hemorrhage', *Journal of Cerebral Blood Flow and Metabolism*, 27(5), pp. 894–908. doi: 10.1038/sj.jcbfm.9600403.

Wang, Q., Tang, X. N. and Yenari, M. A. (2007) 'The inflammatory response in stroke', *Journal of Neuroimmunology*, 184(1–2), pp. 53–68. doi: 10.1016/j.jneuroim.2006.11.014.

Women's Hospitals, B. (2009) 'Ischemic stroke: pathophysiology and principles of localization', *Neurology*.

Xu, J. *et al.* (2020) 'Platelet-rich plasma and regenerative dentistry', *Australian Dental Journal*, 65(2), pp. 131–142. doi: 10.1111/adj.12754.

Yu, S. *et al.* (2018) 'Neutrophil to lymphocyte ratio and early clinical outcomes in patients with acute ischemic stroke', *Journal of the Neurological Sciences*, 387(January), pp. 115–118. doi: 10.1016/j.jns.2018.02.002.

Zhao, L., Dai, Q. and Chen, X. (2015) 'Neutrophil-to-Lymphocyte Ratio Predicts Length of Stay and Acute Hospital Cost in Patients with Acute Ischemic Stroke'. Elsevier Inc., pp. 4–9. doi: 10.1016/j.jstrokecerebrovasdis.2015.11.012.

Zheng, T. *et al.* (2015) 'Impact of early enteral nutrition on short term prognosis

after acute stroke', *Journal of Clinical Neuroscience*. Elsevier Ltd, 22(9), pp. 1473–1476. doi: 10.1016/j.jocn.2015.03.028.

Zhu, Y. *et al.* (2020) 'Mid-upper arm circumference as a simple tool for identifying central obesity and insulin resistance in type 2 diabetes', *PLoS ONE*, 15(5), pp. 1–13. doi: 10.1371/journal.pone.0231308.

