

## DAFTAR PUSTAKA

1. United States Census Bureau 2013, Census.Gov, viewed 01 September 2013, <<http://www.census.gov/population/international/data/idb/informationGateway.php>>.
2. Hip fracture incidence map, International Osteoporosis Foundation, viewed 20 November 2013, <<http://www.iofbonehealth.org/facts-and-statistics/hip-fracture-incidence-map>>.
3. Hagino T, Sato E, Tonotsuka H, Ochiai S, Tokai M, Hamada Y (2006) Prediction of ambulation prognosis in the elderly after hip fracture. *Int Orthop* 30:315–319
4. Hagino, Tetsuo, et al. “A Simple Scoring System to Predict Ambulation Prognosis after Hip Fracture in the Elderly.” *Archives of Orthopaedic and Trauma Surgery*, vol. 127, no. 7, 2007, pp. 607–607., doi:10.1007/s00402-007-0376-x.
5. Ingemarsson, Annika Hellzén, et al. “Walking Ability And Activity Level After Hip Fracture In The Elderly - A Follow-Up.” *Journal of Rehabilitation Medicine*, vol. 35, no. 2, 2003, pp. 76–83., doi:10.1080/16501970306113.
6. Richmond J, Aharonoff GB, Zuckerman JD, Koval KJ (2003). Mortality risk after hip fracture. *J Orthop Trauma* 17:53–56
7. Hagino, T., Maekawa, S., Sato, E., Bando, K., & Hamada, Y. (2006). Prognosis of proximal FEMORAL fracture in PATIENTS aged 90 years and older. *Journal of Orthopaedic Surgery*, 14(2), 122–126. <https://doi.org/10.1177/230949900601400203>
8. Yang, Q.-H., Chen, Y.-X., Zhu, D.-Y., Ai, Z.-S., & Gao, Y.-S. (2019). Geographic variations in Intertrochanteric Femoral fractures in China. *BioMed Research International*, 2019, 1–8. <https://doi.org/10.1155/2019/8396723>

9. Chen, M., Zhang, Y., Du, Y., Hong, W., Tang, W., Li, H., Zheng, S., & Cheng, Q. (2019). Epidemiological and clinical study of hip fracture in hospitalized elderly patients in Shanghai, China. *Archives of Osteoporosis*, 14(1).  
<https://doi.org/10.1007/s11657-019-0580-7>
10. Nherera, L., Trueman, P., Horner, A., Watson, T., & Johnstone, A. J. (2018). Comparison of a twin interlocking derotation and compression screw cephalomedullary nail (intertan) with a single screw derotation cephalomedullary nail (proximal femoral nail antirotation): A systematic review and meta-analysis for intertrochanteric fractures. *Journal of Orthopaedic Surgery and Research*, 13(1).  
<https://doi.org/10.1186/s13018-018-0749-6>
11. Kesmezacar, H., Ayhan, E., Unlu, M. C., Seker, A., & Karaca, S. (2010). Predictors of mortality in elderly patients with AN INTERTROCHANTERIC or a femoral Neck Fracture. *Journal of Trauma: Injury, Infection & Critical Care*, 68(1), 153–158. <https://doi.org/10.1097/ta.0b013e31819adc50>
12. Sexson, S. B., & Lehner, J. T. (1987). Factors affecting hip fracture mortality. *Journal of Orthopaedic Trauma*, 1(4), 298–305. <https://doi.org/10.1097/00005131-198701040-00005>
13. Fazal, M. A., Bagley, C., & Garg, P. (2018). Predictors for perioperative blood transfusion in elderly patients with extra capsular hip fractures treated with cephalo-medullary nailing. *Chinese Journal of Traumatology*, 21(1), 16–19.  
<https://doi.org/10.1016/j.cjtee.2017.09.002>
14. Yoshii, I., Satake, Y., Kitaoka, K., Komatsu, M., & Hashimoto, K. (2016). Relationship between dementia degree and gait ability after surgery of proximal femoral fracture. *Journal of Orthopaedic Science*, 21(4), 481–486.  
<https://doi.org/10.1016/j.jos.2016.03.005>

15. Roshan, A., & Ram, S. (2008). The neglected femoral neck fracture in young adults: review of a challenging problem. *Clinical medicine & research*, 6(1), 33–39.  
<https://doi.org/10.3121/cmr.2008.752>
16. He, W., You, Y. Y., Sun, K., Xie, C., Ming, Y., Yu, L. N., Zhang, F. J., & Yan, M. (2020). Admission delay is associated with worse surgical outcomes for elderly hip fracture patients: A retrospective observational study. *World journal of emergency medicine*, 11(1), 27–32. <https://doi.org/10.5847/wjem.j.1920-8642.2020.01.004>
17. Ryan, Gareth BSc(Hons)1,2,a; Nowak, Lauren PhD1; Melo, Luana PhD1; Ward, Sarah MD, MSc, FRCSC1,2; Atrey, Amit MD, MSc, FRCSC1,2; Schemitsch, Emil H. MD, FRCSC3; Nauth, Aaron MD, MSc, FRCSC1,2; Khoshbin, Amir MD, MSc, FRCSC1,2 Anemia at Presentation Predicts Acute Mortality and Need for Readmission Following Geriatric Hip Fracture, *JBJS Open Access*: July-September 2020 - Volume 5 - Issue 3 - p e20.00048 doi: 10.2106/JBJS.OA.20.00048

## **LAMPIRAN**