

## DAFTAR PUSTAKA

1. KDIGO AKI Work Group. Kidney International Supplements. *KDIGO Clin Prat Guidel ACute Kidney Inj*. 2012;2(1):1-138. doi:10.1038/kisup.2012.3
2. Bellomo R, Ronco C, Kellum JA, et al. Acute renal failure - definition, outcome measures, animal models, fluid therapy and information technology needs: the Second International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Crit Care*. 2004;8(4). doi:10.1186/cc2872
3. Mehta RL, Kellum JA, Shah S V., et al. Acute kidney injury network: Report of an initiative to improve outcomes in acute kidney injury. *Crit Care*. 2007;11(2):1-8. doi:10.1186/cc5713
4. Kim M, Brady JE, Li G. Interaction Effects of Acute Kidney Injury, Acute Respiratory Failure, and Sepsis on 30-Day Postoperative Mortality in Patients Undergoing High-Risk Intraabdominal General Surgical Procedures. *Anesth Analg*. 2015;121(6):1536-1546. doi:10.1213/ANE.0000000000000915
5. Maheshwari K, Turan A, Mao G, et al. The association of hypotension during non-cardiac surgery, before and after skin incision, with postoperative acute kidney injury: a retrospective cohort analysis. *Anaesthesia*. 2018;73(10):1223-1228. doi:10.1111/anae.14416
6. Shim J-W, Kim KR, Jung Y, et al. Role of intraoperative oliguria in risk stratification for postoperative acute kidney injury in patients undergoing colorectal surgery with an enhanced recovery protocol: A propensity score matching analysis. *PLoS One*. 2020;15(4):e0231447. doi:10.1371/journal.pone.0231447
7. Toyonaga Y, Kikura M. Hyperchloremic acidosis is associated with acute kidney injury after abdominal surgery. *Nephrology (Carlton)*. 2017;22(9):720-727.
8. Olufajo OA, Reznor G, Lipsitz SR, et al. Preoperative assessment of surgical risk: creation of a scoring tool to estimate 1-year mortality after emergency abdominal surgery in the elderly patient. *Am J Surg*. 2017;213(4):771-777.e1. doi:10.1016/j.amjsurg.2016.08.007
9. Long TE, Helgason D, Helgadottir S, et al. Acute Kidney Injury After Abdominal Surgery: Incidence, Risk Factors, and Outcome. *Anesth Analg*. 2016;122(6):1912-1920. doi:10.1213/ANE.0000000000001323
10. Göcze I, Jauch D, Götz M, et al. Biomarker-guided Intervention to Prevent Acute Kidney Injury After Major Surgery: the Prospective Randomized BigpAK Study. *Ann Surg*. 2018;267(6):1013-1020. doi:10.1097/SLA.0000000000002485
11. Branch-Elliman W, O'Brien W, Strymish J, et al. Association of Duration and Type of Surgical Prophylaxis With Antimicrobial-Associated Adverse Events. *JAMA Surg*. 2019;154(7):590-598. doi:10.1001/jamasurg.2019.0569
12. Chaudery H, MacDonald N, Ahmad T, et al. Acute Kidney Injury and Risk of Death After Elective Surgery: Prospective Analysis of Data From an International Cohort Study. *Anesth Analg*. 2019;128(5):1022-1029. doi:10.1213/ANE.0000000000003923
13. Kopitkó C, Medve L, Gondos T. The value of combined hemodynamic , respiratory

- and intra-abdominal pressure monitoring in predicting acute kidney injury after major intraabdominal surgeries. *Ren Fail.* 2019;41(1):150-158. doi:10.1080/0886022X.2019.1587467
14. Wu Q, Yang H, Bo H, et al. Predictive role of estimated glomerular filtration rate prior to surgery in postsurgical acute kidney injury among very elderly patients: a retrospective cohort study. *Ren Fail.* 2019;41(1):866-874. doi:10.1080/0886022X.2019.1662440
  15. Tomozawa A, Ishikawa S, Shiota N, et al. Perioperative risk factors for acute kidney injury after liver resection surgery: an historical cohort study. *Can J Anesth.* 2015;62(7):753-761. doi:10.1007/s12630-015-0397-9
  16. Puckett JR, Pickering JW, Palmer SC, et al. Low Versus Standard Urine Output Targets in Patients Undergoing Major Abdominal Surgery: a Randomized Noninferiority Trial. *Ann Surg.* 2017;265(5):874-881. doi:10.1097/SLA.0000000000002044
  17. Wakeam E, Hyder JA, Jiang W, Lipsitz SA, Finlayson S. Risk and patterns of secondary complications in surgical inpatients. *JAMA Surg.* 2015;150(1):65-73. doi:10.1001/jamasurg.2014.1795
  18. Myles PS, Bellomo R, Corcoran T, et al. Restrictive versus liberal fluid therapy for major abdominal surgery. *N Engl J Med.* 2018;378(24):2263-2274. doi:10.1056/NEJMoa1801601
  19. Li S, Wang S, Priyanka P, Kellum JA. Acute Kidney Injury in Critically Ill Patients After Noncardiac Major Surgery: Early Versus Late Onset. *Crit Care Med.* 2019;47(6):e437-e444. doi:10.1097/CCM.00000000000003710
  20. Bang J-Y, Lee J, Oh J, et al. The Influence of Propofol and Sevoflurane on Acute Kidney Injury After Colorectal Surgery: A Retrospective Cohort Study. *Anesth Analg.* 2016;123(2):363-370. doi:10.1213/ANE.0000000000001274
  21. Kim M, Wall MM, Kiran RP, et al. Latent class analysis stratifies mortality risk in patients developing acute kidney injury after high-risk intraabdominal general surgery: a historical cohort study. *Can J Anesth.* 2019;66(1):36-47. doi:10.1007/s12630-018-1221-0
  22. Romagnoli S, Zagli G, Tuccinardi G, et al. Postoperative acute kidney injury in high-risk patients undergoing major abdominal surgery. *J Crit Care.* Published online 2016. doi:10.1016/j.jcrc.2016.05.012
  23. Oh TK, Do S, Kim J, et al. Association of Preoperative Serum Chloride Levels With Mortality and Morbidity After Noncardiac Surgery: A Retrospective Cohort Study. *anesthesia-analgesia.* 2018;XXX(XXX):1-8. doi:10.1213/ANE.0000000000003958
  24. C H, T O-B, A K, et al. Cost and Mortality Associated With Postoperative Acute Kidney Injury. *Ann Surg.* 2015;261(6):1207-1214. <https://pubmed.ncbi.nlm.nih.gov/24887982/>
  25. Sanaiha Y, Kavianpour B, Dobaria V, et al. Acute kidney injury is independently associated with mortality and resource use after emergency general surgery operations. *Surg (United States).* 2020;167(2):328-334. doi:10.1016/j.surg.2019.07.035

26. Joosten A, Tircoveanu R, Arend S, et al. Impact of balanced tetrastarch raw material on perioperative blood loss: a randomized double blind controlled trial. *Br J Anaesth*. 2016;117(4):442-449. doi:10.1093/bja/aew249
27. Schmid S, Kapfer B, Heim M, et al. Algorithm-guided goal-directed haemodynamic therapy does not improve renal function after major abdominal surgery compared to good standard clinical care: a prospective randomised trial. *Crit Care*. 2016;20:50.
28. Kyu T, Sunghee O, Oh AY, et al. Chronic hyperglycemia with elevated glycosylated hemoglobin level and its association with postoperative acute kidney injury after a major laparoscopic abdominal surgery in diabetes patients. *J Anesth*. 2018;0(0):0. doi:10.1007/s00540-018-2551-3
29. Wang W, Wang T, Feng X, et al. Incidence and risk factors of acute kidney injury after esophageal cancer surgery: A nested case-control study. *Int J Surg*. 2017;39:11-15. doi:10.1016/j.ijssu.2017.01.043
30. Engoren M, Maile MD, Heung M, et al. The Association Between Urine Output, Creatinine Elevation, and Death. *Ann Thorac Surg*. 2017;103(4):1229-1237. doi:10.1016/j.athoracsur.2016.07.036