

DAFTAR PUSTAKA

- Akhundov, K. *et al.* (2012) „Development of a cost-effective method for platelet-rich plasma (prp) preparation for topical wound healing“, *Annals of Burns and Fire Disasters*, 25(4), pp. 207–213.
- Cardoso, A. L. *et al.* (2016) „Adipose tissue stromal vascular fraction in the treatment of full thickness burns in rats“, *Acta Cirurgica Brasileira*, 31(9), pp. 578–585. doi: 10.1590/S0102-865020160090000002.
- Deng, C. *et al.* (2018) „Treatment of human chronic wounds with autologous extracellular matrix/stromal vascular fraction gel: A STROBE-compliant study“, *Medicine (United States)*, 97(32), pp. 1–7. doi: 10.1097/MD.00000000000011667.
- Eyuboglu, A. A. *et al.* (2018) „The effect of adipose derived stromal vascular fraction on stasis zone in an experimental burn model“, *Burns*. Elsevier Ltd and International Society of Burns Injuries, 44(2), pp. 386–396. doi: 10.1016/j.burns.2017.08.016.
- Fotouhi, A. *et al.* (2018) „Platelet rich plasma, stromal vascular fraction and autologous conditioned serum in treatment of knee osteoarthritis“, *Biomedicine and Pharmacotherapy*. Elsevier, 104(April), pp. 652–660. doi: 10.1016/j.biopha.2018.05.019.
- Hosny, N. *et al.* (2015) „Assessment of Vascular Endothelial Growth Factor in Fresh versus Frozen Platelet Rich Plasma“, *Journal of Blood Transfusion*. Hindawi

Publishing Corporation, 2015, pp. 1–5. doi: 10.1155/2015/706903.

Infanger, M. *et al.* (2004) „Vascular endothelial growth factor serum level is strongly enhanced after burn injury and correlated with local and general tissue edema“, *Burns*, 30(4), pp. 305–311. doi: 10.1016/j.burns.2003.12.006.

Karina *et al.* (2019) „Combination of the stromal vascular fraction and platelet-rich plasma accelerates the wound healing process: Pre-clinical study in a Sprague-Dawley rat model“, *Stem Cell Investigation*, 6(July), pp. 1–8. doi: 10.21037/sci.2019.06.08.

Kazakos, K. *et al.* (2009) „The use of autologous PRP gel as an aid in the management of acute trauma wounds“, *Injury*, 40(8), pp. 801–805. doi: 10.1016/j.injury.2008.05.002.

Kumar, P. *et al.* (2015) „Role of angiogenesis and angiogenic factors in acute and chronic wound healing“, *Plastic and Aesthetic Research*, 2(5), p. 243. doi: 10.4103/2347-9264.165438.

Liang, Z. J. *et al.* (2018) „Precise Intradermal Injection of Nanofat-Derived Stromal Cells Combined with Platelet-Rich Fibrin Improves the Efficacy of Facial Skin Rejuvenation“, *Cellular Physiology and Biochemistry*, 47(1), pp. 316–329. doi: 10.1159/000489809.

Liu, T. *et al.* (2019) „One-step approach for full-thickness skin defect reconstruction in rats using minced split-thickness skin grafts with Pelnac overlay“, *Burns & Trauma*. *Burns & Trauma*, 7, pp. 1–13. doi: 10.1186/s41038-019-0157-0.

- Liu, Z. *et al.* (2019) „Synergistic Effects of Human Platelet-Rich Plasma Combined with Adipose-Derived Stem Cells on Healing in a Mouse Pressure Injury Model“, *Stem Cells International*, 2019. doi: 10.1155/2019/3091619.
- Mazzocca, A. D. *et al.* (2012) „Platelet-Rich Plasma Differs According to“, *The Journal of bone and joint surgery*, 94-A(4), pp. 308–316.
- PIETRO, GENTILE; MARIA, GIOVANNA SCIOLI; ALESSANDRA, BIELLI; AUGUSTO, ORLANDI; VALERIO, C. (2017) „The use of Adipose-derived Stromal Vascular Fraction Cells and Platelet Rich Plasma in Re- generative Plastic Surgery“, *Stem cells (Dayton, Ohio)*, 29(9), pp. 1338–1348. doi: 10.1002/stem.
- Satoshi, Tajima; Morikuni, Tobita; Hakan, Orbay; Hiko, Hyakusoku; Hiroshi, M. (2014) „Direct and indirect effects on bone regeneration of a combination of adipose-derived stem cells and platelet-rich plasma“, *Tissue Engineering Part A*, 21(5–6), pp. 895–905.
- Seyhan, N. *et al.* (2015) „The Effect of Combined Use of Platelet-Rich Plasma and Adipose-Derived Stem Cells on Fat Graft Survival“, *Annals of Plastic Surgery*, 74(5), pp. 615–620. doi: 10.1097/SAP.0000000000000480.
- Sun, M. *et al.* (2017) „Adipose Extracellular Matrix/Stromal Vascular Fraction Gel Secretes Angiogenic Factors and Enhances Skin Wound Healing in a Murine Model“, *BioMed Research International*, 2017. doi: 10.1155/2017/3105780.
- Vaghardoost, R., Ghavami, Y. and Sobouti, B. (2019) „The Effect of Mentha Pulegium

on Healing of Burn Wound Injuries in Rat", *World Journal of Plastic Surgery*, 8(1), pp. 43–50. doi: 10.29252/wjps.8.1.43.

Zhang, X. G. *et al.* (2019) „The Wound Healing Effect of *Callicarpa nudiflora* in Scalded Rats", *Evidence-based Complementary and Alternative Medicine*, 2019. doi: 10.1155/2019/1860680.

Antonetti DA, Barber AJ, Hollinger LA, Wolpert EB, Gardner TW. Vascular endothelial growth factor induces rapid phosphorylation of tight junction proteins occludin and zonula occludens 1. A potential mechanism for vascular permeability in diabetic retinopathy and tumors. *J Biol Chem*, 1999, 274(33):23463–23467.

Arsic, N., Zacchigna, S., Zentilin, L., Ramirez-Correa, G., Pattarini, L., Salvi, A., Giacca, M. (2004). Vascular endothelial growth factor stimulates skeletal muscle regeneration in Vivo. *Molecular Therapy*, 10(5), 844–854. doi:10.1016/j.ymthe.2004.08.007

Aslı Duran, Şirin Yaşar, Sema Aytakin, et al., 2018. Clinical and histopathological evaluation of the effects of platelet rich plasma, platelet poor plasma and topical serum physiologic treatment on wound healing caused by radiofrequency electrosurgery in rats. *Turkderm-Turk Arch Dermatol Venereology* 2018;52:44-50

Baglioni, S., Francalanci, M., Squecco, R., Lombardi, A., Cantini, G., Angeli, R., Luconi, M. (2009). Characterization of human adult stem-cell populations isolated from visceral and subcutaneous adipose tissue. *The FASEB Journal*, 23(10), 3494–3505. doi:10.1096/fj.08-126946

Bakacak, M., Bostanci, M. S., İnanc, F., Yaylali, A., Serin, S., Attar, R., Yildirim, O. K. (2015). Protective Effect of Platelet Rich Plasma on Experimental

Ischemia/Reperfusion Injury in Rat Ovary. *Gynecologic and Obstetric Investigation*, 81(3), 225–231. doi:10.1159/000440617

Barret J, Herndon D., 2005. *Principles and Practice of Burn Surgery*. New York, USA. Marcel Dekker.

Barrientos S, Stojadinovic O, Golinko MS, et al: Growth factors and cytokines in wound healing. *Wound Rep Regen* 16: 585-601,2008.

Benson, A., W.A. Dickson, and D.E. Boyce, ABC of wound healing: burns. *British Medical Journal*, 2006. 332: p. 649-652.

Borrione, P., Gianfrancesco, A. D., Pereira, M. T., & Pigozzi, F. (2010). Platelet-Rich Plasma in Muscle Healing. *American Journal of Physical Medicine & Rehabilitation*, 89(10), 854–861.

Bosch, J., Abrales, J. G., Fernández, M., & García-Pagán, J. C. (2010). Hepatic endothelial dysfunction and abnormal angiogenesis: New targets in the treatment of portal hypertension. *Journal of Hepatology*, 53(3), 558–567.

Bourin P, Bunnell BA, Casteilla L, Dominici M, Katz AJ, March KL, Redl H, Rubin JP, Yoshimura K, Gimble JM. Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International Society for Cellular Therapy (ISCT). *Cytotherapy*. 2013;15:641–8.

Cébe Suarez, S; Pieren, M; Cariolato, L; Arn, S; Hoffmann, U; Bogucki, A; Manlius, C; Wood, J; Ballmer-Hofer, K (2006). "A VEGF-A splice variant defective for heparan sulfate and neuropilin-1 binding shows attenuated signaling through VEGFR-2" (PDF). *Cellular and Molecular Life Sciences*. 63 (17): 2067–2077

- Cervelli V, et al. Tissue regeneration in loss of substance on the lower limbs through use of platelet-rich plasma, stem cells from adipose tissue, and hyaluronic acid. *Adv Skin Wound Care*. 2010; 23:262-72.)
- Chieregato K, Castegnaro S, Madeo D, Astori G, Pegoraro M, Rodeghiero F: Epidermal growth factor, basic fibroblast growth factor and platelet-derived growth factor-bb can substitute for fetal bovine serum and compete with human platelet-rich plasma in the ex vivo expansion of mesenchymal stromal cells derived from adipose tissue. *Cytotherapy* 2011, 13:933–943.
- Choi, J., Minn, K. W., & Chang, H. (2012). The Efficacy and Safety of Platelet-Rich Plasma and Adipose-Derived Stem Cells: An Update. *Archives of Plastic Surgery*, 39(6), 585. doi:10.5999/aps.2012.39.6.585
- Cohen AW, Carbajal JM, Schaeffer RC Jr. VEGF stimulates tyrosine phosphorylation of beta-catenin and small-pore endothelial barrier dysfunction. *Am J Physiol*, 1999, 277(5 Pt 2): H2038–H2049.
- Comella, K., Silbert, R., & Parlo, M. (2017). Effects of the intradiscal implantation of stromal vascular fraction plus platelet rich plasma in patients with degenerative disc disease. *Journal of Translational Medicine*, 15(1).
- Darinskas, A., Paskevicius, M., Apanavicius, G., Vilkevicius, G., Labanauskas, L., Ichim, T. E., & Rimdeika, R. (2017). Stromal vascular fraction cells for the treatment of critical limb ischemia: a pilot study. *Journal of Translational Medicine*, 15(1). doi:10.1186/s12967-017-1243-3
- DeJana E, Bazzoni G, Lampugnani MG. Vascular endothelial (VE)-cadherin: only an intercellular glue? *Exp Cell Res*, 1999, 252(1):13–19.
- Duffy AM, Bouchier-Hayes DJ, Harmey JH. Vascular endothelial growth factor (VEGF) and its role in non-endothelial cells: autocrine signalling by VEGF. In: *Madame Curie Bioscience Database (formerly, Eureka Bioscience Database). Angiogenesis*. Landes Bioscience, Austin (TX), USA, 2000–2013.

- El-Sharkawy, H., Kantarci, A., Deady, J., Hasturk, H., Liu, H., Alshahat, M., & Van Dyke, T. E. (2007). Platelet-Rich Plasma: Growth Factors and Pro- and Anti-Inflammatory Properties. *Journal of Periodontology*, 78(4), 661–669.
- Emergency Management of Severe Burns (EMSB) Course Manual, 17th Edition, 2013. ANZBA Ltd.
- Eppley BL, Pietrzak WS, Blanton M. Platelet-rich plasma: a review of biology and applications in plastic surgery. *Plast Reconstr Surg* 2006;118:147e-159e
- Esser S, Lampugnani MG, Corada M, DeJana E, Risau W. Vascular endothelial growth factor induces VE-cadherin tyrosine phosphorylation in endothelial cells. *J Cell Sci*, 1998, 111(Pt 13): 1853–1865.
- Ferrara N. From the discovery of vascular endothelial growth factor to the introduction of Avastin in clinical trials – an interview with Napoleone Ferrara by Domenico Ribatti. *Int J Dev Biol*, 2011, 55(4–5):383–388.
- Ferrara N. Vascular endothelial growth factor: basic science and clinical progress. *Endocr Rev*, 2004, 25(4):581–611.
- Ferraro, G. A., Mizuno, H., & Pallua, N. (2016). Adipose Stem Cells: From Bench to Bedside. *Stem Cells International*, 2016, 1–2. doi:10.1155/2016/6484038
- Folkman J. Angiogenesis in cancer, vascular, rheumatoid and other disease. *Nat Med*, 1995, 1(1):27–31.
- Friedenstein AJ, Piatetzky S II, Petrakova KV. Osteogenesis in transplants of bone marrow cells. *J Embryol Exp Morphol*. 1966; 16:381–90. [PubMed: 5336210]
- Gentile, P., Scioli, M. G., Bielli, A., Orlandi, A., & Cervelli, V. (2016). Concise Review: The Use of Adipose-Derived Stromal Vascular Fraction Cells and Platelet Rich Plasma in Regenerative Plastic Surgery. *STEM CELLS*, 35(1), 117–134.

- Germani A, Di Carlo A, Mangoni A, et al. Vascular endothelial growth factor modulates skeletal myoblast function. *Am J Pathol.* 2003;163(4):1417–1428. doi:10.1016/S0002-9440(10)63499-2
- Ghieh, F., Jurjus, R., Ibrahim, A., Geagea, A. G., Daouk, H., El Baba, B., ... Jurjus, A. (2015). The Use of Stem Cells in Burn Wound Healing: A Review. *BioMed Research International*, 2015, 1–9. doi:10.1155/2015/684084
- Gimble J, Guilak F. Adipose-derived adult stem cells: isolation, characterization, and differentiation potential. *Cytotherapy.* 2003; 5:362–9. [PubMed: 14578098]
- Gimble JM, Katz AJ, Bunnell BA. Adipose-derived stem cells for regenerative medicine. *Circ Res.* 2007; 100:1249–60. [PubMed: 17495232]
- Guo HF, Ali RM, Hamid RA, Zaini AA, Khaza'ai H. A new model for studying deep partial-thickness burns in rats. *Int J Burns Trauma.* 2017;7(6):107–114. Published 2017 Oct 25.
- Halim D, dkk., 2010. *Stem Cell: Dasar Teori dan Aplikasi Klinis.* Surabaya, Indonesia. Erlangga.
- Han J, Koh YJ, Moon HR, Ryoo HG, Cho CH, Kim I, Koh GY. Adipose tissue is an extramedullary reservoir for functional hematopoietic stem and progenitor cells. *Blood.* 2010;115:957–64.
- Harris, P.N., S. & Vardaxis, N., ed. *Mosby's Dictionary Of Medicine, Nursing & Health Professionals.* 2nd ed. 2010, Elsevier: Sydney.
- Hausman GJ, Richardson RL. Adipose tissue angiogenesis. *J Anim Sci* 2004;82:925-34.
- Herndon, D.N., ed. *Total Burn Care.* 3rd ed. 2007, Saunders: London.
- Hettiaratchy, S. and R. Papini, Initial management of a major burn: II--assessment and resuscitation. *BMJ,* 2004. 329(7457): p. 101-3.

- Hollenberg CH, Vost A. Regulation of DNA synthesis in fat cells and stromal elements from rat adipose tissue. *J Clin Invest.* 1969; 47:2485–98. [PubMed: 4304653]
- Holmes, Katherine; Roberts, Owain Ll; Thomas, Angharad M.; Cross, Michael J. (2007). "Vascular endothelial growth factor receptor-2: Structure, function, intracellular signalling and therapeutic inhibition". *Cellular Signalling.* 19 (10): 2003–12.
- Horwitz EM, Le Blanc K, Dominici M, et al. Clarification of the nomenclature for MSC: The International Society for Cellular Therapy position statement. *Cytotherapy* 2005; 7:393-5.
- Hu Z, Peel SA, Ho SK, et al. Platelet-rich plasma induces mRNA expression of VEGF and PDGF in rat bone marrow stromal cell differentiation. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;107:43-8.
- Ivan, M., Kondo, K., Yang, H. et al. (2001) HIF α targeted for VHL-mediated destruction by proline hydroxylation: implications for O₂ sensing. *Science* 292, 464–468
- Jaakkola, P., Mole, D. R., Tian, Y. M. et al. (2001) Targeting of HIF- α to the von Hippel–Lindau ubiquitylation complex by O₂-regulated prolyl hydroxylation. *Science* 292, 468–472.
- Josh F, et al., 2012. Accelerated and safe proliferation of human adipose-derived stem cells in medium supplemented with human serum. Tokyo, Japan. *J Nippon Med* (79): pp. 444-452. edition. Denver, USA. RC Press.
- Karagergou, E., Dionyssopoulos, A., Karayannopoulou, M., Psalla, D., Theodoridis, A., Demiri, E., & Koliakos, G. (2018). Adipose-derived stromal vascular fraction aids epithelialisation and angiogenesis in an animal model. *Journal of Wound Care,* 27(10), 637–644.

- Karkkainen, M.J.; Petrova, T.V. (2000). "Vascular endothelial growth factor receptors in the regulation of angiogenesis and lymphangiogenesis". *Oncogene*. 19 (49): 5598–5605
- Kevy S, Jacobson M, Benoit P. The biology of platelet concentrate as prepared by the Harvest Technologies SmartPREP System. Proceedings of the 3rd Annual Meeting of Techvest Conference on Tissue Repair, Replacement, and Regeneration; 2001 Oct 23; New York, USA
- Kim Yeol, et al., 2014. Effects of Platelet-Rich Plasma, Adipose-Derived Stem Cells and Stromal Vascular Fraction on the Survival of Human Transplanted Adipose Tissue. Seoul, Republic of Korea. *J Korean Med Science* (29): pp. 193-200.
- Koch S, Claesson-Welsh L. Signal transduction by vascular endothelial growth factor receptors. *Cold Spring Harb Perspect Med*, 2012, 2(7):a006502.
- Kuwada, S. K., & Li, X. (2000). Integrin $\alpha 5/\beta 1$ Mediates Fibronectin-dependent Epithelial Cell Proliferation through Epidermal Growth Factor Receptor Activation. *Molecular Biology of the Cell*, 11(7), 2485–2496. doi:10.1091/mbc.11.7.2485
- Leong M., Phillips LG., Wound Healing: Chapter 7 (2016) <https://thoracickey.com/wound-healing/>
- Lester SC. The Breast in Text book of Robbin's Pathologic Basic of Disease; Chapter 23. 8th Ed. Phil. W.B Saunders Co. 2010; p: 1083-90.
- Li H, Liu D, Yu Y, Wu T: Experimental research of the promotion effect of autogeneic PRP on osteogenic differentiation of human adipose-derived stem cells in vitro. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi* 2009, 23:732–736
- MacLeod, A. S., & Mansbridge, J. N. (2016). The Innate Immune System in Acute and Chronic Wounds. *Advances in Wound Care*, 5(2), 65–78.

- Maharaj AS, Saint-Geniez M, Maldonado AE, D'Amore PA. Vascular endothelial growth factor localization in the adult. *Am J Pathol*, 2006, 168(2):639–648
- Martins, R. P., Hartmann, D. D., de Moraes, J. P., Soares, F. A. A., & Puntel, G. O. (2016). Platelet-rich plasma reduces the oxidative damage determined by a skeletal muscle contusion in rats. *Platelets*, 27(8), 784–790.
- Matthews, W., Jordan, C. T., Gavin, M., Jenkins, N. A., Copeland, N. G. and Lemischka, I. R. (1991) A receptor tyrosine kinase cDNA isolated from a population of enriched primitive hematopoietic cells and exhibiting close genetic linkage to c-kit. *Proc. Natl. Acad. Sci. U.S.A.* 88, 9026–9030
- Medford ARL, Douglas SK, Godinho SIH, Uppington KM, Armstrong L, Gillespie KM, van Zyl B, Tetley TD, Ibrahim NBN, Mill AB. Vascular endothelial growth factor (VEGF) isoform expression and activity in human and murine lung injury. *Respir Res*, 2009, 10(1):27.
- Melincovici, Carmen Stanca et al. “Vascular endothelial growth factor (VEGF) - key factor in normal and pathological angiogenesis.” *Romanian journal of morphology and embryology = Revue roumaine de morphologie et embryologie* 59 2 (2018): 455-467 .
- Miron L, Gafton B, Marinca M. Angiogeneza tumorală –implicații în terapia cancerelor. *Jurnalul de Chirurgie, Iași*, 2010, 6(2):104–110.
- Moenadjat Y, dkk., 2011. *Luka Bakar: Masalah dan Tatalaksana*, edisi 4. Jakarta, Indonesia. Balai Penerbit FKUI.
- Moss LS. Treatment of the burn patient in primary care. *Adv Skin Wound Care*, 2010. 23(11): 517–24;quiz 525–6.
- Nauta A, et al., 2012. Adipose-derived Stromal Cells Overexpressing Vascular Endothelial Growth Factor Accelerate Mouse Excisional Wound Healing.

California, USA. The American Society of Gene & Cell Therapy.
10.1038/mt.2012.234.

Neufeld G, Cohen T, Gengrinovitch S, Poltorak Z. Vascular endothelial growth factor (VEGF) and its receptors. *FASEB J*, 1999, 13(1):9–22.

Nikolidakis, D., & Jansen, J. A. (2008). The Biology of Platelet-Rich Plasma and Its Application in Oral Surgery: Literature Review. *Tissue Engineering Part B: Reviews*, 14(3), 249–258. doi:10.1089/ten.teb.2008.0062

Niu, Gang, and Xiaoyuan Chen. “Vascular endothelial growth factor as an anti-angiogenic target for cancer therapy.” *Current drug targets* vol. 11,8 (2010): 1000-17.

Olsson AK, Dimberg A, Kreuger J, Claesson-Welsh L. VEGF receptor signalling – in control of vascular function. *Nat Rev Mol Cell Biol*, 2006, 7(5):359–371.

Pajusola K, Aprelikova O, Pelicci G, Weich H, ClaessonWelsh L and Alitalo K. (1994). *Oncogene*, 9, 3545 ± 3555.

Prosnitz LR, Iglehart JD, Winer EP. Breast Cancer. In : Rubin P, editors. *Clinical Oncology, A Multidisciplinary Approach for Physicians and Students*, 8th ed. WB Saunders Co. 2001; p: 267-96.

Qiu Y, Hoareau-Aveilla C, Oltean S, Harper SJ, Bates DO. The anti-angiogenic isoforms of VEGF in health and disease. *Biochem Soc Trans*, 2009, 37(Pt 6):1207–1213.

Rah, D. K., Min, H. J., Kim, Y. W., & Cheon, Y. W. (2017). Effect of Platelet-Rich Plasma on Ischemia-Reperfusion Injury in a Skin Flap Mouse Model. *International Journal of Medical Sciences*, 14(9), 829–839. doi:10.7150/ijms.19573

- Raposo E, et al., 2016. Adipose-derived Stem Cells Added to Platelet-Rich Plasma for Chronic Skin Ulcer Therapy. Parma, Italy. *Wound Research Journal* Vol. 28: pp.434-442.
- Rigotti G, Marchi A., 2009. Adipose-derived Mesenchymal Stem Cells: Past, Present and Future. Verona, Italy. *Aesthetic Plastic Surgery Journal* (33): pp. 271–273.
- Risau W. Mechanisms of angiogenesis. *Nature*, 1997, 386(6626):671–674.
- Rosen LS. Clinical experience with angiogenesis signaling inhibitors: focus on vascular endothelial growth factor (VEGF) blockers. *Cancer Control*, 2002, 9(2 Suppl):36–44.
- Rosen P. Invasif Ductal Carcinoma Assesment of Prognosis, Morphologic Prognostic Markers and Tumor Growth Rate. In : Text book of Rosen’s Breast Pathology, 3rd Ed. 2009; p: 369-79.
- Rumalia VK, Borah GL: Cytokine, growth factor and plastic surgery. *Plast Reconstr Surg* 108:719-733, 2001;
- Schwartz SI (ed): Principles of surgery, ed 7, Newyork, 1999, McGraw-Hill, p269;
- Senger DR, Galli SJ, Dvorak AM, Perruzzi CA, Harvey VS, Dvorak HF. Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid. *Science*, 1983, 219(4587):983–985.
- Sethi A, Kaur T, Malhotra S K, Gambhir M L. Moisturizers: The slippery road. *Indian J Dermatol* 2016;61:279-87
- Shibuya, M., Yamaguchi, S., Yamane, A. et al. (1990) Nucleotide sequence and expression of a novel human receptor-type tyrosine kinase gene (flt) closely related to the fms family. *Oncogene* 5, 519–524
- Shin, Y. J.; Choi, J. S.; et al. (2010). "Induction of vascular endothelial growth factor receptor-3 mRNA in glial cells following focal cerebral ischemia in rats". *J Neuroimmunol.* 229 (1–2): 81–90

- Sidhu SS, Kossiakoff AA. Exploring and Designing Protein Function with Restricted Diversity. *Curr. Opin. Chem. Biol.* 2007; 11; p: 347-54.
- Singh, V. K., Saini, A., Kalsan, M., Kumar, N., & Chandra, R. (2016). Describing the Stem Cell Potency: The Various Methods of Functional Assessment and In silico Diagnostics. *Frontiers in Cell and Developmental Biology*, 4. doi:10.3389/fcell.2016.00134
- Sjamsuhidajat, de Jong., 2016. *Buku Ajar Ilmu Bedah, Sistem Organ dan Tindak Bedahnya*. edisi 4 Vol. 2. Jakarta, Indonesia. EGC.
- Soker, S., Takashima, S., Miao, H., Neufeld, G. & Klagsbrun, M. Neuropilin-1 is expressed by endothelial and tumor cells as an isoform-specific receptor for vascular endothelial growth factor. *Cell* 92, 735–745 (1998).
- Stessuk, T., Puzzi, M. B., Chaim, E. A., Alves, P. C. M., de Paula, E. V., Forte, A., Ribeiro-Paes, J. T. (2016). Platelet-rich plasma (PRP) and adipose-derived mesenchymal stem cells: stimulatory effects on proliferation and migration of fibroblasts and keratinocytes in vitro. *Archives of Dermatological Research*, 308(7), 511–520.
- Tajima, S., Tobita, M., Orbay, H., Hyakusoku, H., & Mizuno, H. (2015). Direct and Indirect Effects of a Combination of Adipose-Derived Stem Cells and Platelet-Rich Plasma on Bone Regeneration. *Tissue Engineering Part A*, 21(5-6), 895–905. doi:10.1089/ten.tea.2014.0336.
- Takahashi, H., & Shibuya, M. (2005). The vascular endothelial growth factor (VEGF)/VEGF receptor system and its role under physiological and pathological conditions. *Clinical Science*, 109(3), 227–241.
- Tantuway V, et al., Autologus grafting of non manipulated freshly isolated-adipose tissue derived stromal vascular fraction in single surgical sitting for treatment of knee osteoarthritis. *Int J Res Orthop.* 2017 Jan;3(1):107-115

- Tavares Pereira, D. dos S., Lima-Ribeiro, M. H. M., de Pontes-Filho, N. T., Carneiro-Leão, A. M. dos A., & Correia, M. T. dos S. (2012). Development of Animal Model for Studying Deep Second-Degree Thermal Burns. *Journal of Biomedicine and Biotechnology*, 2012, 1–7.
- Terman, B. I., Carrion, M. E., Kovacs, E., Rasmussen, B. A., Eddy, R. L. and Shows, T. B. (1991) Identification of a new endothelial cell growth factor receptor tyrosine kinase. *Oncogene* 6, 1677–1683
- Thorne, et al., 2007. *Grabb and Smith Plastic Surgery*, 6th edition. New York, USA. Lippincot Williams and Wilkins.
- Tjwa M, Luttun A, Autiero M, Carmeliet P. VEGF and PlGF: two pleiotropic growth factors with distinct roles in development and homeostasis. *Cell Tissue Res*, 2003, 314(1):5–14.
- Tohidnezhad, M., Varoga, D., Wruck, C.J., et al., 2011. Platelet-released growth factors can accelerate tenocyte proliferation and activate the anti-oxidant response element. *Histochemistry and Cell Biology* 135(5):453-460.
- Van Pham, P., Bui, K. H.-T., Ngo, D., Vu, N., Truong, N., Phan, N. L.-C., Phan, N. (2013). Activated platelet-rich plasma improves adipose-derived stem cell transplantation efficiency in injured articular cartilage. *Stem Cell Research & Therapy*, 4(4), 91.
- Widjajakusumah, Tanzil A., 2014. *Guyton and Hall Buku Ajar Fisiologi Kedokteran*, edisi 12. Singapore. Elsevier.
- Williams N, et al., 2013. *Bailey and Love's Short Practice of Surgery*, 26th
- Williams, C., Assessment and management of paediatric burn injuries. *Nurs Stand*, 2011. 25(25): p. 60-4, 66, 68.
- Witte MB, Barbul A: General principles of wound healing. *Surg Clin North Am* 77:509–528, 1997.

- Yamazaki Y, Morita T. Molecular and functional diversity of vascular endothelial growth factors. *Mol Divers*, 2006, 10(4): 515–527.
- Yancopoulos, G. D., Davis, S., Gale, N. W., Rudge, J. S., Wiegand, S. J., & Holash, J. (2000). Vascular-specific growth factors and blood vessel formation. *Nature*, 407(6801), 242–248. doi:10.1038/35025215
- Zhang YS, He JH, Xiao GY, Li QM: Effect of platelet-rich plasma on the proliferation and adipogenic differentiation of human adipose-derived stem cells in vitro. *Nan Fang Yi Ke Da Xue Xue Bao* 2011, 31:525–528.
- Zuk PA, Zhu M, Ashjian P, De Ugarte DA, Huang JI, Mizuno H, et al. Human adipose tissue is a source of multi-potent stem cells. *Mol Biol Cell*. 2002; 13:4279–95. [PubMed: 12475952]





REKOMENDASI PERSETUJUAN ETIK

Nomor : 216/UN4.6.4.5.31/ PP36/ 2020

Tanggal: 5 Maret 2020

Dengan ini Menyatakan bahwa Protokol dan Dokumen yang Berhubungan Dengan Protokol berikut ini telah mendapatkan Persetujuan Etik :

No Protokol	UH19121108	No Sponsor Protokol	
Peneliti Utama	dr. Sartian Battung	Sponsor	
Judul Peneliti	Efek Penambahan Platelet-Rich Plasma (PRP) Dan Stromal Vascular Fraction (SVFs) Terhadap Kadar VEGF Serum Pada Penyembuhan Luka Bakar Deep Dermal Wistar		
No Versi Protokol	1	Tanggal Versi	16 Desember 2020
No Versi PSP		Tanggal Versi	
Tempat Penelitian	Laboratorium Hewan FKUH, RSUH Laboratorium HUMAN-RC dan Laboratorium PA Cerekang Makassar		
Jenis Review	<input type="checkbox"/> Exempted <input checked="" type="checkbox"/> Expedited <input type="checkbox"/> Fullboard Tanggal	Masa Berlaku 5 Maret 2020 sampai 5 Maret 2021	Frekuensi review lanjutan
Ketua Komisi Etik Penelitian Kesehatan FKUH	Nama Prof.Dr.dr. Suryani As'ad, M.Sc.,Sp.GK (K)	Tanda tangan 	
Sekretaris Komisi Etik Penelitian Kesehatan FKUH	Nama dr. Agussalim Bukhari, M.Med.,Ph.D.,Sp.GK (K)	Tanda tangan 	

Kewajiban Peneliti Utama:

- Menyerahkan Amandemen Protokol untuk persetujuan sebelum di implementasikan
- Menyerahkan Laporan SAE ke Komisi Etik dalam 24 Jam dan dilengkapi dalam 7 hari dan Laporan SUSAR dalam 72 Jam setelah Peneliti Utama menerima laporan
- Menyerahkan Laporan Kemajuan (progress report) setiap 6 bulan untuk penelitian resiko tinggi dan setiap setahun untuk penelitian resiko rendah
- Menyerahkan laporan akhir setelah Penelitian berakhir
- Melaporkan penyimpangan dari protokol yang disetujui (protocol deviation / violation)
- Mematuhi semua peraturan yang ditentukan