

DAFTAR PUSTAKA

- A.L. de Oliveira Manoel, A.C. Neto, P.V. Veigas, S. Rizoli Traumatic brain injury associated coagulopathy Neurocrit Care, 22 (1) (2015), pp. 34-44
- Ahmad. M.R, Hanna, Effect of equiosmolar solutions of hypertonic sodium lactate versus mannitol in craniectomy patients with moderate traumatic brain injury. Med J Indones, Vol. 23, No. 1, February 2014.
<http://dx.doi.org/10.13181/mji.v23i1.686>
- Alexiou GA, Lianos G, Fotakopoulos G, Michos E, Pachatouridis D, dan Voulgaris S. 2014. Admission glucose and coagulopathy occurrence in patients with traumatic brain injury. *Brain Inj* 28(4):439-440
- Algattas H, Huang JH. 2014. Traumatic Brain Injury Pathophysiology and Treatment: Early, Intermediate and Late Phase Post-Injury. International Journal of Molecular Sciences. 15: 309-41.
- Alvis-Miranda HR, Navas-Marrugo SZ, Velasquez-Loperena RA, Adie-Villafaña RJ, Velasquez-Loperena D, Castellar-Leones SM, et al. Effects of Glycemic Level on Outcome of Patients with Traumatic Brain Injury: A Retrospective Cohort Study. Bull Emerg Trauma. 2014;2(2):65-71.
- Amri I. 2017. Pengelolaan Peningkatan Tekanan Intrakranial. MEDIKA TADULAKO Jurnal Ilmiah Kedokteran. 4(3): 1-17.

Arifianto MR, Ma'ruf AZ, Ibrahim A, Bajamal AH. Role of hypertonic sodium lactate in traumatic brain injury management. Asian J Neurosurg 2018;13:971-5.

Blumberg, PC. 1987. Pathology in Head Injury. In: *Neurological Skill* 18(1): 58-62

Budi Harto Batubara, Nazaruddin Umar, Chairul M. Mursin. Perbandingan Osmolaritas Plasma Setelah Pemberian Manitol 20% 3 mL/ kgBB dengan Natrium Laktat Hipertonik 3 mL/kgBB pada Pasien Cedera Otak Traumatik Ringan-Sedang JAP, Volume 4 Nomor 3, Desember 2016

Carpenter KHL, Jalloh I, dan Hutchinson PJ. 2015. Glycolysis and the significance of lactate in traumatic brain injury. *Frontiers in Neuroscience* 9:2-3

Cheng-Shyuan Rau 1,†, Shao-Chun Wu 2,†, Yi-Chun Chen 3, Peng-Chen Chien 3, Hsiao-Yun Hsieh 3, Pao-Jen Kuo 3 and Ching-Hua Hsieh 3,* Stress-Induced Hyperglycemia, but Not Diabetic Hyperglycemia, Is Associated with Higher Mortality in Patients with Isolated Moderate and Severe Traumatic Brain Injury: Analysis of a Propensity Score-Matched Population. Int. J. Environ. Res. Public Health 2017, 14, 1340; doi:10.3390/ijerph14111340

Cottenceau, V., Masson, F., Mahamid, E., Petit, L., Shik, V., Sztark, F., ... Soustiel, J. F. (2011). *Comparison of Effects of Equiosmolar Doses of Mannitol and Hypertonic Saline on Cerebral Blood Flow and*

Metabolism in Traumatic Brain Injury. *Journal of Neurotrauma*, 28(10), 2003–2012. doi:10.1089/neu.2011.1929

Elliot C, McKenzie M, Kelly C. 2015. Mannitol dosing error during interfacility transfer for intracranial emergencies. *Neurosurgeon Journal*. 123: 1166-9.

Folkerson, L. E., Sloan, D., Davis, E., Kitagawa, R. S., Cotton, B. A., Holcomb, J. B., ... Wade, C. E. (2018). Coagulopathy as a predictor of mortality after penetrating traumatic brain injury. *The American Journal of Emergency Medicine*, 36(1), 38–42

Fuentes-García, Diego & Hernández-Palazón, Joaquín & Falcon-Araña, Luis & Piqueras-Pérez, Claudio & Doménech-Asensi, Paloma. (2013). Effect of Equiosmolar Solutions of Mannitol Versus Hypertonic Saline on Blood Coagulation Assessed by Thromboelastometry During Elective Craniotomy. *American Society Of Anaesthesiology*.

George A. Alexiou¹, Georgios Lianos², George Fotakopoulos¹, Evangelos Michos¹, Dimitrios Pachatouridis¹, & Spyridon Vouglaris¹. Admission glucose and coagulopathy occurrence in patients with traumatic brain injury. *Brain Inj*, 2014; 28(4): 438–441. DOI: 10.3109/02699052.2014.888769

Haddad and Arabi: Critical care management of severe traumatic brain injury in adults. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine* 2012 20:12.

Haddad and Arabi: Critical care management of severe traumatic brain injury in adults. Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine 2012 20:12.

Horng S, Therattil A et al. 2017. Astrocytic tight junction control inflammatory CNS lesion pathogenesis. J Clin Invest. 127(8): 3136-51.

Ichai, C., Armando, G., Orban, J.-C., Berthier, F., Rami, L., Samat-Long, C., ... Leverve, X. (2008). *Sodium lactate versus mannitol in the treatment of intracranial hypertensive episodes in severe traumatic brain-injured patients*. Intensive Care Medicine, 35(3), 471–479. doi:10.1007/s00134-008-1283-5

Iwan AR, Sri R, et all. 2015. Terapi Hiperosmolar pada Cedera Otak Traumatika. JNI; 4 (2): 119–33

Jaya I, Widodo D, Ganda I.J. *Comparison of Giving Hipertonic Saline Solution 3% and Manitol 20% on Conservative Moderate Head Injury Patients* JST Kesehatan, Oktober 2017, Vol. 7 No. 4 : 374 – 380

Jeroen Hermanides^{1,5*}, Mark P. Plummer², Mark Finnis³, Adam M. Deane⁴, Jonathan P. Coles¹ and David K. Menon¹ Glycaemic control targets after traumatic brain injury: a systematic review and meta-analysis. Critical Care (2018) 22:11. DOI 10.1186/s13054-017-1883-y

- Jha, R. M., Kochanek, P. M., & Simard, J. M. (2018). Pathophysiology and treatment of cerebral edema in traumatic brain injury. *Neuropharmacology*. doi:10.1016/j.neuropharm.2018.08.004
- Kafaki, S. B., Alaeddini, K., Qorbani, A., Asadian, L., & Haddadi, K. (2016). *Hyperglycemia: A Predictor of Death in Severe Head Injury Patients. Clinical Medicine Insights: Endocrinology and Diabetes*, 9, CMED.S40330. doi:10.4137/cmed.s40330
- Koenig MA. 2018. Cerebral edema and Elevated Intracranial Pressure. *Minneap Minn.* 24(6): 1588-1602
- Komarova YA, Kruse K, Mehta D. 2017. Protein Interaction at Endothelial Junctions and Signaling Mechanism Regulating Endothelial Permeability. *Circ Res.* 120(1): 179-206.
- Kumar V , Shende N , Singh A , Sahu A. 2015. Blood Glucose Levels and Outcome in Traumatic Brain Injury: An Observational Study. *International Journal of Science and Research (IJSR)* Vol. 4; 8.
- Li J, Wang B. 2013. Hyperosmolar therapy for intracranial pressure in neurological practice manitol versus hypertonic saline *International journal of Anesthesiologi Reseach* 1:56-61
- Li, M., Chen, T., Chen, S., Cai, J., & Hu, Y. (2015). Comparison of Equimolar Doses of Mannitol and Hypertonic Saline for the Treatment of Elevated Intracranial Pressure After Traumatic Brain Injury. *Medicine*, 94(17), e668. doi:10.1097/md.0000000000000668

- Liu J, dan Tian HL. 2016. Relationship between trauma-induced coagulopathy and progressive hemorrhagic injury in patients with traumatic brain injury. *Chinese Journal of Traumatology* 19:173-4
- Liu-DeRyke X, Collingridge DS, Orme J, Roller D, Zurasky J, Rhoney DH. Clinical impact of early hyperglycemia during acute phase of traumatic brain injury. *Neurocrit Care*. 2009;11:151–7.
- Luostarinens, T., Niiya, T., Schramko, A., Rosenberg, P., & Niemi, T. (2010). *Comparison of Hypertonic Saline and Mannitol on Whole Blood Coagulation In Vitro Assessed by Thromboelastometry*. *Neurocritical Care*, 14(2), 238–243. doi:10.1007/s12028-010-9475-6
- Maguigan KL et al. Method of Hypertonic Saline Administration: Effects on Osmolality in Traumatic Brain Injury Patients. *J Clin Neurosci* (2017),
- Malik, Z., Mir, S., & Naqash, I Sofi, K., Wani, A., (2014). *A prospective, randomized, double blind study to compare the effects of equiosmolar solutions of 3% hypertonic saline and 20% mannitol on reduction of brain-bulk during elective craniotomy for supratentorial brain tumor resection*. *Anesthesia: Essays and Researches*, 8(3), 388. doi:10.4103/0259-1162.143155
- Mikkelsen, M., et al. 2012. The Acute Coagulopathy of Trauma: Mechanisms and Tools for Risk Stratification. *Shock Society*. *Shock* 38(35):2-3
- Miranda H, Marrugo S, Loperena RA, Villafane RJ, Loperena DV, Leones SM, et. al. 2014. Effects of Glycemic Level on Outcome of Patients

- with Traumatic Brain Injury: A Retrospective Cohort Study. *Bull Emerg Trauma* 2(2):65-71
- Mustafa AG, Alshboul GA. 2013. Pathophysiology of Traumatic Brain Injury. *Neurosciences Journal*. 18(3): 222-34.
- Oliveira M. Neto AC. 2015. Traumatic Brain Injury Associated Coagulopathy. *Neuro Critical Care Journal*. 22:34-44.
- Oliveira DV, Amorim RL, Viera RC, Paiva WS. 2017. Traumatic Brain Injury and Hyperglycemia. *Impact Journal Oncotarget*. 8(12): 18622.
- Palmaers T, Krämer E, Hinsenkamp J, Eismann H, Scheinichen D, Frank P. Mannitol and the Combination of Mannitol and Gelatin Impair Whole Blood Coagulation and the Platelet Function In Vitro. *Turk J Anaesthesiol Reanim* 2019; 47(3): 199-205.
- Patil, H., & Gupta, R. (2019). "A comparative study of bolus dose of hypertonic saline, mannitol and mannitol plus glycerol combination in patients with severe traumatic brain injury." *World Neurosurgery*.
- Peters N, Farrel L, Smith J. 2018. Hyperosmolar therapy for the treatment of cerebral edema. *US Pharm*. 43(1):8-11.
- Piper, B. J., & Harrigan, P. W. (2015). *Hypertonic Saline in Paediatric Traumatic Brain Injury: A Review of Nine Years' Experience with 23.4% Hypertonic Saline as Standard Hyperosmolar Therapy*. *Anaesthesia and Intensive Care*, 43(2), 204–210. doi:10.1177/0310057x1504300210 Sofi,

Rani P, Panda NB, Hazarika A, Ahluwalia J, Chhabra R. Trauma-induced coagulopathy: Incidence and outcome in patients with isolated traumatic brain injury in a level I trauma care center in India. Asian J Neurosurg 2019;14:1175-80

Rozet I, Tontisirin N, Muangman S, Vavilala MS, Souter MJ, Lee LA, dkk. Effect of equiosmolar solutions of mannitol versus hypertonic saline on intraoperative brain relaxation and electrolyte balance. Anesthesiology. 2007;107(5):697–704.

Sakellaridis, N., Pavlou, E., Karatzas, S., Chroni, D., Vlachos, K., Chatzopoulos, K., ... Karaouli, V. (2011). *Comparison of mannitol and hypertonic saline in the treatment of severe brain injuries*. Journal of Neurosurgery, 114(2), 545–548. doi:10.3171/2010.5.jns091685

Sekeon SAS, Maja PS, Kembuan M. Executive function disorderin acute traumatic brain injury in Manado, Indonesia. Global Journal of Medicine and Public Health. 2015: 4(2). ISSN 2277-9604Shawkat H, Westwood M, Mortimer A. 2012. Mannitol: A review of its clinical uses : Continuing Education in Anasthesia, Critial Care and Pain Journal.12;2:1-4

Seo, W., & Oh, H. (2010). *Alterations in Serum Osmolality, Sodium, and Potassium Levels After Repeated Mannitol Administration*. Journal of Neuroscience Nursing, 42(4), 201–207. doi:10.1097/jnn.0b013e3181e26b4a

- Sharma, R., Setlur, R., & Swamy, M. (2011). *Evaluation of mannitol as an osmotherapeutic agent in traumatic brain injuries by measuring serum osmolality*. *Medical Journal Armed Forces India*, 67(3), 230–233. doi:10.1016/s0377-1237(11)60047-6
- Shawkat, H., Westwood, M.-M., & Mortimer, A. (2012). *Mannitol: a review of its clinical uses*. *Continuing Education in Anaesthesia Critical Care & Pain*, 12(2), 82–85. doi:10.1093/bjaceaccp/mkr063
- Shi J, Dong B, Mao Y, Guan W, et. al. 2016. Review: Traumatic brain injury and hyperglycemia, a potentially modifiable risk factor. *Oncotarget* 7(43):71052.
- Shrestha A, Joshi RM, dan Devkota UP. 2015. Coagulation disorder in moderate to severe traumatic brain injuries. *Asian Journal of Medical Sciences* 6(3):23-24
- Sorani, M. D., Morabito, D., Rosenthal, G., Giacomini, K. M., & Manley, G. T. (2008). *Characterizing the Dose-Response Relationship between Mannitol and Intracranial Pressure in Traumatic Brain Injury Patients Using a High-Frequency Physiological Data Collection System*. *Journal of Neurotrauma*, 25(4), 291–298. doi:10.1089/neu.2007.0411
- Stovell MG, Yan JL, Sleigh A, Mada OM, Carpenter TA, Hutchinson PJA, et al. 2017. Assessing Metabolism and Injury in Acute Human Traumatic Brain Injury with Magnetic Resonance Spectroscopy: Current and Future Applications. *Frontiers in Neurology* 8(1):9

Tamer Abdallah Helmy, Dina Hassan Zidan and Ahmed Mohamed Abdallah. (2016). Hypertonic saline versus mannitol in treatment of diffuse brain edema in patients with moderate and severe traumatic brain injury. *Biolife*, 4(3), pp 471-474 doi:10.17812/blj.2016.4308

Terzioglu, B., Ekinci, O. and Berkman, Z., 2015. Hyperglycemia is a predictor of prognosis in traumatic brain injury: Tertiary intensive care unit study. *Journal of Research in Medical Sciences*, 20(12), p.1166.

van Gent, J.A.N., van Essen, T.A., Bos, M.H.A. et al. Coagulopathy after hemorrhagic traumatic brain injury, an observational study of the incidence and prognosis. *Acta Neurochir* 162, 329–336 (2020).
<https://doi.org/10.1007/s00701-019-04111-z>

Vos PE, Arrastia RD. Traumatic Brain Injury. 1st ed. Oxford: Wiley Blackwell. 2015. Pp 3.

Wang, H., Cao, H., Zhang, X., Ge, L., & Bie, L. (2017). *The effect of hypertonic saline and mannitol on coagulation in moderate traumatic brain injury patients. The American Journal of Emergency Medicine*, 35(10), 1404–1407. doi:10.1016/j.ajem.2017.04.020

Wei HJ, Liu L, Chen FL, Wang D, Wang ZG, Jiang RC, et. al. 2019. Decreased numbers of circulating endothelial progenitor cells are associated with hyperglycemia in patients with traumatic brain injury. *Neural Regen Res* 14(6):985-88

Widodo. D, Islam.A.A, Bahar. B. 2012. Dinamika Kadar Glukosa, Laktat dan Glial Fibrillary Acidic Protein Serum sebagai Prediktor Luaran

Pasien Cedera Otak Tertutup. *Jurnal Berkala Ilmiah Arsip Ilmu Bedah* 4(3)::188-91

Winkler EA et al. Cerebral Edema in Traumatic Brain Injury Neurosurgery Clinics of North America. Elsevier Inc. 2015. 27(4): 473-88.

Winkler, E. A., Minter, D., Yue, J. K., & Manley, G. T. (2016). Cerebral Edema in Traumatic Brain Injury. *Neurosurgery Clinics of North America*, 27(4), 473–488. doi:10.1016/j.nec.2016.05.008

Xue, B., Ruan, S., Xie, P., Yan, K., Gu, Z., Meng, N., ... Zhang, H. (2018). Prospective analysis of glycemic variability in patients with severe traumatic brain injury: modified Leuven's adjustment process versus conventional adjustment process. *Journal of International Medical Research*, 46(4), 1505–1516. doi:10.1177/0300060517738396

Yuan et al. 2018. Prognostic Value of Coagulation tests for in hospital mortality in patients with traumatic brain injury. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine*. 26:3

LAMPIRAN



KEMENTERIAN PENDIDIKAN DAN KEBUDAYAAN
 UNIVERSITAS HASANUDDIN FAKULTAS KEDOKTERAN
 KOMITE ETIK PENELITIAN KESEHATAN
 RSPTN UNIVERSITAS HASANUDDIN
 RSUP Dr. WAHIDIN SUDIROHUSODO MAKASSAR



Sekeerariat : Lantai 3 Gedung Laboratorium Terpadu
 Jl. PERINTIS KEMERDEKAAN KAMPUS TAMALANREA KM.10 MAKASSAR 90245.
 Contact Person: dr. Agussalim Bukhari,M.Med,PhD, Sp.GK TELP. 081241850858, 0411 5780103, Fax: 0411-581431

REKOMENDASI PERSETUJUAN ETIK

Nomor : 359/UN4.6.4.5.31/PP36/2020

Dengan Ini Menyatakan bahwa Protokol dan Dokumen yang Berhubungan Dengan Protokol berikut ini telah mendapatkan Persetujuan Etik :

Tanggal: 8 Juli 2020

No Protokol	UH20010041	No Sponsor Protokol	
Peneliti Utama	dr Martin Saputro Wardoyo	Sponsor	
Judul Peneliti	Hubungan Manitol 20% Dengan Dosis Berbeda Terhadap Osmolaritas Darah,Gula Darah Serum dan Faktor Koagulasi Pasien Cedera Otak Sedang Dengan Peningkatan Tekanan Intrakranial		
No Versi Protokol	2	Tanggal Versi	3 Juli 2020
No Versi PSP	2	Tanggal Versi	3 Juli 2020
Tempat Penelitian	RSUP Dr Wahidin Sudirohusodo dan RS Jejaring		
Jenis Review	<input type="checkbox"/> Exempted <input type="checkbox"/> Expedited <input checked="" type="checkbox"/> Fullboard 5 Februari 2020	Masa Berlaku 8 Juli 2020 Sampai 8 Juli 2021	Frekuensi review lanjutan
Ketua Komisi Etik Penelitian Kesehatan FK UH	Nama Prof.Dr.dr.Suryani As'ad.,MSc,Sp.GK (K)	Tanda tangan	Tanggal
Sekretaris Komisi Etik Penelitian Kesehatan FK UH	Nama dr. Agussalim Bukhari,M.Med,PhD,Sp.GK (K)	Tanda tangan	Tanggal

Kewajiban Peneliti Utama:

- Menyerahkan Amandemen Protokol untuk persetujuan sebelum di implementasikan
- Menyerahkan Laporan SAE ke Komisi Etik dalam 24 Jam dan dilengkapi dalam 7 hari dan Lapor SUSAR dalam 72 Jam setelah Peneliti Utama menerima laporan
- Menyerahkan Laporan Kemajuan (progress report) setiap 6 bulan untuk penelitian resiko tinggi dan setiap setahun untuk penelitian resiko rendah
- Menyerahkan laporan akhir setelah Penelitian berakhir
- Melaporkan penyimpangan dari protokol yang disetujui (protocol deviation / violation)
- Mematuhi semua peraturan yang ditentukan