

## DAFTAR PUSTAKA

Agre, P. (2006) 'The aquaporin water channels', *Proceedings of the American Thoracic Society*, 3(1), pp. 5–13. doi: 10.1513/pats.200510-109JH.

Ågren, J. *et al.* (2003) 'Transepidermal water loss in developing rats: Role of aquaporins in the immature skin', *Pediatric Research*, 53(4), pp. 558–565. doi: 10.1203/01.PDR.0000055777.25933.98.

Aisyah, I., Zulkarnain, I. and Sawitri, S. (2019) 'Profil Nilai pH dan Transepidermal Water Loss (TEWL) Pada Pasien Dermatitis Atopik Anak', *Berkala Ilmu Kesehatan Kulit dan Kelamin*, 31(2), pp. 138–143.

Aoki, M. *et al.* (2019) 'Comparison of ceramide retention in the stratum corneum between dry skin and normal skin using animal model with fluorescent imaging method', *Skin Research and Technology*, 25(2), pp. 158–164. doi: 10.1111/srt.12625.

Aydin, S. (2015) 'A short history, principles, and types of ELISA, and our laboratory experience with peptide/protein analyses using ELISA', *Peptides*, 72, pp. 4–15. doi: 10.1016/j.peptides.2015.04.012.

Azad, A. K. *et al.* (2021) 'Human Aquaporins: Functional Diversity and Potential Roles in Infectious and Non-infectious Diseases', *Frontiers in Genetics*, 12(March). doi: 10.3389/fgene.2021.654865.

Bock, M., Wulfhorst, B. and John, S. M. (2007) 'Site variations in susceptibility to SLS', *Contact Dermatitis*, 57(2), pp. 94–96. doi: 10.1111/j.1600-0536.2007.01159.x.

Bollag, W. B. *et al.* (2020) 'Aquaporin-3 in the epidermis: More than skin deep', *American Journal of Physiology - Cell Physiology*, 318(6), pp. C1144–C1153. doi: 10.1152/ajpcell.00075.2020.

Cleary, G. (2013) 'Transdermal DDS', *Biomaterials Science: An Introduction to Materials: Third Edition*, pp. 1073–1082. doi: 10.1016/B978-0-08-087780-8.00101-7.

Coderch, L. *et al.* (2003) 'Ceramides and skin function', *American Journal of Clinical Dermatology*, 4(2), pp. 107–129. doi: 10.2165/00128071-200304020-00004.

D.C. Soler, Ph.D. eD.C. Soler, Ph.D., X. Bai, Ph.D., L. Ortega, T. Pethukova, M.D., S.T. Nedorost, M.D.&, D.L. Popkin, MD/PhD, &, K. D. Cooper, M.D.&, and T.S. McCormick, P. . (2011) 'The Key Role of Aquaporin 3 and Aquaporin 10 in the Pathogenesis of Pompholyx', *Physiology & behavior*, 176(1), pp. 139–148. doi: 10.1016/j.mehy.2015.02.006.The.

Darlenski, R., Kazandjieva, J. and Tsankov, N. (2011) 'Skin barrier function: morphological basis and regulatory mechanisms', *Journal of Clinical Medicine*, 4(1), pp. 36–45.

Darmawan, H. and Wijayadi, L. Y. (2015) 'Aktivasi Protein Aquaporin-3 Pada Keratinosit Kulit Manusiasebagai Target Pengobatan Hidrasi Kulit Oleh Asiaticosida Yang diisolasi Dari Centella Asiatica', *Journal Untar*. Available at: <https://www.scribd.com/doc/292075231/Aktivasi-Protein-Aquaporin>.

Davis, D., Kannan, M. and Wattenberg, B. (2018) 'Orm/ORMDL proteins: Gate guardians and master regulators', *Advances in Biological Regulation*, 70, pp. 3–18. doi: 10.1016/j.jbior.2018.08.002.

Day, R. E. *et al.* (2014) 'Human aquaporins: Regulators of transcellular water flow', *Biochimica et Biophysica Acta - General Subjects*, 1840(5), pp. 1492–1506. doi: 10.1016/j.bbagen.2013.09.033.

Draelos, Z. D. (2008) 'The effect of ceramide-containing skin care products on eczema resolution duration', *Cutis*, 81(1), pp. 87–91.

Engvall, E. (2010) 'The ELISA, enzyme-linked immunosorbent assay', *Clinical Chemistry*, 56(2), pp. 319–320. doi: 10.1373/clinchem.2009.127803.

Farwanah, H. *et al.* (2005) 'Ceramide profiles of the uninvolved skin in atopic dermatitis and psoriasis are comparable to those of healthy skin', *Archives of Dermatological Research*, 296(11), pp. 514–521. doi: 10.1007/s00403-005-0551-2.

Fujimura, T. *et al.* (2017) 'Differences in the stratum corneum of Indonesian infants and adults', 57(1), pp. 35–40. doi: 10.14238/pi51.1.2011.35-40.

Giusti, F. *et al.* (2001) 'Skin barrier, hydration, and pH of the skin of infants under 2 years of age', *Pediatric Dermatology*, 18(2), pp. 93–96. doi: 10.1046/j.1525-1470.2001.018002093.x.

Gorzelanny, C. *et al.* (2020) 'Skin barriers in dermal drug delivery: Which barriers have to be overcome and how can we measure them?', *Pharmaceutics*, 12(7), pp. 1–31. doi: 10.3390/pharmaceutics12070684.

Hannun, Y. A. and Obeid, L. M. (2008) 'Principles of bioactive lipid signalling: Lessons from sphingolipids', *Nature Reviews Molecular Cell Biology*, 9(2), pp. 139–150. doi: 10.1038/nrm2329.

Hans Schaefer; Thomas E. Redelmeier (1996) *Skin Barrier*. Switzerland: karger.

Hara-Chikuma, M. and Verkman, A. S. (2008) 'Roles of aquaporin-3 in the epidermis', *Journal of Investigative Dermatology*, 128(9), pp. 2145–2151. doi: 10.1038/jid.2008.70.

Hara, M., Ma, T. and Verkman, A. S. (2002) 'Selectively reduced glycerol in skin of aquaporin-3-deficient mice may account for impaired skin hydration, elasticity, and barrier recovery', *Journal of Biological Chemistry*, 277(48), pp. 46616–46621. doi: 10.1074/jbc.M209003200.

Hayati, L. *et al.* (2015) 'Perbandingan Efektivitas Krim Urea 10% dan Krim Niasinamid 4% Terhadap Hidrasi Kulit Pasien Dermatitis Atopik', *Majalah Kedokteran Sriwijaya*, 47(01), pp. 2–5.

Hnasko, R. (2015) 'ELISA: Methods and Protocols', *ELISA: Methods and Protocols*, 1318, pp. 1–216. doi: 10.1007/978-1-4939-2742-5.

Hoeger, P. H. *et al.* (2002) 'Epidermal barrier lipids in human vernix caseosa: Corresponding ceramide pattern in vernix and fetal skin', *British Journal of Dermatology*, 146(2), pp. 194–201. doi: 10.1046/j.1365-2133.2002.04584.x.

Jang, H. H. and Lee, S. N. (2016) 'Epidermal Skin Barrier', *Asian Journal of Beauty and Cosmetology*, 14(3), pp. 339–347. doi: 10.20402/ajbc.2016.0039.

Janssens, M. *et al.* (2012) 'Increase in short-chain ceramides correlates with an altered lipid organization and decreased barrier function in atopic eczema patients', *Journal of Lipid Research*, 53(12), pp. 2755–2766. doi: 10.1194/jlr.P030338.

Kahraman, E. *et al.* (2019) 'Recent advances on topical application of ceramides to restore barrier function of skin', *Cosmetics*, 6(3). doi: 10.3390/COSMETICS6030052.

Keles, F. F., Pandaleke, H. E. J. and Mawu, F. O. (2016) 'Profil dermatitis atopik pada anak di Poliklinik Kulit dan Kelamin RSUP Prof. Dr. R. D. Kandou Manado periode Januari 2013 – Desember 2015', *e-CliniC*, 4(2). doi: 10.35790/ecl.4.2.2016.14456.

Kelleher, M. *et al.* (2015) 'Skin barrier dysfunction measured by transepidermal water loss at 2 days and 2 months predates and predicts atopic dermatitis at 1 year', *Journal of Allergy and Clinical Immunology*, 135(4), pp. 930-935.e1. doi: 10.1016/j.jaci.2014.12.013.

Keng Wooi Ng and Wing Man Lau (2015) 'Skin Deep: The Basics of Human Skin Structure and Drug Penetration', *Percutaneous Penetration Enhancers Chemical Methods in Penetration Enhancement: Drug Manipulation Strategies and Vehicle Effects*, pp. 1–341. doi: 10.1007/978-3-662-45013-0.

Kottner, J., Lichterfeld, A. and Blume-Peytavi, U. (2013) 'Transepidermal water loss in young and aged healthy humans: A systematic review and meta-analysis', *Archives of Dermatological Research*, 305(4), pp. 315–323. doi: 10.1007/s00403-012-1313-6.

Kurniadi (2012) 'Peran Aquaporin Dalam Pengaturan Transport Air Kurniadi',

*Jurnal Kesehatan Prima*, 6(2), pp. 976–981.

Lawton, S. (2019) ‘Skin 1: the structure and functions of the skin’, *Nursing Times*, 115(12), pp. 30–33.

Lee, Y. *et al.* (2012) ‘Changes in transepidermal water loss and skin hydration according to expression of aquaporin-3 in psoriasis’, *Annals of Dermatology*, 24(2), pp. 168–174. doi: 10.5021/ad.2012.24.2.168.

Liu, H. and Wintour, E. M. (2005) ‘Aquaporins in development - A review’, *Reproductive Biology and Endocrinology*, 3, pp. 1–10. doi: 10.1186/1477-7827-3-18.

Loeffelholz, M. J. *et al.* (2020) ‘Multicenter evaluation of the cepheid xpert xpress sars-cov-2 test’, *Journal of Clinical Microbiology*, 58(8). doi: 10.1128/JCM.00926-20.

Luisa Coderch, Olga López, A. de la M. & J. L. P. (2012) ‘Ceramides and Skin Function’, *American Journal of Clinical Dermatology volume 4, pages107–129(2003)*. Available at: <https://link.springer.com/article/10.2165/00128071-200304020-00004>.

Made, N., Savitri, A. and Ayu, P. R. (2017) ‘Sel punca mesenkimal sebagai terapi dermatitis atopik yang menjanjikan mesenchymal stem cell as promising therapy of atopic dermatitis’, *Fakultas Kedokteran Universitas Lampung*, 7(November), pp. 83–88.

Mandy Alhadj and Aisha Farhana (2020) *Enzyme Linked Immunosorbent Assay, StatPearls Publishing, Treasure Island (FL)*. StatPearls Publishing LLC.

Mathanda, T. R. *et al.* (2016) ‘Transepidermal Water Loss in Neonates: Baseline Values Using a Closed-Chamber System’, *Pediatric Dermatology*, 33(1), pp. 33–37. doi: 10.1111/pde.12704.

Meckfessel, M. H. and Brandt, S. (2014) ‘The structure, function, and importance of ceramides in skin and their use as therapeutic agents in skin-care products’, *Journal of the American Academy of Dermatology*, 71(1), pp. 177–184. doi: 10.1016/j.jaad.2014.01.891.

Meeusen, J. W. *et al.* (2018) ‘Plasma ceramides a novel predictor of major adverse cardiovascular events after coronary angiography’, *Arteriosclerosis, Thrombosis, and Vascular Biology*, 38(8), pp. 1933–1939. doi: 10.1161/ATVBAHA.118.311199.

Minami-Hori, M. *et al.* (2014) ‘Developmental alterations of physical properties and components of neonatal-infantile stratum corneum of upper thighs and diaper-covered buttocks during the 1st year of life’, *Journal of Dermatological Science*, 73(1), pp. 67–73. doi: 10.1016/j.jdermsci.2013.08.015.

Mizutani, Y. *et al.* (2009) ‘Ceramide biosynthesis in keratinocyte and its role in skin function’, *Biochimie*, 91(6), pp. 784–790. doi: 10.1016/j.biochi.2009.04.001.

Moore, D. J. and Rawlings, A. V. (2017) ‘The chemistry, function and (patho)physiology of stratum corneum barrier ceramides’, *International Journal of Cosmetic Science*, 39(4), pp. 366–372. doi: 10.1111/ics.12399.

Nakahigashi, K. *et al.* (2011) ‘Upregulation of aquaporin-3 is involved in keratinocyte proliferation and epidermal hyperplasia’, *Journal of Investigative Dermatology*, 131(4), pp. 865–873. doi: 10.1038/jid.2010.395.

Nareswari, A. *et al.* (2019) ‘Perbedaan Transepidermal Water Loss Pada Neonatus Neonatal Transepidermal Water Loss Difference Based on’, (132), pp. 9–15.

Nikam, V. N. *et al.* (2019) ‘Transepidermal Water Loss in Psoriasis: A Case-control Study Abstract’, *Indian Dermatology Online Journal*. doi: 10.4103/idoj.IDOJ\_180\_18.

Nomoto, K. *et al.* (2015) ‘Increasing effect of ceramides in skin by topical application of sphingosine’, *Global Dermatology*, 2(4), pp. 154–158. doi: 10.15761/god.1000143.

Olsson, M. *et al.* (2006) ‘Increased expression of aquaporin 3 in atopic eczema’, *Allergy: European Journal of Allergy and Clinical Immunology*, 61(9), pp. 1132–1137. doi: 10.1111/j.1398-9995.2006.01151.x.

Park, J. W., Park, W. J. and Futerman, A. H. (2014) ‘Ceramide synthases as potential targets for therapeutic intervention in human diseases’, *Biochimica et Biophysica Acta - Molecular and Cell Biology of Lipids*, 1841(5), pp. 671–681. doi: 10.1016/j.bbalip.2013.08.019.

Partogi, D. (2008) ‘Kulit kering’, *Departemen Ilmu Kesehatan Kulit Dan Kelamin FK.USU*, pp. 1–12.

Pratima, N. A. (2018) ‘Liquid Chromatography-Mass Spectrometry and Its Applications: A Brief Review’, *Archives of Organic and Inorganic Chemical Sciences*, 1(1), pp. 26–34. doi: 10.32474/aoics.2018.01.000103.

Qin, H. *et al.* (2011) ‘Aquaporin-3 in keratinocytes and skin: Its role and interaction with phospholipase D2’, *Archives of Biochemistry and Biophysics*, 508(2), pp. 138–143. doi: 10.1016/j.abb.2011.01.014.

Radityastuti, R. and Anggraeni, P. (2017) ‘Karakteristik Penyakit Kulit Akibat Infeksi di Poliklinik Kulit Dan Kelamin RSUP Dr. Kariadi Semarang Periode Januari 2008 – Desember 2010’, *Media Medika Muda*, 2(2), pp. 137–142.

Raone, B. *et al.* (2014) ‘Transepidermal water loss in newborns within the first 24 hours of life: Baseline values and comparison with adults’, *Pediatric*

*Dermatology*, 31(2), pp. 191–195. doi: 10.1111/pde.12274.

Rohan, H. H. dan S. H. S. (2013) *Kesehatan Reproduksi*. Yogyakarta: Nuha Medika).

Roudier, N. *et al.* (1998) ‘Evidence for the presence of aquaporin-3 in human red blood cells’, *Journal of Biological Chemistry*, 273(14), pp. 8407–8412. doi: 10.1074/jbc.273.14.8407.

Sadick, N. S. and Krueger, N. (2014) *Cosmetic dermatology, Dermatologic Clinics*. doi: 10.1016/j.det.2013.10.001.

Schachner, L. A. *et al.* (2020) ‘A Consensus About the Importance of Ceramide Containing Skincare for Normal and Sensitive Skin Conditions in Neonates and Infants’, *Journal of drugs in dermatology: JDD*, 19(8), pp. 769–776. doi: 10.36849/JDD.2020.5252.

Shimada, K. *et al.* (2009) ‘Increased transepidermal water loss and decreased ceramide content in lesional and non-lesional skin of dogs with atopic dermatitis’, *Veterinary Dermatology*, 20(5–6), pp. 541–546. doi: 10.1111/j.1365-3164.2009.00847.x.

Sinulingga, E. H., Budiastuti, A. and Widodo, A. (2018) ‘Efektivitas Madu Dalam Formulasi Pelembap Pada Kulit Kering’, *Diponegoro Medical Journal (Jurnal Kedokteran Diponegoro)*, 7(1), pp. 146–157.

Siskind, L. J., Mullen, T. D. and Obeid, L. M. (2010) *The role of ceramide in cell regulation*. Second Edi, *Handbook of Cell Signaling*, 2/e. Second Edi. Elsevier Inc. doi: 10.1016/B978-0-12-374145-5.00148-0.

Stamatas, G. N. *et al.* (2010) ‘Infant skin microstructure assessed in vivo differs from adult skin in organization and at the cellular level’, *Pediatric Dermatology*, 27(2), pp. 125–131. doi: 10.1111/j.1525-1470.2009.00973.x.

Szpilbarg, N. *et al.* (2018) ‘New insights into the role of placental aquaporins and the pathogenesis of preeclampsia’, *Frontiers in Physiology*, 9(OCT), pp. 1–7. doi: 10.3389/fphys.2018.01507.

Tachi, M. and Iwamori, M. (2008) ‘Mass spectrometric characterization of cholesterol esters and wax esters in epidermis of fetal, adult and keloidal human skin’, *Experimental Dermatology*, 17(4), pp. 318–323. doi: 10.1111/j.1600-0625.2007.00647.x.

Taïeb, A. (2018) ‘Skin barrier in the neonate’, *Pediatric Dermatology*, 35, pp. s5–s9. doi: 10.1111/pde.13482.

Toncic, R. J. *et al.* (2020) ‘Altered levels of sphingosine, sphinganine and their ceramides in atopic dermatitis are related to skin barrier function, disease severity and local cytokine milieu’, *International Journal of Molecular Sciences*, 21(6), pp. 1–14. doi: 10.3390/ijms21061958.

- Trisnowati, N. and Soebono, H. (2015) 'Hubungan Hidrasi Stratum Korneum, Kehilangan Air Melalui Epidermis dan Kadar Keperahan Dermatitis Atopik', *Media Dermatovenereologica Indonesiana*, 42(4), pp. 164–166.
- Uchida, Y. (2017) *Skin Lipids, Cosmetic Science and Technology: Theoretical Principles and Applications*. Elsevier Inc. doi: 10.1016/B978-0-12-802005-0.00041-0.
- Uchida, Y. and Park, K. (2021) 'Ceramide in Skin Health and Disease: An Update', *American Journal of Clinical Dermatology*, 22(6), pp. 853–866. doi: 10.1007/s40257-021-00619-2.
- Vashist, S. K. and Luong, J. H. T. (2018) *Enzyme-linked immunoassays, Handbook of Immunoassay Technologies: Approaches, Performances, and Applications*. Elsevier Inc. doi: 10.1016/B978-0-12-811762-0.00005-0.
- Vavrova, K., Kovačik, A. and Opalka, L. (2017) 'Ceramide in the skin barrier', *European Pharmaceutical Journal*, 64(2), pp. 28–35. doi: 10.1515/afpuc-2017-0004.
- Verbavatz, J. M. *et al.* (2009) 'Skin aquaporins: Function in hydration, wound healing, and skin epidermis homeostasis', *Handbook of Experimental Pharmacology*, 190, pp. 205–217. doi: 10.1007/978-3-540-79885-9\_10.
- Verkman, A. . (2000) 'Structure and function of aquaporin water channels (Invited review)', *Environmental Pollution*, 138(3) Verkman, A. . and A. K. M. (2005) 'Structure and function of aquaporin water channels (Invited review)', *Environmental Pollution*, 138(3), p. 377. doi: 10.1016/j.envpol.2004.10.021.), p. 377. doi: 10.1016/j.envpol.2004.10.021.
- Verkman, A. S. (2013) 'Aquaporins', *Current Biology*, 23(2), pp. 52–55. doi: 10.1016/j.cub.2012.11.025.
- Wijayadi, L. Y. (2021) 'Aktivitas Ekspresi Protein dan Gen Aquaporin 3 (AQP3) Sebagai Target Pengobatan Hidrasi Kulit', *Jurnal Muara Medika dan Psikologi Klinis*, 1(1), pp. 89–97.
- Yousef H, Alhajj M, S. S. (2020) *Anatomy, Skin (Integument), Epidermis*. StatPearls Publishing LLC.
- Yuan, C. *et al.* (2017) 'Properties of Skin in Chinese Infants: Developmental Changes in Ceramides and in Protein Secondary Structure of the Stratum Corneum', *BioMed Research International*, 2017. doi: 10.1155/2017/3594629.
- Yulianti, L. *et al.* (2015) 'Artikel Asli Ekspresi Protein Aquaporin3 Pada Fibroblas Dan Keratinosit Kulit Manusia Yang Diinduksioleh Nanopartikel Kitosan', (1), pp. 7–17.

Zannetti, A. *et al.* (2020) 'Mesenchymal stem cells', *Role of Aquaporins in the Physiological Functions of Mesenchymal Stem Cells*, pp. 147–156. doi: 10.1007/978-4-431-56059-3\_12.

Zelenina, M., Zelenin, S. and Aperia, A. (2005) 'Water Channels (Aquaporins) and Their Role for Postnatal Adaptation', *Pediatric Research*, 57(5), pp. 47R-53R. doi: 10.1203/01.PDR.0000159572.79074.0B.

## LAMPIRAN

### Lampiran 1. Lembar *Informed Consent*

No. Etik Penelitian : 448/UN4.6.4.5.31/PP36/2021, Tanggal 12 Juli 2021

Judul Penelitian : Peranan Aquaporine-3 dan Ceramide Terhadap SawarKulit Pada Bayi

### **FORMULIR PERSETUJUAN SETELAH PENJELASAN (INFORMED CONSENT)**

Berikut adalah naskah yang akan dibacakan kepada Ibu responden penelitian :

Saya adalah mahasiswa S3 program studi ilmu kedokteran Universitas Hasanuddin yang akan melaksanakan penelitian sebagai salah satu kegiatan dalam menyelesaikan tugas akhir di program studi S3 universitas hasanuddin.

Tujuan penelitian ini adalah untuk mengetahui kadar Aqaporine-3 dan ceramide sebagai penanda sawar kulit pada bayi, sehingga dengan adanya data kadar Aquaporine3 dan Ceramide kedepannya dapat dijadikan salah satu penanda penentuan kondisi sawar kulit sejak dini.

Saya mengharapkan partisipasi Ibu untuk bersedia menjadi koresponden dalam penelitian ini dengan berpartisipasi untuk menjawab beberapa pertanyaan seputar masalah kondisi riwayat penyakit kulit ibu dan bayinya serta bersedia dan sukarela untuk ibu dan bayi di periksa kondisi kulitnya dengan menggunakan Trans Epidermal Water Lost (TEWL).

Pemeriksaan TEWL dilakukan dalam waktu sekitar 30 menit. Pertama-tama calon responden mengkondisikan suhu tubuh dengan suhu ruangan dengan cara duduk sekitar 20 menit di dalam ruang pemeriksaan. Pemeriksaan dilakukan pada lengan dalam bagian bawah. Selanjutnya bagian lengan dalam bagian bawah di bersihkan dengan menggunakan kapas pembersih yang mengandung antiseptik. Selanjutnya alat TEWL di lekatkan/ditempelkan pada bagian lengan dalam bagian bawah yang telah di bersihkan untuk dilakukan pengukuran nilai TEWL, pengukuran berlangsung sekitar 30 detik. Tahapan terakhir adalah pembacaan.

Tanggapan atau jawaban yang Ibu akan kami jaga kerahasiannya dan hanya digunakan untuk kepentingan penelitian ini

Terima kasih atas kerjasama Ibu Setelah mendengar dan memahami penjelasan penelitian ini, dengan ini saya menyatakan

SETUJU/TIDAK SETUJU\*

Untuk ikut sebagai respondent/ sample penelitian

Jakarta, \_\_\_\_\_ 2021

Nama Terang \_\_\_\_\_ :

Alamat \_\_\_\_\_ :

\_\_\_\_\_


**Lampiran 2. Lembar Rekomendasi Persetujuan Etik**

Lampiran 3. Dokumentasi Pengukuran TEWL



Gambar Pemeriksaan TEWL Bayi Umur 9-10 Bulan menggunakan alat Tewameter pada posisi lengan polar selama 30 detik

**Lampiran 4. Dokumentasi Pengukuran Kadar AQP3 dan CER**



Gambar Preparasi Sampel Plasma Arteri umbilikal



Gambar Pengukuran AQP3 dan CER menggunakan alat Elisa Reader