

DAFTAR PUSTAKA

- Abdel-Sayed, P. *et al.* (2019) ‘Cell therapies for skin regeneration: an overview of 40 years of experience in burn units.’, *Swiss medical weekly*. Switzerland, 149, p. w20079. doi: 10.4414/smw.2019.20079.
- Abraham, J. A. and Klagsbrun, M. (1988) ‘Modulation of Wound Repair by Members of the Fibroblast Growth Factor Family’, in *The Molecular and Cellular Biology of Wound Repair*. Boston, MA: Springer US, pp. 195–248. doi: 10.1007/978-1-4899-0185-9_6.
- Aleckovic, M. and Simon, C. (2008) ‘Is teratoma formation in stem cell research a characterization tool or a window to developmental biology?’, *Reproductive biomedicine online*. Netherlands, 17(2), pp. 270–280. doi: 10.1016/s1472-6483(10)60206-4.
- Anderson, J. H., Mandell, S. P. and Gibran, N. S. (2019) ‘Burns’, in Brunnicardi, F. C. *et al.* (eds) *Schwartz’s Principles of Surgery*. Eleventh. New York: McGraw-Hill, pp. 251–270.
- ANZBA (2016) *Emergency Management of Severe Burns (EMSB)*. 18th edn. Albany Creek: THE EDUCATION COMMITTEE OF AUSTRALIA AND NEW ZEALAND BURN ASSOCIATION LTD.
- Baglioni, S. *et al.* (2009) ‘Characterization of human adult stem-cell populations isolated from visceral and subcutaneous adipose tissue.’, *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. United States, 23(10), pp. 3494–3505. doi: 10.1096/fj.08-126946.
- Baird, A. (1997) ‘The Regulation of Basic Fibroblast Growth Factor (FGF-2) Through Limited Bioavailability’, in Ziegler, T. R., Pierce, G. F., and Herndon, D. N. (eds) *Growth Factors and Wound Healing*. New York, NY: Springer New York, pp. 27–36.
- Bakacak, M. *et al.* (2016) ‘Protective Effect of Platelet Rich Plasma on Experimental Ischemia/Reperfusion Injury in Rat Ovary.’, *Gynecologic and obstetric investigation*. Switzerland, 81(3), pp. 225–231. doi: 10.1159/000440617.
- Barret-Nerin, J. (2004) *Principles and Practice of Burn Surgery*. CRC Press. doi: 10.1201/b21634.
- Benson, A., Dickson, W. A. and Boyce, D. E. (2006) ‘ABC of wound healing: Burns’, *BMJ*, 333(Suppl S3), p. 609324. doi: 10.1136/sbmj.0609324.
- Borrione, P. *et al.* (2010) ‘Platelet-rich plasma in muscle healing.’, *American journal of physical medicine & rehabilitation*. United States, 89(10), pp. 854–861. doi: 10.1097/PHM.0b013e3181f1c1c7.

- Bourin, P. *et al.* (2013) 'Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International Society for Cell Therapy (ISCT)', *Cytotherapy*. England, 15(6), pp. 641–648. doi: 10.1016/j.jcyt.2013.02.006.
- Braund, R., Hook, S. and Medlicott, N. J. (2007) 'The role of topical growth factors in chronic wounds.', *Current drug delivery*. United Arab Emirates, 4(3), pp. 195–204. doi: 10.2174/156720107781023857.
- Cerqueira, M. T., Pirraco, R. P. and Marques, A. P. (2016) 'Stem Cells in Skin Wound Healing: Are We There Yet?', *Advances in Wound Care*, 5(4), pp. 164–175. doi: 10.1089/wound.2014.0607.
- Cervelli, V. *et al.* (2010) 'Tissue regeneration in loss of substance on the lower limbs through use of platelet-rich plasma, stem cells from adipose tissue, and hyaluronic acid.', *Advances in skin & wound care*. United States, 23(6), pp. 262–272. doi: 10.1097/01.ASW.0000363551.82058.36.
- Chieriegato, K. *et al.* (2011) 'Epidermal growth factor, basic fibroblast growth factor and platelet-derived growth factor-bb can substitute for fetal bovine serum and compete with human platelet-rich plasma in the ex vivo expansion of mesenchymal stromal cells derived from adipose tissue', *Cytotherapy*. England, 13(8), pp. 933–943. doi: 10.3109/14653249.2011.583232.
- Choi, J., Minn, K. W. and Chang, H. (2012) 'The efficacy and safety of platelet-rich plasma and adipose-derived stem cells: an update', *Archives of plastic surgery*. 2012/11/14. The Korean Society of Plastic and Reconstructive Surgeons, 39(6), pp. 585–592. doi: 10.5999/aps.2012.39.6.585.
- Comella, K., Silbert, R. and Parlo, M. (2017) 'Effects of the intradiscal implantation of stromal vascular fraction plus platelet rich plasma in patients with degenerative disc disease', *Journal of translational medicine*. BioMed Central, 15(1), p. 12. doi: 10.1186/s12967-016-1109-0.
- Dailey, L. *et al.* (2005) 'Mechanisms underlying differential responses to FGF signaling.', *Cytokine & growth factor reviews*. England, 16(2), pp. 233–247. doi: 10.1016/j.cytogfr.2005.01.007.
- Darinskas, A. *et al.* (2017) 'Stromal vascular fraction cells for the treatment of critical limb ischemia: a pilot study.', *Journal of translational medicine*. England, 15(1), p. 143. doi: 10.1186/s12967-017-1243-3.
- El-Sharkawy, H. *et al.* (2007) 'Platelet-rich plasma: growth factors and pro- and anti-inflammatory properties.', *Journal of periodontology*. United States, 78(4), pp. 661–669. doi: 10.1902/jop.2007.060302.
- Eppley, B. L., Pietrzak, W. S. and Blanton, M. (2006) 'Platelet-rich plasma: a review of

biology and applications in plastic surgery.’, *Plastic and reconstructive surgery*. United States, 118(6), pp. 147e-159e. doi: 10.1097/01.prs.0000239606.92676.cf.

Eswarakumar, V. P., Lax, I. and Schlessinger, J. (2005) ‘Cellular signaling by fibroblast growth factor receptors.’, *Cytokine & growth factor reviews*. England, 16(2), pp. 139–149. doi: 10.1016/j.cytogfr.2005.01.001.

Ferraro, G. A., Mizuno, H. and Pallua, N. (2016) ‘Adipose Stem Cells: From Bench to Bedside’, *Stem Cells International*. Hindawi Publishing Corporation, 2016, p. 6484038. doi: 10.1155/2016/6484038.

Finch, P. W. *et al.* (1989) ‘Human KGF is FGF-Related with Properties of a Paracrine Effector of Epithelial Cell Growth’, *Science*. American Association for the Advancement of Science, 245(4919), pp. 752–755. Available at: <http://www.jstor.org/stable/1704044>.

Foubert, P. *et al.* (2016) ‘Adipose-derived regenerative cell therapy for burn wound healing: A comparison of two delivery methods’, *Advances in Wound Care*, 5(7), pp. 288–298. doi: 10.1089/wound.2015.0672.

Fu, Y.-S. *et al.* (2006) ‘Conversion of human umbilical cord mesenchymal stem cells in Wharton’s jelly to dopaminergic neurons in vitro: potential therapeutic application for Parkinsonism.’, *Stem cells (Dayton, Ohio)*. United States, 24(1), pp. 115–124. doi: 10.1634/stemcells.2005-0053.

Gentile, P. *et al.* (2017) ‘Concise Review: The Use of Adipose-Derived Stromal Vascular Fraction Cells and Platelet Rich Plasma in Regenerative Plastic Surgery.’, *Stem cells (Dayton, Ohio)*. United States, 35(1), pp. 117–134. doi: 10.1002/stem.2498.

Ghadially, R., Halkier-Sorensen, L. and Elias, P. M. (1992) ‘Effects of petrolatum on stratum corneum structure and function’, *Journal of the American Academy of Dermatology*, 26(3), pp. 387–396. doi: [https://doi.org/10.1016/0190-9622\(92\)70060-S](https://doi.org/10.1016/0190-9622(92)70060-S).

Ghieh, F. *et al.* (2015) ‘The Use of Stem Cells in Burn Wound Healing: A Review’, *BioMed Research International*. Edited by C. Kasper. Hindawi Publishing Corporation, 2015, p. 684084. doi: 10.1155/2015/684084.

Gillenwater, J. and Garner, W. L. (2020) ‘Thermal, Chemical, and Electrical Injuries’, in Kevin C. Chung (ed.) *Grabb and Smith’s plastic surgery*. Eighth. Philadelphia: Wolters Kluwer Health.

Gimble, J. and Guilak, F. (2003) ‘Adipose-derived adult stem cells: isolation, characterization, and differentiation potential.’, *Cytotherapy*. England, 5(5), pp. 362–369. doi: 10.1080/14653240310003026.

Gimble, J. M., Katz, A. J. and Bunnell, B. A. (2007) ‘Adipose-derived stem cells for regenerative medicine.’, *Circulation research*. United States, 100(9), pp. 1249–1260. doi: 10.1161/01.RES.0000265074.83288.09.

- Goldfarb, M. (2005) 'Fibroblast growth factor homologous factors: evolution, structure, and function', *Cytokine & growth factor reviews*. 2005/03/23, 16(2), pp. 215–220. doi: 10.1016/j.cytogfr.2005.02.002.
- Guo, H.-F. *et al.* (2017) 'A new model for studying deep partial-thickness burns in rats.', *International journal of burns and trauma*. United States, 7(6), pp. 107–114.
- Han, J. *et al.* (2010) 'Adipose tissue is an extramedullary reservoir for functional hematopoietic stem and progenitor cells.', *Blood*. United States, 115(5), pp. 957–964. doi: 10.1182/blood-2009-05-219923.
- Harrison, D. A. (2012) 'The Jak/STAT pathway.', *Cold Spring Harbor perspectives in biology*. United States, 4(3). doi: 10.1101/cshperspect.a011205.
- Hayes, M. *et al.* (2012) 'Clinical review: Stem cell therapies for acute lung injury/acute respiratory distress syndrome - hope or hype?', *Critical care (London, England)*. England, 16(2), p. 205. doi: 10.1186/cc10570.
- Herndon, D. N. *et al.* (1997) 'Growth Hormone Therapy in Human Burn Injury', in Ziegler, T. R., Pierce, G. F., and Herndon, D. N. (eds) *Growth Factors and Wound Healing*. New York, NY: Springer New York, pp. 231–244.
- Hirase, T. *et al.* (2018) 'Topical application of platelet-rich plasma for diabetic foot ulcers: A systematic review', *World Journal of Diabetes*, 9(10), pp. 172–179. doi: 10.4239/wjd.v9.i10.172.
- Holland, E. C. and Varmus, H. E. (1998) 'Basic fibroblast growth factor induces cell migration and proliferation after glia-specific gene transfer in mice.', *Proceedings of the National Academy of Sciences of the United States of America*. United States, 95(3), pp. 1218–1223. doi: 10.1073/pnas.95.3.1218.
- Hombach-Klonisch, S. *et al.* (2008) 'Adult stem cells and their trans-differentiation potential—perspectives and therapeutic applications', *Journal of molecular medicine (Berlin, Germany)*. 2008/07/16, 86(12), pp. 1301–1314. doi: 10.1007/s00109-008-0383-6.
- Horwitz, E. M. *et al.* (2005) 'Clarification of the nomenclature for MSC: The International Society for Cellular Therapy position statement.', *Cytotherapy*. England, 7(5), pp. 393–395. doi: 10.1080/14653240500319234.
- Hosni Ahmed, H. *et al.* (2017) 'Can mesenchymal stem cells pretreated with platelet-rich plasma modulate tissue remodeling in a rat with burned skin?', *Biochemistry and Cell Biology*, 95(5), pp. 537–548. doi: 10.1139/bcb-2016-0224.
- Hu, Z. *et al.* (2009) 'Platelet-rich plasma induces mRNA expression of VEGF and PDGF in rat bone marrow stromal cell differentiation.', *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics*. United States, 107(1), pp. 43–48. doi: 10.1016/j.tripleo.2008.06.023.

- Itoh, N. and Ornitz, D. M. (2004) 'Evolution of the Fgf and Fgfr gene families', *Trends in Genetics*. Department of Genetic Biochemistry, Kyoto University Graduate School of Pharmaceutical Sciences, Yoshida-Shimoadachi, Sakyo, Kyoto 606-8501, Japan. itohnobu@pharm.kyoto-u.ac.jp, 20(11), pp. 563–569. doi: 10.1016/j.tig.2004.08.007.
- Itoh, N. and Ornitz, D. M. (2008) 'Functional evolutionary history of the mouse Fgf gene family', *Developmental Dynamics*, 237(1), pp. 18–27. doi: 10.1002/dvdy.21388.
- Josh, F. *et al.* (2012) 'Accelerated and safe proliferation of human adipose-derived stem cells in medium supplemented with human serum.', *Journal of Nippon Medical School = Nippon Ika Daigaku zasshi*. Japan, 79(6), pp. 444–452. doi: 10.1272/jnms.79.444.
- Karagergou, E. *et al.* (2018) 'Adipose-derived stromal vascular fraction aids epithelialisation and angiogenesis in an animal model', *Journal of Wound Care*, 27, pp. 637–644. doi: 10.12968/jowc.2018.27.10.637.
- Karina *et al.* (2019) 'Combination of the stromal vascular fraction and platelet-rich plasma accelerates the wound healing process: pre-clinical study in a Sprague-Dawley rat model', *Stem cell investigation*. AME Publishing Company, 6, p. 18. doi: 10.21037/sci.2019.06.08.
- Kevy, S., Jacobson, M. and Benoit, P. (2001) 'The biology of platelet concentrate as prepared by the Harvest Technologies Smart-PReP System', in *Proceedings of the 3rd Annual Meeting of Techvest Conference on Tissue Repair, Replacement, and Regeneration*.
- Kim, D.-Y. *et al.* (2014) 'Effects of platelet-rich plasma, adipose-derived stem cells, and stromal vascular fraction on the survival of human transplanted adipose tissue', *Journal of Korean medical science*. 2014/11/21. The Korean Academy of Medical Sciences, 29 Suppl 3(Suppl 3), pp. S193–S200. doi: 10.3346/jkms.2014.29.S3.S193.
- Kirschstein, R. (2001) 'Stem cells: scientific progress and future directions', *National Institutes of Health*, pp. 1–222.
- Kolkova, K. *et al.* (2000) 'Neural cell adhesion molecule-stimulated neurite outgrowth depends on activation of protein kinase C and the Ras-mitogen-activated protein kinase pathway', *The Journal of neuroscience : the official journal of the Society for Neuroscience*. Protein Laboratory, Institute of Molecular Pathology, University of Copenhagen, DK-2200, Copenhagen N, Denmark., 20(6), pp. 2238–2246. doi: 10.1523/JNEUROSCI.20-06-02238.2000.
- Komi-Kuramochi, A. *et al.* (2005) 'Expression of fibroblast growth factors and their receptors during full-thickness skin wound healing in young and aged mice', *Journal of Endocrinology*, 186(2), pp. 273–289. doi: 10.1677/joe.1.06055.
- Kotani, T. *et al.* (2017) 'Anti-inflammatory and anti-fibrotic effects of intravenous adipose-derived stem cell transplantation in a mouse model of bleomycin-induced interstitial pneumonia', *Scientific Reports*. Springer US, 7(1), pp. 1–10. doi:

10.1038/s41598-017-15022-3.

Lawrence, W. T. (1998) 'Physiology of the acute wound.', *Clinics in plastic surgery*. United States, 25(3), pp. 321–340.

Le, T. M. *et al.* (2019) 'The sustained release of basic fibroblast growth factor accelerates angiogenesis and the engraftment of the inactivated dermis by high hydrostatic pressure', *PLOS ONE*. Edited by F. Zhao. Public Library of Science, 14(2), p. e0208658. doi: 10.1371/journal.pone.0208658.

Li, H. *et al.* (2009) '[Experimental research of the promotion effect of autogeneic PRP on osteogenic differentiation of human adipose-derived stem cells in vitro].', *Zhongguo xiu fu chong jian wai ke za zhi = Zhongguo xiufu chongjian waike zazhi = Chinese journal of reparative and reconstructive surgery*. China, 23(6), pp. 732–736.

Lo, B. and Parham, L. (2009) 'Ethical issues in stem cell research', *Endocrine reviews*. 2009/04/14. The Endocrine Society, 30(3), pp. 204–213. doi: 10.1210/er.2008-0031.

MacLeod, A. S. and Mansbridge, J. N. (2016) 'The Innate Immune System in Acute and Chronic Wounds', *Advances in wound care*. Mary Ann Liebert, Inc., 5(2), pp. 65–78. doi: 10.1089/wound.2014.0608.

Maddaluno, L., Urwyler, C. and Werner, S. (2017) 'Fibroblast growth factors: key players in regeneration and tissue repair.', *Development (Cambridge, England)*. England, 144(22), pp. 4047–4060. doi: 10.1242/dev.152587.

Mansoub, N. H. *et al.* (2018) 'The role of PRP and adipose tissue-derived keratinocytes on burn wound healing in diabetic rats', *BioImpacts*, 8(1), pp. 5–12. doi: 10.15171/bi.2018.02.

Marck, R., Middelkoop, M. and Breederveld, R. (2018) 'Considerations on the use of platelet rich plasma, specifically for burn treatment: Journal of burn care & research May-Jun 2014;35(3):219-227', in *On PLATELETS and burns*, pp. 20–34.

McGee, G. S. *et al.* (1988) 'Recombinant basic fibroblast growth factor accelerates wound healing.', *The Journal of surgical research*. United States, 45(1), pp. 145–153. doi: 10.1016/0022-4804(88)90034-0.

Moenadjat, Y. *et al.* (2013) 'The application of human umbilical cord blood mononuclear cells in the management of deep partial thickness burn', *Medical Journal of Indonesia*, 22(2 SE-Clinical Research). doi: 10.13181/mji.v22i2.534.

Motamed, S. *et al.* (2017) 'Cell-based skin substitutes accelerate regeneration of extensive burn wounds in rats', *American Journal of Surgery*, 214(4), pp. 762–769. doi: 10.1016/j.amjsurg.2017.04.010.

Nauta, A. *et al.* (2013) 'Adipose-derived stromal cells overexpressing vascular endothelial growth factor accelerate mouse excisional wound healing.', *Molecular*

therapy: the journal of the American Society of Gene Therapy. United States, 21(2), pp. 445–455. doi: 10.1038/mt.2012.234.

Nazzal, M. *et al.* (2019) ‘Wound Healing’, in F. Charles Brunnicardi *et al.* (eds) *Schwartz’s Principles of Surgery*. Eleventh. New York: McGraw-Hill, pp. 271–304.

Nikolidakis, D. and Jansen, J. A. (2008) ‘The biology of platelet-rich plasma and its application in oral surgery: literature review.’, *Tissue engineering. Part B, Reviews*. United States, 14(3), pp. 249–258. doi: 10.1089/ten.teb.2008.0062.

Oryan, A., Alemzadeh, E. and Moshiri, A. (2017) ‘Burn wound healing: Present concepts, treatment strategies and future directions’, *Journal of Wound Care*, 26(1), pp. 5–19. doi: 10.12968/jowc.2017.26.1.5.

Park, J. W., Hwang, S. R. and Yoon, I.-S. (2017) ‘Advanced Growth Factor Delivery Systems in Wound Management and Skin Regeneration’, *Molecules (Basel, Switzerland)*. MDPI, 22(8), p. 1259. doi: 10.3390/molecules22081259.

Patel, D. P. *et al.* (2015) ‘High-grade renal injuries are often isolated in sports-related trauma’, *Injury*. 2015/03/01, 46(7), pp. 1245–1249. doi: 10.1016/j.injury.2015.02.008.

Patel, P., Duttaroy, D. and Kacheriwala, S. (2014) ‘Management of renal injuries in blunt abdominal trauma’, *Journal of Research in Medical and Dental Science*, 2(2), p. 38. doi: 10.5455/jrmds.2014229.

Pearson, G. *et al.* (2001) ‘Mitogen-activated protein (MAP) kinase pathways: regulation and physiological functions.’, *Endocrine reviews*. United States, 22(2), pp. 153–183. doi: 10.1210/edrv.22.2.0428.

Petry, T. *et al.* (2017) ‘Review of data on the dermal penetration of mineral oils and waxes used in cosmetic applications’, *Toxicology Letters*, 280, pp. 70–78. doi: <https://doi.org/10.1016/j.toxlet.2017.07.899>.

Van Pham, P. *et al.* (2013) ‘Activated platelet-rich plasma improves adipose-derived stem cell transplantation efficiency in injured articular cartilage.’, *Stem cell research & therapy*. England, 4(4), p. 91. doi: 10.1186/scrt277.

Plichta, J. K. and Radek, K. A. (2012) ‘Sugar-Coating Wound Repair: A Review of FGF-10 and Dermatan Sulfate in Wound Healing and Their Potential Application in Burn Wounds’, *Journal of Burn Care & Research*, 33(3), pp. 299–310. doi: 10.1097/BCR.0b013e318240540a.

Powers, C. J., McLeskey, S. W. and Wellstein, A. (2000) ‘Fibroblast growth factors, their receptors and signaling.’, *Endocrine-related cancer*. England, 7(3), pp. 165–197. doi: 10.1677/erc.0.0070165.

Raposio, E. *et al.* (2016) ‘Adipose-derived Stem Cells Added to Platelet-rich Plasma for Chronic Skin Ulcer Therapy.’, *Wounds: a compendium of clinical research and*

practice. United States, 28(4), pp. 126–131.

Rigotti, G., Marchi, A. and Sbarbati, A. (2009) ‘Adipose-derived mesenchymal stem cells: past, present, and future.’, *Aesthetic plastic surgery*. United States, pp. 271–273. doi: 10.1007/s00266-009-9339-7.

Rohovsky, S. and D’Amore, P. A. (1997) ‘Growth Factors and Angiogenesis in Wound Healing’, in Thomas R. Ziegler, Pierce, G. F., and Herndon, D. N. (eds) *Growth Factors and Wound Healing*. New York, NY: Springer New York, pp. 8–26. doi: 10.1007/978-1-4612-1876-0_2.

Rose, L. F. and Chan, R. K. (2016) ‘The Burn Wound Microenvironment’, *Advances in Wound Care*, 5(3), pp. 106–118. doi: 10.1089/wound.2014.0536.

Rosenstrauch, D. *et al.* (2005) ‘Stem celltherapy for ischemic heart failure’, *Texas Heart Institute journal*, 32(3), pp. 339–347. Available at: <https://pubmed.ncbi.nlm.nih.gov/16392214>.

Rumalla, V. K. and Borah, G. L. (2001) ‘Cytokines, growth factors, and plastic surgery.’, *Plastic and reconstructive surgery*. United States, 108(3), pp. 719–733. doi: 10.1097/00006534-200109010-00019.

Said, A. *et al.* (2019) ‘Sauromatum guttatum extract promotes wound healing and tissue regeneration in a burn mouse model via up-regulation of growth factors’, *Pharmaceutical Biology*. Taylor & Francis, 57(1), pp. 736–743. doi: 10.1080/13880209.2019.1676266.

Schaffer, C. J. and Nanney, L. B. (1996) ‘Cell biology of wound healing.’, *International review of cytology*. United States, 169, pp. 151–181. doi: 10.1016/s0074-7696(08)61986-5.

Schöler, H. R. (2016) ‘The Potential of Stem Cells: An Inventory’, in Schipanski, D., Knoepfle, N., and S L Sorgner (eds) *Humanbiotechnology as Social Challenge An Interdisciplinary Introduction to Bioethics*. First. London: Taylor & Francis, pp. 1–28.

Sethi, A. *et al.* (2016) ‘Moisturizers: The Slippery Road’, *Indian journal of dermatology*. Medknow Publications & Media Pvt Ltd, 61(3), pp. 279–287. doi: 10.4103/0019-5154.182427.

Seyhan, N. *et al.* (2015) ‘The Effect of Combined Use of Platelet-Rich Plasma and Adipose-Derived Stem Cells on Fat Graft Survival’, *Annals of Plastic Surgery*, 74(5), pp. 615–620. doi: 10.1097/SAP.0000000000000480.

Shpichka, A. *et al.* (2019) ‘Skin tissue regeneration for burn injury’, *Stem Cell Research and Therapy*. Stem Cell Research & Therapy, 10(1), pp. 1–16. doi: 10.1186/s13287-019-1203-3.

Shukla, S. K. *et al.* (2020) ‘Can miRNAs serve as potential markers in thermal burn

injury: An in silico approach', *Journal of Burn Care and Research*, 41(1), pp. 57–64. doi: 10.1093/jbcr/irz183.

Singh, V. K. *et al.* (2016) 'Describing the Stem Cell Potency: The Various Methods of Functional Assessment and In silico Diagnostics.', *Frontiers in cell and developmental biology*. Switzerland, 4, p. 134. doi: 10.3389/fcell.2016.00134.

Tajima, S. *et al.* (2014) 'Direct and Indirect Effects of a Combination of Adipose-Derived Stem Cells and Platelet-Rich Plasma on Bone Regeneration', *Tissue engineering. Part A*, 21. doi: 10.1089/ten.TEA.2014.0336.

Tantuway, V. *et al.* (2016) 'Autologous grafting of non manipulated freshly isolated - adipose tissue derived stromal vascular fraction in single surgical sitting for treatment of knee osteoarthritis', *International Journal of Research in Orthopaedics*, 3, p. 107. doi: 10.18203/issn.2455-4510.IntJResOrthop20164834.

Tavares Pereira, D. dos S. *et al.* (2012) 'Development of Animal Model for Studying Deep Second-Degree Thermal Burns', *Journal of Biomedicine and Biotechnology*. Edited by M. Fedele. Hindawi Publishing Corporation, 2012, p. 460841. doi: 10.1155/2012/460841.

Thisse, B. and Thisse, C. (2005) 'Functions and regulations of fibroblast growth factor signaling during embryonic development.', *Developmental biology*. United States, 287(2), pp. 390–402. doi: 10.1016/j.ydbio.2005.09.011.

Tohidnezhad, M. *et al.* (2011) 'Platelet-released growth factors can accelerate tenocyte proliferation and activate the anti-oxidant response element', *Histochemistry and Cell Biology*, 135(5), pp. 453–460. doi: 10.1007/s00418-011-0808-0.

Toussaint, J. and Singer, A. J. (2014) 'The evaluation and management of thermal injuries: 2014 update', *Clinical and experimental emergency medicine*. The Korean Society of Emergency Medicine, 1(1), pp. 8–18. doi: 10.15441/ceem.14.029.

Tsien, L. (2006) 'Stem Cell Basics', *Postgraduate Obstetrics & Gynecology*, 26(24), pp. 1–6. doi: 10.1097/00256406-200612310-00001.

Turner, N. and Grose, R. (2010) 'Fibroblast growth factor signalling: from development to cancer.', *Nature reviews. Cancer*. England, 10(2), pp. 116–129. doi: 10.1038/nrc2780.

Vaghardoost, R. *et al.* (2018) 'The Healing Effect of Sesame Oil, Camphor and Honey on Second Degree Burn Wounds in Rat.', *World journal of plastic surgery*, 7(1), pp. 67–71.

Wang, Y. *et al.* (2018) 'Burn injury: Challenges and advances in burn wound healing, infection, pain and scarring', *Advanced Drug Delivery Reviews*. Elsevier B.V., 123, pp. 3–17. doi: 10.1016/j.addr.2017.09.018.

- Werner, S. (1998) 'Keratinocyte growth factor: a unique player in epithelial repair processes.', *Cytokine & growth factor reviews*. England, 9(2), pp. 153–165. doi: 10.1016/s1359-6101(98)00010-0.
- Werner, S. and Grose, R. (2003) 'Regulation of wound healing by growth factors and cytokines.', *Physiological reviews*. United States, 83(3), pp. 835–870. doi: 10.1152/physrev.2003.83.3.835.
- Widowati, W. and Widyanto, R. M. (2013) 'Sel Punca sebagai Transformasi Alternatif Terapi', *Zenit*, 2(1), pp. 1–5.
- Witte, M. B. and Barbul, A. (1997) 'General principles of wound healing.', *The Surgical clinics of North America*. United States, 77(3), pp. 509–528. doi: 10.1016/s0039-6109(05)70566-1.
- Wobus, A. M. and Boheler, K. R. (2005) 'Embryonic Stem Cells: Prospects for Developmental Biology and Cell Therapy', *Physiological Reviews*. American Physiological Society, 85(2), pp. 635–678. doi: 10.1152/physrev.00054.2003.
- Wong, A. *et al.* (2002) 'FRS2 alpha attenuates FGF receptor signaling by Grb2-mediated recruitment of the ubiquitin ligase Cbl.', *Proceedings of the National Academy of Sciences of the United States of America*. United States, 99(10), pp. 6684–6689. doi: 10.1073/pnas.052138899.
- Yun, Y.-R. *et al.* (2010) 'Fibroblast growth factors: biology, function, and application for tissue regeneration', *Journal of tissue engineering*. SAGE-Hindawi Access to Research, 2010, p. 218142. doi: 10.4061/2010/218142.
- Zakrzewski, W. *et al.* (2019) 'Stem cells: past, present, and future', *Stem Cell Research & Therapy*, 10(1), p. 68. doi: 10.1186/s13287-019-1165-5.
- Zhang, Y. *et al.* (2011) '[Effect of platelet-rich plasma on the proliferation and adipogenic differentiation of human adipose-derived stem cells in vitro].', *Nan fang yi ke da xue xue bao = Journal of Southern Medical University*. China, 31(3), pp. 525–528.
- Zuk, P. A. *et al.* (2002) 'Human adipose tissue is a source of multipotent stem cells', *Molecular biology of the cell*. The American Society for Cell Biology, 13(12), pp. 4279–4295. doi: 10.1091/mbc.e02-02-0105.