

DAFTAR PUSTAKA

- Abu-Humaidan, A. H. A. *et al.* (2014) 'The Epidermal Growth Factor Receptor Is a Regulator of Epidermal Complement Component Expression and Complement Activation', *The Journal of Immunology*, 192(7), pp. 3355 LP – 3364. doi: 10.4049/jimmunol.1302305.
- Aleckovic, M. and Simon, C. (2008) 'Is teratoma formation in stem cell research a characterization tool or a window to developmental biology?', *Reproductive biomedicine online*. Netherlands, 17(2), pp. 270–280. doi: 10.1016/s1472- 6483(10)60206-4.
- ANZBA (2016) Emergency Management of Severe Burns (EMSB). 18th edn. Albany Creek: THE EDUCATION COMMITTEE OFAUSTRALIA AND NEW ZEALAND BURN ASSOCIATION LTD.
- Baglioni, S. *et al.* (2009) 'Characterization of human adult stem-cell populations isolated from visceral and subcutaneous adipose tissue.', *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. United States, 23(10), pp. 3494–3505. doi: 10.1096/fj.08-126946.
- Bakacak, M. *et al.* (2016) 'Protective Effect of Platelet Rich Plasma on Experimental Ischemia/Reperfusion Injury in Rat Ovary.', *Gynecologic and obstetric investigation*. Switzerland, 81(3), pp. 225–231. doi: 10.1159/000440617.
- Barret-Nerin, J. (2004) Principles and Practice of Burn Surgery. CRC Press. doi: 10.1201/b21634.
- Bodnar, R. J. (2013) 'Epidermal Growth Factor and Epidermal Growth Factor Receptor: The Yin and Yang in the Treatment of Cutaneous Wounds and Cancer', *Advances in wound care*. Mary Ann Liebert, Inc., 2(1), pp. 24–29. doi: 10.1089/wound.2011.0326.
- Borrione, P. *et al.* (2010) 'Platelet-rich plasma in muscle healing.', *American journal of physical medicine & rehabilitation*. United States, 89(10), pp. 854–861. doi: 10.1097/PHM.0b013e3181f1c1c7.
- Bourin, P. *et al.* (2013) 'Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International Society for Cytotherapy', *Cytotherapy*. England, 15(6), pp. 641–648. doi: 10.1016/j.jcyt.2013.02.006.
- Braund, R., Hook, S. and Medlicott, N. J. (2007) 'The role of topical growth

- factors in chronic wounds.', Current drug delivery. United Arab Emirates, 4(3), pp. 195–204. doi: 10.2174/156720107781023857.
- Cerqueira, M. T., Pirraco, R. P. and Marques, A. P. (2016) 'Stem Cells in Skin Wound Healing: Are We There Yet?', Advances in Wound Care, 5(4), pp. 164–175. doi: 10.1089/wound.2014.0607.
- Cervelli, V. et al. (2010) 'Tissue regeneration in loss of substance on the lower limbs through use of platelet-rich plasma, stem cells from adipose tissue, and hyaluronic acid.', Advances in skin & wound care. United States, 23(6), pp. 262–272. doi: 10.1097/01.ASW.0000363551.82058.36.
- Chieregato, K. et al. (2011) 'Epidermal growth factor, basic fibroblast growth factor and platelet-derived growth factor-bb can substitute for fetal bovine serum and compete with human platelet-rich plasma in the ex vivo expansion of mesenchymal stromal cells derived from adipose tissue', Cytotherapy. England, 13(8), pp. 933–943. doi: 10.3109/14653249.2011.583232.
- Choi, J., Minn, K. W. and Chang, H. (2012) 'The efficacy and safety of platelet-rich plasma and adipose-derived stem cells: an update', Archives of plastic surgery. 2012/11/14. The Korean Society of Plastic and Reconstructive Surgeons, 39(6), pp. 585–592. doi: 10.5999/aps.2012.39.6.585.
- Comella, K., Silbert, R. and Parlo, M. (2017) 'Effects of the intradiscal implantation of stromal vascular fraction plus platelet rich plasma in patients with degenerative disc disease', Journal of translational medicine. BioMed Central, 15(1), p. 12. doi: 10.1186/s12967-016-1109-0.
- Dailey, L. et al. (2005) 'Mechanisms underlying differential responses to FGF signaling.', Cytokine & growth factor reviews. England, 16(2), pp. 233–247. doi: 10.1016/j.cytopfr.2005.01.007.
- Darinskas, A. et al. (2017) 'Stromal vascular fraction cells for the treatment of critical limb ischemia: a pilot study.', Journal of translational medicine. England, 15(1), p. 143. doi: 10.1186/s12967-017-1243-3.
- Desai, T. J. and Cardoso, W. V (2002) 'Growth factors in lung development and disease: friends or foe?', Respiratory research. 2001/10/09. BioMed Central, 3(1), p. 2. doi: 10.1186/rr169.
- Duran, Aslı et al. (2016) Clinical and histopathological evaluation of the effects of platelet rich plasma, platelet poor plasma and topical serum physiologic treatment on wound healing caused by radiofrequency electrosurgery in

rats.

- Eppley, B. L., Pietrzak, W. S. and Blanton, M. (2006) ‘Platelet-rich plasma: a review of biology and applications in plastic surgery.’, Plastic and reconstructive surgery. United States, 118(6), pp. 147e-159e. doi: 10.1097/01.prs.0000239606.92676.cf.
- Esquirol Caussa, J. and Herrero Vila, E. (2015) ‘Epidermal growth factor, innovation and safety’, Medicina Clínica (English Edition), 145(7), pp. 305–312. doi: 10.1016/j.medcle.2016.02.028.
- Falanga, V. et al. (1992) ‘Topical use of human recombinant epidermal growth factor (h-EGF) in venous ulcers.’, The Journal of dermatologic surgery and oncology. United States, 18(7), pp. 604–606. doi: 10.1111/j.1524-4725.1992.tb03514.x.
- Ferraro, G. A., Mizuno, H. and Pallua, N. (2016) ‘Adipose Stem Cells: From Bench to Bedside’, Stem Cells International. Hindawi Publishing Corporation, 2016, p. 6484038. doi: 10.1155/2016/6484038.
- Foubert, P. et al. (2016) ‘Adipose-derived regenerative cell therapy for burn wound healing: A comparison of two delivery methods’, Advances in Wound Care, 5(7), pp. 288–298. doi: 10.1089/wound.2015.0672.
- Fu, Y.-S. et al. (2006) ‘Conversion of human umbilical cord mesenchymal stem cells in Wharton’s jelly to dopaminergic neurons in vitro: potential therapeutic application for Parkinsonism.’, Stem cells (Dayton, Ohio). United States, 24(1), pp. 115–124. doi: 10.1634/stemcells.2005-0053.
- Gentile, P. et al. (2017) ‘Concise Review: The Use of Adipose-Derived Stromal Vascular Fraction Cells and Platelet Rich Plasma in Regenerative Plastic Surgery.’, Stem cells (Dayton, Ohio). United States, 35(1), pp. 117–134. doi: 10.1002/stem.2498.
- Ghadially, R., Halkier-Sorensen, L. and Elias, P. M. (1992) ‘Effects of petrolatum on stratum corneum structure and function’, Journal of the American Academy of Dermatology, 26(3), pp. 387–396. doi: [https://doi.org/10.1016/0190-9622\(92\)70060-S](https://doi.org/10.1016/0190-9622(92)70060-S).
- Ghieh, F. et al. (2015) ‘The Use of Stem Cells in Burn Wound Healing: A Review’, BioMed Research International. Edited by C. Kasper. Hindawi Publishing Corporation, 2015, p. 684084. doi: 10.1155/2015/684084.
- Gillenwater, J. and Garner, W. L. (2020) ‘Thermal, Chemical, and Electrical

- Injuries', in Kevin C. Chung (ed.) Grabb and Smith's plastic surgery. Eighth. Philadelphia: Wolters Kluwer Health.
- Gimble, J. and Guilak, F. (2003) 'Adipose-derived adult stem cells: isolation, characterization, and differentiation potential.', *Cyotherapy*. England, 5(5), pp. 362– 369. doi: 10.1080/14653240310003026.
- Gimble, J. M., Katz, A. J. and Bunnell, B. A. (2007) 'Adipose-derived stem cells for regenerative medicine.', *Circulation research*. United States, 100(9), pp. 1249–1260. doi: 10.1161/01.RES.0000265074.83288.09.
- Grant, M. B. et al. (1992) 'Effects of epidermal growth factor, fibroblast growth factor, and transforming growth factor-beta on corneal cell chemotaxis.', *Investigative ophthalmology & visual science*. United States, 33(12), pp. 3292–3301.
- Grigore, T. V. and Cozma, C. (2018) 'Platelet-rich plasma as a site-targeted approach in wound healing: a molecular perspective', *Discoveries*, 6(4), pp. 1–8. doi: 10.15190/d.2018.8.
- Guo, H.-F. et al. (2017) 'A new model for studying deep partial-thickness burns in rats.', *International journal of burns and trauma*. United States, 7(6), pp. 107–114.
- Hardwicke, J. et al. (2008) 'Epidermal growth factor therapy and wound healing-- past, present and future perspectives.', *The surgeon : journal of the Royal Colleges of Surgeons of Edinburgh and Ireland*. Scotland, 6(3), pp. 172–177. doi: 10.1016/s1479- 666x(08)80114-x.
- Harris, R. C., Chung, E. and Coffey, R. J. (2003) 'EGF receptor ligands.', *Experimental cell research*. United States, 284(1), pp. 2–13. doi: 10.1016/s0014-4827(02)00105-2.
- Harrison, D. A. (2012) 'The Jak/STAT pathway.', *Cold Spring Harbor perspectives in biology*. United States, 4(3). doi: 10.1101/cshperspect.a011205.
- Hausman, G. J. and Richardson, R. L. (2004) 'Adipose tissue angiogenesis.', *Journal of animal science*. United States, 82(3), pp. 925–934. doi: 10.2527/2004.823925x.
- Hayes, M. et al. (2012) 'Clinical review: Stem cell therapies for acute lung injury/acute respiratory distress syndrome - hope or hype?', *Critical care* (London, England), 16(2), p. 205. doi: 10.1186/cc10570.

- Herndon, D. N. et al. (1997) ‘Growth Hormone Therapy in Human Burn Injury’, in Ziegler, T. R., Pierce, G. F., and Herndon, D. N. (eds) *Growth Factors and Wound Healing*. New York, NY: Springer New York, pp. 231–244.
- Hombach-Klonisch, S. et al. (2008) ‘Adult stem cells and their trans-differentiation potential--perspectives and therapeutic applications’, *Journal of molecular medicine (Berlin, Germany)*. 2008/07/16, 86(12), pp. 1301–1314. doi: 10.1007/s00109-008-0383-6.
- Hu, Z. et al. (2009) ‘Platelet-rich plasma induces mRNA expression of VEGF and PDGF in rat bone marrow stromal cell differentiation.’, *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics*. United States, 107(1), pp. 43–48. doi: 10.1016/j.tripleo.2008.06.023.
- Josh, F. et al. (2012) ‘Accelerated and safe proliferation of human adipose-derived stem cells in medium supplemented with human serum.’, *Journal of Nippon Medical School*
= Nippon Ika Daigaku zasshi. Japan, 79(6), pp. 444–452. doi: 10.1272/jnms.79.444.
- Ju, W. D. et al. (1993) ‘TGF alpha enhances locomotion of cultured human keratinocytes.’, *The Journal of investigative dermatology*. United States, 100(5), pp. 628–632.
- Karagergou, E. et al. (2018) ‘Adipose-derived stromal vascular fraction aids epithelialisation and angiogenesis in an animal model’, *Journal of Wound Care*, 27, pp. 637–644. doi: 10.12968/jowc.2018.27.10.637.
- Karina et al. (2019) ‘Combination of the stromal vascular fraction and platelet-rich plasma accelerates the wound healing process: pre-clinical study in a Sprague-Dawley rat model’, *Stem cell investigation*. AME Publishing Company, 6, p. 18. doi: 10.21037/sci.2019.06.08.
- Kim, D.-Y. et al. (2014) ‘Effects of platelet-rich plasma, adipose-derived stem cells, and stromal vascular fraction on the survival of human transplanted adipose tissue’, *Journal of Korean medical science*. 2014/11/21. The Korean Academy of Medical Sciences, 29 Suppl 3(Suppl 3), pp. S193–S200. doi: 10.3346/jkms.2014.29.S3.S193.
- Kirschstein, R. (2001) ‘Stem cells: scientific progress and future directions’, National Institutes of Health, pp. 1–222.
- Kuwada, S. K. and Li, X. (2000) ‘Integrin alpha5/beta1 mediates fibronectin-dependent epithelial cell proliferation through epidermal growth factor receptor activation’, *Molecular biology of the cell*. The American Society for Cell Biology, 11(7), pp. 2485–2496. doi: 10.1091/mbc.11.7.2485.

- Lawrence, W. T. (1998) 'Physiology of the acute wound.', *Clinics in plastic surgery*. United States, 25(3), pp. 321–340.
- Liu, L. et al. (2009) 'Evaluation of Interleukin-8 in Expressed Prostatic Secretion as a Reliable Biomarker of Inflammation in Benign Prostatic Hyperplasia', *Urology*. Elsevier Inc., 74(2), pp. 340–344. doi: 10.1016/j.urology.2009.02.064.
- Lo, B. and Parham, L. (2009) 'Ethical issues in stem cell research', *Endocrine reviews*. 2009/04/14. The Endocrine Society, 30(3), pp. 204–213. doi: 10.1210/er.2008-0031.
- MacLeod, A. S. and Mansbridge, J. N. (2016) 'The Innate Immune System in Acute and Chronic Wounds', *Advances in wound care*. Mary Ann Liebert, Inc., 5(2), pp. 65–78. doi: 10.1089/wound.2014.0608.
- Mansoub, N. H. et al. (2018) 'The role of PRP and adipose tissue-derived keratinocytes on burn wound healing in diabetic rats', *BioImpacts*, 8(1), pp. 5–12. doi: 10.15171/bi.2018.02.
- Marck, R., Middelkoop, M. and Breederveld, R. (2018) 'Considerations on the use of platelet rich plasma, specifically for burn treatment: Journal of burn care & research May-Jun 2014;35(3):219-227', in *On PLATELETS and burns*, pp. 20–34.
- McGee, G. S. et al. (1988) 'Recombinant basic fibroblast growth factor accelerates wound healing.', *The Journal of surgical research*. United States, 45(1), pp. 145–153. doi: 10.1016/0022-4804(88)90034-0.
- Minori, N. (2019) 'Penyembuhan Luka Bakar Derajat Mid Dermal Pasca Pemberian PRP (Platelet Rich Plasma) pada Tikus Wistar'.
- Moenadjat, Y. et al. (2013) 'The application of human umbilical cord blood mononuclear cells in the management of deep partial thickness burn', *Medical Journal of Indonesia*, 22(2 SE-Clinical Research). doi: 10.13181/mji.v22i2.534.
- Mohammed, M. O. et al. (2019) 'Role of Bovine Fetal Platelet-rich Plasma (PRP) on Skin Wound Healing in Mice', *International Journal of Pharmacology*, 16(1), pp. 18– 26. doi: 10.3923/ijp.2020.18.26.
- Momeni, M. et al. (2019) 'A randomized, double-blind, phase I clinical trial of fetal cell-based skin substitutes on healing of donor sites in burn patients', *Burns*. Elsevier Ltd and International Society of Burns Injuries, 45(4), pp. 914–922. doi: 10.1016/j.burns.2018.10.016.

- Moss, L. S. (2010) ‘Treatment of the burn patient in primary care.’, Advances in skin & wound care. United States, 23(11), pp. 516–517. doi: 10.1097/01.ASW.0000390374.34201.c8.
- Motamed, S. et al. (2017) ‘Cell-based skin substitutes accelerate regeneration of extensive burn wounds in rats’, American Journal of Surgery, 214(4), pp. 762–769. doi: 10.1016/j.amjsurg.2017.04.010.
- Nauta, A. et al. (2013) ‘Adipose-derived stromal cells overexpressing vascular endothelial growth factor accelerate mouse excisional wound healing.’, Molecular therapy : the journal of the American Society of Gene Therapy. United States, 21(2), pp. 445–455. doi: 10.1038/mt.2012.234.
- Nazzal, M. et al. (2019) ‘Wound Healing’, in F. Charles Brunicardi et al. (eds) Schwartz’s Principles of Surgery. Eleventh. New York: McGraw-Hill, pp. 271–304.
- Nexø, E., Jørgensen, P. E. and Hansen, M. R. (1992) ‘Human epidermal growth factor—on molecular forms present in urine and blood’, Regulatory Peptides, 42(1), pp. 75–84. doi: [https://doi.org/10.1016/0167-0115\(92\)90025-P](https://doi.org/10.1016/0167-0115(92)90025-P).
- Nikolidakis, D. and Jansen, J. A. (2008) ‘The biology of platelet-rich plasma and its application in oral surgery: literature review.’, Tissue engineering. Part B, Reviews. United States, 14(3), pp. 249–258. doi: 10.1089/ten.teb.2008.0062.
- Oryan, A., Alemzadeh, E. and Moshiri, A. (2017) ‘Burn wound healing: Present concepts, treatment strategies and future directions’, Journal of Wound Care, 26(1), pp. 5–19. doi: 10.12968/jowc.2017.26.1.5.
- Park, J. W., Hwang, S. R. and Yoon, I.-S. (2017) ‘Advanced Growth Factor Delivery Systems in Wound Management and Skin Regeneration’, Molecules (Basel, Switzerland). MDPI, 22(8), p. 1259. doi: 10.3390/molecules22081259.
- Pearson, G. et al. (2001) ‘Mitogen-activated protein (MAP) kinase pathways: regulation and physiological functions.’, Endocrine reviews. United States, 22(2), pp. 153–183. doi: 10.1210/edrv.22.2.0428.
- Petry, T. et al. (2017) ‘Review of data on the dermal penetration of mineral oils and waxes used in cosmetic applications’, Toxicology Letters, 280, pp. 70–78. doi: <https://doi.org/10.1016/j.toxlet.2017.07.899>.
- Van Pham, P. et al. (2013) ‘Activated platelet-rich plasma improves adipose-derived stem cell transplantation efficiency in injured articular cartilage.’,

- Stem cell research & therapy. England, 4(4), p. 91. doi: 10.1186/scrt277.
- Rah, D. K. et al. (2017) ‘Effect of Platelet-Rich Plasma on Ischemia-Reperfusion Injury in a Skin Flap Mouse Model’, International journal of medical sciences. Ivyspring International Publisher, 14(9), pp. 829–839. doi: 10.7150/ijms.19573.
- Raposio, E. et al. (2016) ‘Adipose-derived Stem Cells Added to Platelet-rich Plasma for Chronic Skin Ulcer Therapy.’, Wounds : a compendium of clinical research and practice. United States, 28(4), pp. 126–131.
- Rigotti, G., Marchi, A. and Sbarbati, A. (2009) ‘Adipose-derived mesenchymal stem cells: past, present, and future.’, Aesthetic plastic surgery. United States, pp. 271–273. doi: 10.1007/s00266-009-9339-7.
- Rohovsky, S. and D’Amore, P. A. (1997) ‘Growth Factors and Angiogenesis in Wound Healing’, in Thomas R. Ziegler, Pierce, G. F., and Herndon, D. N. (eds) Growth Factors and Wound Healing. New York, NY: Springer New York, pp. 8–26. doi: 10.1007/978- 1-4612-1876-0_2.
- Rosadi Seswandhana, M. et al. (2020) ‘A case series of negative pressure wound therapy as a promising treatment in patients with burn injury’, International Journal of Surgery Case Reports. Surgical Associates Ltd, 69, pp. 64–67. doi: 10.1016/j.ijscr.2020.03.034.
- Rosenstrauch, D. et al. (2005) ‘Stem celltherapy for ischemic heart failure’, Texas Heart Institute journal, 32(3), pp. 339–347. Available at: <https://pubmed.ncbi.nlm.nih.gov/16392214/>.
- Rumalla, V. K. and Borah, G. L. (2001) ‘Cytokines, growth factors, and plastic surgery.’, Plastic and reconstructive surgery. United States, 108(3), pp. 719–733. doi: 10.1097/00006534-200109010-00019.
- Salehi, H. et al. (2018) ‘Comparing the effect of colactive plus ag dressing versus nitrofurazone and vaseline gauze dressing in the treatment of second-degree burns’, Annals of Burns and Fire Disasters, 31(3), pp. 204–208.
- Satriyo, A., Djukardi, E. K. and Zubier, F. (2011) ‘Peran plasma kaya trombosit (platelet-rich plasma) di bidang dermatologi’, Mdvi, 38(1), pp. 22–28.
- Schaffer, C. J. and Nanney, L. B. (1996) ‘Cell biology of wound healing.’, International review of cytology. United States, 169, pp. 151–181. doi: 10.1016/s0074- 7696(08)61986-5.
- Schöler, H. R. (2016) ‘The Potential of Stem Cells: An Inventory’, in Schipanski, D., Knoepfle, N., and S L Sorgner (eds) Humanbiotechnology as Social

- Challenge An Interdisciplinary Introduction to Bioethics. First. London: Taylor & Francis, pp. 1–28.
- Sethi, A. et al. (2016) ‘Moisturizers: The Slippery Road’, Indian journal of dermatology. Medknow Publications & Media Pvt Ltd, 61(3), pp. 279–287. doi: 10.4103/0019-5154.182427.
- Shpichka, A. et al. (2019) ‘Skin tissue regeneration for burn injury’, Stem Cell Research and Therapy. Stem Cell Research & Therapy, 10(1), pp. 1–16. doi: 10.1186/s13287-019-1203-3.
- Singh, V. K. et al. (2016) ‘Describing the Stem Cell Potency: The Various Methods of Functional Assessment and In silico Diagnostics.’, Frontiers in cell and developmental biology. Switzerland, 4, p. 134. doi: 10.3389/fcell.2016.00134.
- Stessuk, T. et al. (2016) ‘Platelet-rich plasma (PRP) and adipose-derived mesenchymal stem cells: stimulatory effects on proliferation and migration of fibroblasts and keratinocytes in vitro.’, Archives of dermatological research. Germany, 308(7), pp. 511–520. doi: 10.1007/s00403-016-1676-1.
- Tajima, S. et al. (2014) ‘Direct and Indirect Effects of a Combination of Adipose-Derived Stem Cells and Platelet-Rich Plasma on Bone Regeneration’, Tissue engineering. Part A, 21. doi: 10.1089/ten.TEA.2014.0336.
- Tantuway, V. et al. (2016) ‘Autologous grafting of non manipulated freshly isolated - adipose tissue derived stromal vascular fraction in single surgical sitting for treatment of knee osteoarthritis’, International Journal of Research in Orthopaedics, 3, p. 107. doi: 10.18203/issn.2455-4510.IntJResOrthop20164834.
- Tarnuzzer, R. W. et al. (1997) ‘Epidermal Growth Factor in Wound Healing: A Model for the Molecular Pathogenesis of Chronic Wounds.’, in Ziegler, T. R., Pierce, G. F., and Herndon, D. N. (eds) Growth Factors and Wound Healing. New York, NY: Springer New York, pp. 206–228.
- Tavares Pereira, D. dos S. et al. (2012) ‘Development of Animal Model for Studying Deep Second-Degree Thermal Burns’, Journal of Biomedicine and Biotechnology. Edited by M. Fedele. Hindawi Publishing Corporation, 2012, p. 460841. doi: 10.1155/2012/460841.
- Tohidnezhad, M. et al. (2011) ‘Platelet-released growth factors can accelerate tenocyte proliferation and activate the anti-oxidant response element’, Histochemistry and Cell Biology, 135(5), pp. 453–460. doi: 10.1007/s00418-011-0808-0.
- Toussaint, J. and Singer, A. J. (2014) ‘The evaluation and management of thermal

- injuries: 2014 update', Clinical and experimental emergency medicine. The Korean Society of Emergency Medicine, 1(1), pp. 8–18. doi: 10.15441/ceem.14.029.
- Tsien, L. (2006) 'Stem Cell Basics', Postgraduate Obstetrics & Gynecology, 26(24), pp. 1–6. doi: 10.1097/00256406-200612310-00001.
- Vaghardoost, R. et al. (2018) 'The Healing Effect of Sesame Oil, Camphor and Honey on Second Degree Burn Wounds in Rat.', World journal of plastic surgery, 7(1), pp. 67–71.
- Venturi, S. and Venturi, M. (2009) 'Iodine in Evolution of Salivary Glands and in Oral Health', Nutrition and health, 20, pp. 119–134. doi: 10.1177/026010600902000204.
- Widowati, W. and Widyanto, R. M. (2013) 'Sel Punca sebagai Transformasi Alternatif Terapi', Zenit, 2(1), pp. 1–5.
- Williams, C. (2011) 'Assessment and management of paediatric burn injuries.', Nursing standard (Royal College of Nursing (Great Britain): 1987). England, 25(25), pp. 60–64,66,68. doi: 10.7748/ns2011.02.25.25.60.c8349.
- Witte, M. B. and Barbul, A. (1997) 'General principles of wound healing.', The Surgical clinics of North America. United States, 77(3), pp. 509–528. doi: 10.1016/s0039-6109(05)70566-1.
- Wobus, A. M. and Boheler, K. R. (2005) 'Embryonic Stem Cells: Prospects for Developmental Biology and Cell Therapy', Physiological Reviews. American Physiological Society, 85(2), pp. 635–678. doi: 10.1152/physrev.00054.2003.
- Wong, A. et al. (2002) 'FRS2 alpha attenuates FGF receptor signaling by Grb2-mediated recruitment of the ubiquitin ligase Cbl.', Proceedings of the National Academy of Sciences of the United States of America. United States, 99(10), pp. 6684–6689. doi: 10.1073/pnas.052138899.
- Yun, Y.-R. et al. (2010) 'Fibroblast growth factors: biology, function, and application for tissue regeneration', Journal of tissue engineering. SAGE-Hindawi Access to Research, 2010, p. 218142. doi: 10.4061/2010/218142.
- Zakrzewski, W. et al. (2019) 'Stem cells: past, present, and future', Stem Cell Research & Therapy, 10(1), p. 68. doi: 10.1186/s13287-019-1165-5.
- Zuk, P. A. et al. (2002) 'Human adipose tissue is a source of multipotent stem cells', Molecular biology of the cell. The American Society for Cell Biology, 13(12), pp. 4279–4295. doi: 10.1091/mbc.e02-02-0105.