

Daftar Pustaka

- Abdallah, E. M. (2017). Black Seed (*Nigella sativa*) As Antimicrobial Drug: A Mini-Review. *Novel Approaches in Drug Designing and Development*, 3(2), 1–4. [https://www.semanticscholar.org/paper/Black-Seed \(Nigella-sativa\) As Antimicrobial-Drug](https://www.semanticscholar.org/paper/Black-Seed-(Nigella-sativa)-As-Antimicrobial-Drug)
- Amin, B., & Hosseinzadeh, H. (2016). Black Cumin (*Nigella sativa*) and Its Active Constituent, Thymoquinone: An Overview on the Analgesic and Anti-inflammatory Effects. In *Planta Medica* (Vol. 82, Issues 1–2, pp. 8–16). <https://doi.org/10.1055/s-0035-1557838>
- Arquam, M. (2021) ‘Pure honey and black seed (*Nigella sativa*) is the treatment and prevention of pandemic disease (corona virus)’, *Annals of the Romanian Society for Cell Biology*, 25(4), pp. 3359–3363.
- Arquam, M. (2021). Pure honey and black seed (*Nigella sativa*) is the treatment and prevention of pandemic disease (corona virus). *Annals of the Romanian Society for Cell Biology*, 25(4), 3359–3363.
- Asgharzadeh, F. et al. (2017) ‘Thymoquinone restores liver fibrosis and improves oxidative stress status in a lipopolysaccharide-induced inflammation model in rats.’, *Avicenna Journal of Phytomedicine*, 7(6), pp. 502–510.
- Beheshti, F. et al. (2014) ‘The effects of *Nigella sativa* extract on hypothyroidism-associated learning and memory impairment during neonatal and juvenile growth in rats’, *Nutritional Neuroscience*, 20(1), pp. 49–59. doi: 10.1179/1476830514Y.0000000144.

- Beheshti, F. et al. (2018) 'of Kidney Diseases and Transplantation Original Article Nigella sativa Prevented Liver and Renal Tissue Damage in', 29(3), pp. 554–566.
- Belgaumi, U. I., Patil, S., Gandhi, J. M., & Shete, A. S. (2020). The Many Therapeutic Applications of Nigella sativa - A Review of Literature. Journal of Evolution of Medical and Dental Sciences, 9(30), 2151–2157. <https://doi.org/10.14260/jemds/2020/469>
- Brown, G. T., & Kleiner, D. E. (2016). Histopathology of nonalcoholic fatty liver disease and nonalcoholic steatohepatitis. *Metabolism: clinical and experimental*, 65(8), 1080–1086. <https://doi.org/10.1016/j.metabol.2015.11.008>
- Collier, J. and Joekes, A. M. (1976) 'Severe allergic reaction to chlormethiazole', British Medical Journal, 2(6044), p. 1105. doi: 10.1136/bmj.2.6044.1105.
- Fajar, D. R., Akrom, & Darmawan, E. (2017). The influence of black cumin seed oil therapy with dosage of 1.5 mL/day and 3 mL/day to interleukin-21 (IL-21) expression of the patients with metabolic syndrome risk. IOP Conference Series: Materials Science and Engineering, 259(1). <https://doi.org/10.1088/1757-899X/259/1/012012>
- Garcia-Martin, F. et al. (1991) 'Acute interstitial nephritis induced by ethambutol [13]', Nephron, 59(4), pp. 679–680. doi: 10.1159/000186675.
- Hannan, Abdul, Aatur Rahman, A. A. M. S. 2 et al. (2021) 'Black Cumin (Nigella sativa L.): A Comprehensive Review on Phytochemistry, Health Benefits, Molecular Pharmacology, and Safety', Nutrients.

- Hannan, Abdul, Aatur Rahman, A. A. M. S. 2 et al. (2021) 'Black Cumin (*Nigella sativa* L.): A Comprehensive Review on Phytochemistry, Health Benefits, Molecular Pharmacology, and Safety', *Nutrients*.
- Hasan, D. I., Sabri, M., Hambal, M., & Balqis, U. (2019). Effect of Black Cumin Oil Administration on Cortisol Level and Liver Histopathology of Heat Stressed Broiler Chickens. *Jurnal Kedokteran Hewan*, 13(1), 31–36.
- Hosseini, S. M. et al. (2017) 'Protective effect of thymoquinone, the active constituent of *nigella sativa* fixed oil, against ethanol toxicity in rats', *Iranian Journal of Basic Medical Sciences*, 20(8), pp. 927–939. doi: 10.22038/ijbms.2017.9116.
- Hussein, M. M., Abdel-Azeem, A. S., & El-Damhougy, S. T. (2016). The health benefits of black seed (*Nigella sativa*). *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 7(1), 1109–1113.
- Islam, M. N. et al. (2021) 'Revisiting pharmacological potentials of *Nigella sativa* seed: A promising option for COVID-19 prevention and cure', *Phytotherapy Research*, 35(3), pp. 1329–1344. doi: 10.1002/ptr.6895.
- Jamel, E. G. and Ahmed, S. (2019) 'DRESS Syndrome and Chronic Renal Failure Induced by Ethambutol', *American Journal of the Medical Sciences*, 358(5), p. e19. doi: 10.1016/j.amjms.2019.06.003.
- Jaswal, A. et al. (2013) 'Therapeutic potential of thymoquinone against anti-tuberculosis drugs induced liver damage', *Environmental Toxicology and Pharmacology*, 36(3), pp. 779–786. doi: 10.1016/j.etap.2013.07.010.
- Jeong, I. et al. (2011) 'Drug Induced Hepatotoxicity of Antituberculosis Drugs and Their Serum Levels', *Chest*, 140(4), p. 771A. doi: 10.1378/chest.1116568.

- Kooti, W. et al. (2016) 'Phytochemistry, pharmacology, and therapeutic uses of black seed (*Nigella sativa*)', *Chinese Journal of Natural Medicines*, 14(10), pp. 732–745. doi: 10.1016/S1875-5364(16)30088-7.
- Kooti, W., Hasanzadeh-Noohi, Z., Sharafi-Ahvazi, N., Asadi-Samani, M., & Ashtary-Larky, D. (2016). Phytochemistry, pharmacology, and therapeutic uses of black seed (*Nigella sativa*). *Chinese Journal of Natural Medicines*, 14(10), 732–745. [https://doi.org/10.1016/S1875-5364\(16\)30088-7](https://doi.org/10.1016/S1875-5364(16)30088-7)
- Latif, M., Faheem, M., Asmatullah, Hoseinifar, S. H., & Van Doan, H. (2021). Dietary black seed effects on growth performance, proximate composition, antioxidant and histo-biochemical parameters of a culturable fish, rohu (*Labeo rohita*). *Animals*, 11(1), 1–14. <https://doi.org/10.3390/ani11010048>
- Li, X. et al. (2015) 'Liver transplantation in antituberculosis drugs-induced fulminant hepatic failure: A case report and review of the literature', *Medicine (United States)*, 94(49), pp. 1–4. doi: 10.1097/MD.0000000000001665.
- Makhlouf, H. A. et al. (2008) 'A prospective study of antituberculous drug-induced hepatotoxicity in an area endemic for liver diseases', *Hepatology International*, 2(3), pp. 353–360. doi: 10.1007/s12072-008-9085-y.
- Mariod, A. A., Saeed Mirghani, M. E., & Hussein, I. (2017). *Nigella sativa* L. Black Cumin. *Unconventional Oilseeds and Oil Sources*, 2019, 73–80. <https://doi.org/10.1016/b978-0-12-809435-8.00013-5>
- Nili-Ahmadabadi, A. et al. (2018) 'Thymoquinone attenuates hepatotoxicity and oxidative damage caused by diazinon: An in vivo study', *Research in Pharmaceutical Sciences*, 13(6), pp. 493–499. doi: 10.4103/1735-5362.245962.

- Nordin, A. et al. (2019) 'Effect of *Nigella sativa* and its bioactive compound on type 2 epithelial to mesenchymal transition: a systematic review', *BMC complementary and alternative medicine*, 19(1), p. 290. doi: 10.1186/s12906-019-2706-2.
- Perveen, R. (2019). Therapeutic effects of black cumin (*Nigella sativa*). A systematic review. *Progress in Nutrition*, 21, 40–49. <https://doi.org/10.23751/pn.v21i1-S.5796>
- Ramazan, I. et al. (2006) 'Antituberculous Therapy–Induced Fulminant Hepatic Failure: Successful Treatment With Liver Transplantation and Nonstandard Antituberculous Therapy Ramazan', *Liver Transplantation*, 12, pp. 1427–1430. doi: 10.1002/lt.
- Rohman, A. et al. (2019) 'Nigella sativa oil: Physico-chemical properties, authentication analysis and its antioxidant activity', *Food Research*, 3(6), pp. 628–634. doi: 10.26656/fr.2017.3(6).115.
- Sahu, N. et al. (2020) 'Naringenin mitigates antituberculosis drugs induced hepatic and renal injury in rats', *Journal of Traditional and Complementary Medicine*, 10(1), pp. 26–35. doi: 10.1016/j.jtcme.2019.01.001.
- Sallehuddin, N. et al. (2020) 'Nigella sativa and its active compound, thymoquinone, accelerate wound healing in an in vivo animal model: A comprehensive review', *International Journal of Environmental Research and Public Health*, 17(11), pp. 1–17. doi: 10.3390/ijerph17114160.
- Strunk, A. K. et al. (2016) 'Single- and multiple-dose pharmacokinetics of ethambutol and rifampicin in a tuberculosis patient with acute respiratory distress syndrome undergoing extended daily dialysis and ECMO treatment',

International Journal of Infectious Diseases, 42, pp. 1–3. doi: 10.1016/j.ijid.2015.10.018.

Subratti, A., Lalgee, L. J. and Jalsa, N. K. (2019) ‘Efficient extraction of black cumin (*Nigella sativa* L.) seed oil containing thymol, using liquefied dimethyl ether (DME)’, *Journal of Food Processing and Preservation*, 43(4), pp. 1–7. doi: 10.1111/jfpp.13913.

Yee, D. et al. (2003) ‘Incidence of Serious Side Effects from First-Line Antituberculosis Drugs among Patients Treated for Active Tuberculosis’, *American Journal of Respiratory and Critical Care Medicine*, 167(11), pp. 1472–1477. doi: 10.1164/rccm.200206-626OC.

Younossian, A. B. et al. (2005) ‘High hepatotoxicity of pyrazinamide and ethambutol for treatment of latent tuberculosis’, *European Respiratory Journal*, 26(3), pp. 462–464. doi: 10.1183/09031936.05.00006205.

(MD): National Institute of Diabetes and Digestive and Kidney Diseases; 2012-. Ethambutol. [Updated 2020 Dec 24]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK548745/>

Galli, S. J., Tsai, M., & Piliponsky, A. M. (2008). The development of allergic inflammation. *Nature*, 454(7203), 445–454. <https://doi.org/10.1038/nature07204>

Gunawan, B.K. and Kaplowitz, N. (2007) “Mechanisms of Drug-Induced Liver Disease,” *Clinics in Liver Disease*, pp. 459–475. Available at: <https://doi.org/10.1016/j.cld.2007.06.001>.

- Larrey, D. *et al.* (2020) “Drug induced liver injury and vascular liver disease,” *Clinics and Research in Hepatology and Gastroenterology*, 44(4), pp. 471–479. Available at: <https://doi.org/10.1016/j.clinre.2020.03.020>.
- Satapathy, S. K., Kuwajima, V., Nadelson, J., Atiq, O., & Sanyal, A. J. (2015). Drug-induced fatty liver disease: An overview of pathogenesis and management. *Annals of hepatology*, 14(6), 789–806. <https://doi.org/10.5604/16652681.1171749>
- Patel, V., & Sanyal, A. J. (2013). Drug-induced steatohepatitis. *Clinics in liver disease*, 17(4), 533–vii. <https://doi.org/10.1016/j.cld.2013.07.012>
- Lackner, C., Gogg-Kamerer, M., Zatloukal, K., Stumptner, C., Brunt, E. M., & Denk, H. (2008). Ballooned hepatocytes in steatohepatitis: the value of keratin immunohistochemistry for diagnosis. *Journal of hepatology*, 48(5), 821–828. <https://doi.org/10.1016/j.jhep.2008.01.026>
- Ethambutol Hydrochloride/Ethionamide 275* (no date). Available at: <http://aappolicy.aappublications.org/cgi/content/full/>.
- Rakhmawatie, M.D. *et al.* (2020) “Effect of essential oil of black cumin (*Nigella sativa*) on hematological profiles and total cholesterol levels of Wistar rats exposed by cigarette smoke,” *Indonesian Journal of Pharmacology and Therapy*, 1(1). Available at: <https://doi.org/10.22146/ijpther.495>.
- Takahashi, Y., & Fukusato, T. (2014). Histopathology of nonalcoholic fatty liver disease/nonalcoholic steatohepatitis. *World journal of gastroenterology*, 20(42), 15539–15548. <https://doi.org/10.3748/wjg.v20.i42.15539>

Gaut, J. P., & Liapis, H. (2020). Acute kidney injury pathology and pathophysiology: a retrospective review. *Clinical kidney journal*, 14(2), 526–536. <https://doi.org/10.1093/ckj/sfaa142>

Wang J, Liu J, Wu W, Yang S, Liu L, Fu Q, Li J, Chen X, Deng R, Wu C, Long S, Zhang W, Zhang H, Mao H and Chen W (2022) Combining Clinical Parameters and Acute Tubular Injury Grading Is Superior in Predicting the Prognosis of Deceased-Donor Kidney Transplantation: A 7-Year Observational Study. *Front. Immunol.* 13:912749. doi: 10.3389/fimmu.2022.912749