

DAFTAR PUSTAKA

- Adham M, AN K, AI M, et al. 2012 Nasopharyngeal carcinoma in Indonesia: epidemiology, incidence, signs, and symptoms at presentation. Chin J Cancer. doi:10.5732/cjc.011.10328.
- Alsaab, HO, Sau S, Alzhrani R. et.al. 2017. PD-1 and PD-L1 Checkpoint Signaling Inhibition for Cancer Immunotherapy : Mechanism, Combinations, and Clinical Outcome. Front. Pharmacol. 8 : 561. Doi : 10.3389/fphar. 2017.00561.
- Argirion I, Zarins KR, Julie J, et al. 2019. Ruterbusch Increasing Incidence of Epstein-Barr Virus-Related Nasopharyngeal Carcinoma in the United States. American Cancer Society 2019.
- Bachri, A. & Djufri, N.I., 2020. Retrospective Review of Nasofaring Carcinoma in Wahidin Sudirohusodo. International Journal of Nasopharyngeal Carcinoma (IJNPC) , 02, pp.82-84.
- Bintoro, S. 2015. Perbedaan eskresi hsa-miRNA-21-5p dan mRNA PTEN berdasarkan stadium klinis pada plasma darah pasien karsinoma nasofaring. Tesis dan Disertasi Universitas Gajah Mada
- Bruce, J.P., Liu F.F. 2014. MicroRNAs in nasopharyngeal carcinoma. Chinese journal of cancer. 33(11): 539-544
- Chan OSH, Kowanetz M,et al. 2017. Characterization of PD-L1 expression and immune cell infiltration in nasopharyngeal cancer. Oral Oncology 67 (2017) 52-60 57. 2017
- Chen, W. & Hu, G.-H. 2015. Biomarkers for enhancing the radiosensitivity of nasopharyngeal carcinoma. *Cancer Biol Med*, 12, 23-32.
- Chua DTT, 2014. Willam Ignace Wei. Nasopharyngeal Carcinoma. In: Johnson JT, Rosen CA, eds. Bailey's Head and Neck Surgery. Vol 5th ed. Philadelphia: Lippincot Williams & wilkins.1875-1897.
- Frigola X, Inman BA, Lohse CM, Krco CJ, Cheville JC, Houston TR, Leibovich B, Blute ML, Dong H and Kwon ED: Identification of a soluble form of B7-H1 that retains immuno-suppressive activity and is associated with aggressive renal cell carcinoma. Clin Cancer Res 17: 1915-1923, 2011.

GLOBOCAN. 2018. The global cancer observatory (WHO) International Agency for Research on Cancer 2018.p 2.

Green MR, Rodig S, Juszczynski P, et al. 2012. Constitutive AP-1 activity and EBV infection induce PD-L1 in Hodgkin lymphomas and posttransplant lymphoproliferative disorders: implications for targeted therapy. Clin Cancer Res. 2012;18:1611–8.

Gondhowiardjo SA, Handoko, Adham M,et al. 2020. Tumor microenvironment predicts local tumor extensiveness in PD-L1 positive nasopharyngeal cancer. PLoS ONE 15(3):e0230449. <https://doi.org/10.1371/journal.pone.0230449>

He, M.L., Luo M.X, Lin M.C., Kung H.F., 2012. MicroRNAs: potential diagnostic markers and therapeutic targets for EBV-associated nasopharyngeal carcinoma. Biochimica et biophysica,1-10

Hsu, M. C. et al. 2010. Increase of programmed death-1-expressing intratumoral CD8 T cells predicts a poor prognosis for nasopharyngeal carcinoma. Mod Pathol. 23, 1393–1403 (2010).

Hui A.B., lenarduzzi M., Krushel T., Waldron L., Pintilie M., Shi W., et.al., 2010. Comprehensive microRNA profiling for head and neck squamous cell carcinomas. American association for cancer research. 16 (4):1129-1139

Iliopoulos D., jaeger H.A., Hirsch M.L, Bulyk K. 2010. STAT3 activation of miRNA-21 and miR-181b-1, via PTEN and CYLD, are part of the epigenetic switch linking inflammation to cancer. Mol Cell (4) 39: 493-506

Kadir, A. ,Retnowati, K. , Akil, M.A. , Usman, A.N., 2018. The Evaluation of Serum IL-6 Changes as Proliferative Cytokines in Patients With Nasopharyngeal Carcinoma Before and After the Ionizing Radiotherapy. *Global Journal of Health Science*, 10(8), p. 55-6

Kataki AC, Simons MJ, Das AK, et al. 2011. Nasopharyngeal carcinoma in the northeastern states of india. Chin J Cancer 2011; 30(1): p.106-13.

Kim H.R, Ha S.J, Hong M.H, et al. 2016. PD-L1 expression on immune cells, but not on tumor cells, is a favorable prognostic factor for head and neck cancer patients. Scientific Reports.6: 36956.DOI : 10.1038/srep36956. 2016

Kolegium Ilmu Kesehatan THT-KL, Karsinoma Nasofaring, Modul Onkologi 2010.1-51.

Kwong D.L.W., Sham J.S.T., Au G.K.H. et al. 2004. Concurrent and Adjuvant Chemotherapy for Nasopharyngeal Carcinoma: A Factorial Study. Journal of Clinical Oncology. 22: 2643-53.

Kresno S.B., 2018. Imuno-Onkologi edisi ke-1. Sagung Seto. 119-120.

Kristianti, A., Madiadipoera, T. & Soeseno, B., 2010. Pengaruh sisplatin dosis tinggi terhadap penurunan. ORLI, 40(2), pp. 84-85.

Lee K.T., Tan J.K., Lam A.K, Gan S.Y., 2016. MicroRNAs serving as potential biomarkers and therapeutic targets in nasopharyngeal carcinoma: a critical review, Critical Reviews in Oncology/Hematology, 1-9.

Lee H.V, Lo, A.W, Leung, CY., 2016. Correlation of Expression of Tumor Cells with Survival Outcomes after Radical Intensity Modulated Radiation Therapy for Non-Metastatic Nasopharyngeal Carcinoma : a research article. Plos ONE 11 (6): e0157969. Doi : 10.371/journal.pone.0157969. 2016

Liu, N., Tang, L.L., Sun, Y., Cui, R.X., Wang, H.Y., Huang, B.J., et al., 2013. MiR-29 c suppresses invasion and metastasis by targeting TIAM1 in nasopharyngeal carcinoma. Cancer Lett. 329, 181–188.

Liu X, Shan C, Song Y, Du J. 2019. Prognostic Value of Programmed Death Ligand-1 Expression in Nasopharyngeal Carcinoma : A Meta-Analysis of 1.315 Patients. Front. Oncol. 9:1111. Doi :10.3389/fonc.2019.01111

Lin, S., HO; Willard, Fee; Francisco, Talavera. 2016. Malignant Nasopharyngeal Tumors. <http://emedicine.medscape.com/article/848163-overview>. [Accessed 24/4/2020].

Li Y., Yan L., Zhang W., Wang H., Chen W., Hu N., et.al. 2014. miRNA-21 inhibitor suppresses proliferation and migration of nasopharyngeal carcinoma cells through down-regulation of BCL2 expression. Int J Clin Exp Pathol. 7(6):3478-3487.

Li YF, Ding JW, Liao LM, Zhang ZL, Liao SS, Wu Y, Zhou DY, Liu AW and Huang L. 2017. Expression of programmed death ligand-1 predicts poor outcome in nasopharyngeal carcinoma. Mol Clin Oncol 7: 378-382, 2017.

- Lo SS, Lu JJ. 2010. Natural history, presenting symptoms, and diagnosis of nasopharyngeal carcinoma. In: Brady LW, Heilmann H-P, Molls M, Nieder C, eds. Nasopharyngeal Cancer Multidisciplinary Management. Vol Germany: Springer. 41-52.
- Luan J., Wang J., Su Q., Chen X., jiang G., Xu X., 2015. Meta –analysis of the differentially expressed microRNA profiles in nasopharyngeal carcinoma. *Oncotarget*. 7 (9). 10513-10521.
- Melani W, Sofyan F. 2013. Karakteristik penderita kanker nasofaring di Rumah Sakit H . Adam Malik Medan tahun 2011. *E-Jurnal FK-USU*, 1(1):1-5
- Miao B.P., Zhang R.S., Fu Y.T., Zhao M., Liu Z.G, Yang P.C., 2014. Nasopharyngeal cancer-derived microRNA-21 promotes immune suppressive B cells. *Cellular & Molecular Immunology*, 1–7.
- National Comprehensive Cancer Network Guidelines for head and neck cancer (version 4.2020)
- Pardoll DM. 2012. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer*. 2012; 12(4): 252–64. doi: [10.1038/nrc3239](https://doi.org/10.1038/nrc3239) PMID: [22437870](https://pubmed.ncbi.nlm.nih.gov/22437870/)
- Perez CA, Brady LW, Wazer DE, Halperin EC. 2013. Principles and Practice of Radiation Oncology Sixth Edition. Lippincott Willian. America
- Pieter N.A.L.,2013. Profil Iga(Vca-P18+Ebna1) Dan Viral Load DNA EBV Sebagai Faktor Resiko Keluarga Penderita Karsinoma Nasofaring Dengan EBV Positif. Disertasi tidak diterbitkan. Program Pasca Sarjana UNHAS. 1-50.
- Plant, L., R 2009. Neoplasms of the Nasopharynx. In: SNOW, J., B; WACKIM,P,A (ed.) *Ballenger's Otorhinolaryngology Head and Neck Surgery*. 17 ed. Shelton, Connecticut: BC Decker.
- Punagi, A.Q., Mubarika. S., Yususuf. I. 2013. Vascular Endothelial Growth Factor Receptor (VEGFR Flt-4) and Latent Membrane Protein (LMP-1) Expression in Nasopharyngeal Carcinoma. *The Open Otorhinolaryngology Journal*, Volume 7, p. 1.

Putri EB. 2011. Karakteristik penderita karsinoma nasofaring di departemen ilmu kesehatan THT-KL FKUP/RSUP Dr. Hasan Sadikin Bandung Periode Tahun 2006 - 2010 [skripsi]. Bandung: Universitas Padjadjaran.

Ran X, Yang K. 2018. Inhibitors of the PD-1/PD-L1 Axis for the Treatment of Head and Neck Cancer : Current status and Future Prospective in Drug Design, Development and Therapy 2017:11 2007-2014.
<https://www.doverpress.com/by76:102.64.113on30-Jul.2018>

Registrasi Kanker Nasional, Indonesia, Kementerian Kesehatan Republik Indonesia. 2019.

Roezin A, Adham M. 2014. Karsinoma Nasofaring. Dalam: Soepardi EA, Iskandar N, Bashirudin J, Restuti RD. Buku ajar ilmu kesehatan telinga hidung tenggorokan. Edisi ke-7. Jakarta: Badan Penerbit FKUI. hlm. 158–63.

Sabarin MSM, Permana AD, Soeseno B. 2014. Epidemiologi penderita tumor ganas kepala leher di Departemen telinga hidung tenggorok-Kepala leher rumah sakit Dr. Hasan Sadikin Bandung, Indonesia, periode 2010-2014. p 2-3.

Salehiniya H, Mohammadian M., Mohammadian Hafshejani A, et al. 2018. Nasopharyngeal Cancer In The World: Epidemiology, Incidence, Mortality And Risk Factors. World cancer research journal WCRJ 2018; 5 (1): e1046.

Salmaninejad A, Valilou AF, Shabgah, AG, et.al. 2019. PD-1/PD-L1 pathway : Basic biology and role in cancer immunotherapy. J Cell Physiol. 2019; 1-14. <https://doi.org/10.1002/jcp.28358>

Savitri E., Haryana MS. (2015). Expression of interleukin-8, interleukin-10 and Ebstein-Barr Viral Load as prognostic indicator in nasopharyngeal carcinoma. Global journal of Health science.Vol 7(3) 362-372.

Sekar M. 2018. Perbedaan Kualitas Hidup Pasien Karsinoma Nasofaring Dengan Karakteristik (Usia, Jenis Kelamin Dan Stadium) Tumor Nasofaring Di RSUD Dr. H. Abdoel Moeloek Bandar Lampung [skripsi] Lampung: Universitas Lampung; 2018.

Sengupta, S., den Boon, J.A., Chen, I.H., Newton, M.A., Stanhope, S.A., Cheng, Y.J.,et al., 2008. MicroRNA 29c is down-regulated in nasopharyngeal carcinomas, up-regulating mRNAs encoding

- extracellular matrix proteins. Proc. Natl. Acad.Sci. U. S. A. 105, 5874–5878.
- Shah, J., Patel, S., Singh, B. & Wong, R., 2020. Head And Neck Surgery And Oncology. Fifth Edition ed. London: Elsevier.
- Sharif S.E, Zawawi N, Yajid A.I, et al. 2020. Pathology classification of nasopharyngeal carcinoma. DOI: <https://doi.org/10.1016/B978-0-12-814403-9.00005-7>
- Shi SJ, Wang LJ, Wang GD, Guo ZY, Wei M, Meng YL, Yang AG and Wen WH.2013. B7-H1 expression is associated with poor prognosis in colorectal carcinoma and regulates the proliferation and invasion of HCT116 colorectal cancer cells. PLoS One 8: e76012, 2013.
- Spence T., Bruce J., Yip K.W., Liu F.F., 2015. MicroRNAs in nasopharyngeal carcinoma. Chin clin Oncol, 5(2).1-11.
- Sun Z.F., Li S.G., Kaufmann A.M., Albers A.L. 2014., miRNA-21 increases the programmed cell death 4 gene-regulated cell proliferation in head and neck squamous carcinoma cell lines. Oncology Reports 32: 2283-2289.
- Tan G., Tang X., tang F., 2015. The role of microRNAs in nasopharyngeal carcinoma. Tumor Biol. 69-79.
- Tsao SW, Yip YL, Tsang CM, Pang PS, Lau VMY, Zhang G, Lo KW. 2014. Etiological factors of nasopharyngeal carcinoma. Oral Oncol, 2014;50:330-8 [SEP]
- Wang Z, Peng SH, Xie H, Guo LP, Cai QL, Shang ZQ, et al. 2018. Prognostic and clinicopathological significance of PD-L1 in patients with renal cell carcinoma: a meta-analysis based on 1863 individuals. Clin Exp Med. (2018) 18:165–75. doi: 10.1007/s10238-018-0488-3
- Wargasetia TL.,2016. The potential of miRNAs as biomarker and therapy targets for cancer. Journal of medicine and health. 1, 277-286.
- Wotman M, Herman SW, et al. 2020. The prognostic Role of Programmed Death-Ligand 1 in Nasopharyngeal Carcinoma. Laryngoscope, 00:1-9, 2020.

- Zhang J.X., Qian D., Wang F.W., Liao D.Z., Wei J.H., Tong Z.T., et.al. 2013. MicroRNA-29c enhances the sensitivities of human nasopharyngeal carcinoma to cisplatin-based chemotherapy and radiotherapy. *Cancer Letters.* 91-98.
- Zerde, I., et al. 2018. Genetic, transcriptional and post-Translational Regulation of the Programmed Death Protein Ligand 1 in Cancer: Biology and Clinical Correlation.<https://doi.org/10.1038/s41388-018-0303-3>
- Zhang J., Fang W., Qin T., Yang Y., Hong S., Liang W., et.al. 2015. Co-expression of PD-1 and PD-L1 predict poor outcome in nasopharyngeal carcinoma. *Med Oncol* (2015) 32:86. DOI 10.1007/s12032-015-0501-6
- Zheng L., Cao C., Cheng G., Hu Q., Chen X. 2017. Cytomembranic PD-L1 expression in locoregionally advanced nasopharyngeal carcinoma. *Onco Targets and Therapy* 2017 : 10 5483-587.
- Zhou Y., Shi D., Miao J., Wu H., Chen J., Chen J., et.al. 2017. PD-L1 predict poor prognosis for nasopharyngeal carcinoma irrespective of PD-1 and EBV-DNA load. *Scientific Reports.* 7: 43627. DOI: 10.1038/srep43627
- Zhu Q, Cai M, Chen CL, et al. 2017. Tumor cells PD-L1 expression as a favorable prognosis factor in nasopharyngeal carcinoma patients with pre-existing intratumor- infiltrating lymphocytes. *OncolImmunol*, DOI : 10.1080/2162402X.2017.1312240
- .

Lampiran 1. Lembar informed consent.

FORMULIR PERSETUJUAN SETELAH PENJELASAN

Yang bertanda tangan dibawah ini :

Nama :
Umur :
Jenis Kelamin :
Alamat :
Pekerjaan :

Setelah mendapatkan penjelasan dari peneliti dengan ini saya menyatakan bersedia secara sukarela tanpa paksaan untuk mengikuti penelitian ini dan mentaati semua prosedur yang akan dilakukan pada penelitian ini.

Saya tahu bahwa saya berhak untuk bertanya apabila masih ada hal-hal yang saya tidak mengerti.

Saya mengerti bahwa prosedur pengambilan jaringan tumor nasofaring untuk pemeriksaan Programmed Death Ligand 1 terhadap diri saya berlangsung kurang lebih 15 menit dan dapat menyebabkan hal-hal yang merugikan, namun saya percaya bahwa akan dilakukan tindakan-tindakan kewaspadaan untuk mencegah hal-hal tersebut. Saya mengerti bahwa resiko yang mungkin terjadi dalam pengambilan jaringan tumor nasofaring adalah rasa tidak nyaman dan nyeri dan saya juga mengerti bahwa resiko pengambilan jaringan tumor nasofaring di ruang operasi adalah resiko terjadinya perdarahan. Saya juga telah mengerti bahwa resiko ini dapat diantisipasi dengan cara pengambilan jaringan tumor nasofaring secara legeartis dan anestesi lokal dan umum, serta adanya tampon hidung dan obat obatan anti nyeri dan anti perdarahan. Saya juga telah mengerti bahwa hasil pemeriksaan yang didapatkan akan dijamin kerahasiaannya dan semua biaya pemeriksaan adalah tanggungan sepenuhnya dari peneliti.

Saya juga tahu bahwa saya berhak mendapat penanganan atau kompensasi biaya dari peneliti bila terjadi efek samping dari pemeriksaan yang dilakukan.

Saya juga berhak menolak untuk tidak ikut dalam penelitian ini tanpa kehilangan hak saya untuk mendapat pelayanan kesehatan dari dokter.

Makassar

Nama Saksi

1. (.....)
Nama jelas/Tanda tangan

2. (.....) (.....)
Nama jelas/Tanda tangan Nama jelas/Tanda tangan

Peneliti Utama : **dr. Mila Habibasari**

Komplek Graha Sari Blok C/2, Makassar / Telpon : 082291590001

Dokter Penanggung jawab medis : **Prof. Dr.dr. Abdul Kadir, Ph. D, Sp.T.H.T.K.L(K),M.Kes**
Hertasning Blok E14/11, Makassar/Telepon: 0816 253 509

Lampiran 2. Data Dasar Penelitian

DATA SAMPEL PENELITIAN

NO	NAMA	UMUR (THN)	JK (L/P)	HASIL BIOPSI	DIAGNOSA KLINIS	PD-L1
1.	Tn.GD	40	L	Squamous Cell Carcinoma tidak berkeratin tidak berdiferensiasi	KNF Stadium IVB (T4N0M0)	LOW
2.	Ny.NM		P	Squamous Cell Carcinoma tidak berkeratin tidak berdiferensiasi	KNF Stadium IVC (T4N3M1)	LOW
3.	Ny.R		P	Squamous Cell Carcinoma	KNF Stadium III (T3N1M0)	LOW
4.	Tn.ZB		L	Squamous Cell Carcinoma, WHO tipe II	KNF Stadium II (T2N0M0)	NEG/0
5.	Tn.A		L	Squamous Cell Carcinoma	KNF Stadium II (T2N0M0)	LOW
6.	Tn.OL		L	Squamous Cell Carcinoma, WHO tipe II	KNF Stadium II (T2N0M0)	NEG/0
7.	Ny.BW		P	Squamous Cell Carcinoma	KNF Stadium IVA (T4N0M0)	HIGH
8.	Ny.M		P	Squamous Cell Carcinoma tidak berkeratin tidak berdiferensiasi	KNF Stadium III (T3N1M0)	LOW
9.	Tn.RA		L	Squamous Cell Carcinoma	KNF Stadium II (T2N0M0)	LOW
10.	Tn.SP		L	Squamous Cell Carcinoma	KNF Stadium IVA (T4N3M0)	LOW
11.	Tn.DM		L	Squamous Cell Carcinoma	KNF Stadium IVB (T4N3M1)	HIGH
12.	Tn.TW		L	Squamous Cell Carcinoma, WHO tipe II	KNF Stadium II (T2N0M0)	LOW

13.	Tn.OLH		L	Squamous Cell Carcinoma tidak berkeratin	KNF Stadium II (T2N0M0)	NEG/0
14.	Tn.SM		L	Squamous Cell Carcinoma	KNF Stadium IVB (T4N2M1)	HIGH
15.	Ny.MS		P	Squamous Cell Carcinoma	KNF Stadium IVA (T4N0M0)	HIGH
16.	Nn.ML		P	Squamous Cell Carcinoma	KNF Stadium IVA (T4N0M0)	HIGH
17.	Tn.TL		L	Squamous Cell Carcinoma	KNF Stadium III (T3N0M0)	LOW
18.	Tn.MYN		L	Squamous Cell Carcinoma	KNF Stadium IVA (T3N1M0)	HIGH
19.	Tn.YSF		L	Squamous Cell Carcinoma	KNF Stadium IVA (T3N1M0)	HIGH
20.	Tn.IW		L	Squamous Cell Carcinoma	KNF Stadium III (T3N0M0)	LOW
21.	Tn.WH		L	Squamous Cell Carcinoma	KNF Stadium III (T3N1M0)	HIGH
22.	Tn.SK		L	Squamous Cell Carcinoma	KNF Stadium III (T3N1M0)	LOW
23.	Tn.MR		L	Squamous Cell Carcinoma	KNF Stadium II (T2N0M0)	LOW
24.	Tn.LM		L	Squamous Cell Carcinoma	KNF Stadium IVA (T4N1M0)	HIGH
25	Tn.NM		L	Squamous Cell Carcinoma	KNF Stadium IVA (T4N1M0)	HIGH

	no	jk	umur	stadium	stadium	pdl1	VAR000	VAR000
1	1	1	40 IVB		1	1	1	3
2	2	2	34 IVC		1	1	1	3
3	3	2	39 III		1	1	1	3
4	4	1	35 II		2	0	2	3
5	5	1	37 II		2	1	1	3
6	6	1	41 II		2	0	2	3
7	7	2	41 IVA		1	2	1	2
8	8	2	48 III		1	1	1	3
9	9	1	56 II		2	1	1	3
10	10	2	51 IVA		1	1	1	3
11	11	1	61 IVB		1	2	1	2
12	12	1	58 II		2	1	1	3
13	13	1	49 II		2	0	2	3
14	14	1	49 IVB		1	2	1	2
15	15	2	30 IVA		1	2	1	2
16	16	2	24 IVA		1	2	1	2
17	17	1	45 III		1	1	1	3
18	18	1	45 IVA		1	2	1	2
19	19	1	51 IVA		1	2	1	2
20	20	1	45 III		1	1	1	3
21	21	2	54 III		1	2	1	2
22	22	1	54 III		1	1	1	3
23	23	1	26 II		2	1	1	3
24	24	1	46 IVA		1	2	1	2
25	25	1	46 IVA		1	2	1	2

Frequencies

Statistics

	Jenis Kelamin	Stadium Kanker	Stadium Kanker	PD-L1
N	Valid	25	25	25
	Missing	0	0	0

Frequency Table

Jenis Kelamin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Laki-laki	17	68.0	68.0	68.0
	Perempuan	8	32.0	32.0	100.0
	Total	25	100.0	100.0	

Stadium Kanker

			Valid	Cumulative
Valid	II III	7	28.0	28.0
	IVA	6	24.0	52.0
	IVB	8	32.0	84.0
	IVC	3	12.0	96.0
	Total			

Stadium Kanker

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid Lanjut (III/IV)		18	72.0	72.0	72.0
Dini (I/II)		7	28.0	28.0	100.0
Total		25	100.0	100.0	

PD-L1

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Negative	3	12.0	12.0	12.0
	Low	12	48.0	48.0	60.0
	High	10	40.0	40.0	100.0
	Total	25	100.0	100.0	

Means

Case Processing Summary

	Cases					
	Included		Excluded		Total	
	N	Percent	N	Percent	N	Percent
Umur (Tahun) * Stadium Kanker	25	100.0%	0	0.0%	25	100.0%

Report

Umur (Tahun)

Stadium Kanker	Mean	N	Std. Deviation	Median	Minimum	Maximum
Lanjut (III/IV)	44.61	18	9.004	45.50	24	61
Dini (I/II)	43.14	7	11.711	41.00	26	58
Total	44.20	25	9.600	45.00	24	61

T-Test

Group Statistics

Stadium Kanker		N	Mean	Std. Deviation	Std. Error Mean
Umur (Tahun)	Lanjut (III/IV)	18	44.61	9.004	2.122
	Dini (I/II)	7	43.14	11.711	4.426

Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means		
		F	Sig.	t	df	Sig. (2-tailed)
Umur (Tahun)	Equal variances assumed	1.328	.261	.337	23	.739
	Equal variances not assumed			.299	8.910	.772

Independent Samples Test

		t-test for Equality of Means					
		C a			95% Confidence Interval of the Difference		
	V a	Miss ing	T o	Mean Difference	Std. Error Difference	Lower	Upper
	N	Per cent	N				
	Jenis Kelamin *	25	100. 0%	0.	2	100.0 %	
Stadium Kanker							
Umur (Tahun)	Equal variances assumed			1.468	4.358	-7.546	10.483
	Equal variances not assumed			1.468	4.909	-9.653	12.590

Crosstabs

Case Processing Summary

Jenis Kelamin * Stadium Kanker Crosstabulation

Jenis Kelamin	Laki-laki		Stadium Kanker		
			Lanjut (III/IV)	Dini (I/II)	
Jenis Kelamin	Laki-laki	Count	10	7	17
		% within Jenis Kelamin	58.8%	41.2%	100.0%
		% within Stadium Kanker	55.6%	100.0%	68.0%
	Perempuan	Count	8	0	8
		% within Jenis Kelamin	100.0%	0.0%	100.0%
		% within Stadium Kanker	44.4%	0.0%	32.0%
Total		Count	18	7	25
		% within Jenis Kelamin	72.0%	28.0%	100.0%
		% within Stadium Kanker	100.0%	100.0%	100.0%

Chi-Square Tests

			Asymp. Sig. (2- sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.575 ^a	1	.032		
Continuity Correction ^b	2.761	1	.097		
Likelihood Ratio Fisher's Exact Test	6.613	1	.010		
Linear-by-Linear Association				.057	.040
N of Valid Cases					

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 2.24.

b. Computed only for a 2x2 table

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
PD-L1 * Stadium Kanker	25	100.0%	0	0.0%	25	100.0%

PD-L1 * Stadium Kanker Crosstabulation

	PD-L1	Negative	Stadium Kanker		
			Count	Lanjut	
			0	3	3
	% within PD-L1		0.0%	100.0%	100.0%

Low	Count	8	4	12
	% within PD-L1	66.7%	33.3%	100.0%
High	Count	10	0	10
	% within PD-L1	100.0%	0.0%	100.0%
Total	Count	18	7	25
	% within PD-L1	72.0%	28.0%	100.0%

Chi-Square Tests

			Asymp. Sig. (2- sided)
Pearson Chi-Square	11.772 ^a	2	.003
Likelihood Ratio	14.371	2	.001
Linear-by-Linear Association	10.611	1	.001

a. 4 cells (66.7%) have expected count less than 5. The minimum expected count is .84.

NPar Tests

Kruskal-Wallis Test

Ranks

Stadium Kanker		N	Mean Rank
PD-L1	Lanjut (III/IV) Dini (I/II)	18	15.61
	Total	7	6.29
		25	

Test Statistics^{a,b}

	PD-L1
Chi-Square df	9.808
Asymp. Sig.	1

a. Kruskal Wallis Test

b. Grouping Variable: Stadium Kanker


ROC Curve

Case Processing Summary

	Valid N (listwise)
Positive ^a	18

Larger values of the test result variable(s) indicate stronger evidence for a positive actual state.

a. The positive actual state is Lanjut (III/IV).



Diagonal segments are produced by ties.

Area Under the Curve

Test Result Variable(s): PD-L1

		Asymptotic Sig. ^b	Asymptotic 95% Confidence Interval	
			Lower Bound	Upper Bound
.873	.073	.004	.730	1.000

The test result variable(s): PD-L1 has at least one tie between the positive actual state group and the negative actual state group. Statistics may be biased.

a. Under the nonparametric assumption

b. Null hypothesis: true area = 0.5

Coordinates of the Curve

Test Result Variable(s): PD-L1

Positive if Greater Than or Equal To ^a		
-1.00	1.000	1.000
.50	1.000	.571

The test result variable(s): PD-L1 has at least one tie between the positive actual state group and the negative actual state group.

a. The smallest cutoff value is the minimum observed test value minus 1, and the largest cutoff value is

the maximum observed test value plus 1. All the other cutoff values are the averages of two consecutive ordered observed test values.

Crosstab

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
PD-L1 * Stadium Kanker	25	100.0%	0	0.0%	25	100.0%
PD-L1 * Stadium Kanker	25	100.0%	0	0.0%	25	100.0%

PD-L1 * Stadium Kanker

Crosstab

			Stadium Kanker		Total
			Lanjut (III/IV)	Dini (I/II)	
PD-L1	Low/High	Count	18	4	22
		% within PD-L1	81.8%	18.2%	100.0%
		% within Stadium Kanker	100.0%	57.1%	88.0%
	Negative	Count	0	3	3
		% within PD-L1	0.0%	100.0%	100.0%
		% within Stadium Kanker	0.0%	42.9%	12.0%
	Total	Count	18	7	25
		% within PD-L1	72.0%	28.0%	100.0%
		% within Stadium Kanker	100.0%	100.0%	100.0%

Chi-Square Tests

			Asymp. Sig. (2- sided)	Exact Sig.	Exact Sig.
Pearson Chi-Square	8.766 ^a	1	.003		
Continuity Correction ^b					
Likelihood Ratio	5.178	1	.023		
Fisher's Exact Test					
Linear-by-Linear Association	8.786	1	.003		

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is .84.

b. Computed only for a 2x2 table

PD-L1 Stadium Kanker

Crosstab

			Stadium Kanker		
			Lanjut (III/IV)	Dini (I/II)	
PD-L1	High	Count	10	0	10
		% within PD-L1	100.0%	0.0%	100.0%
		% within Stadium Kanker	55.6%	0.0%	40.0%
	Negative/Low	Count	8	7	15
		% within PD-L1	53.3%	46.7%	100.0%
		% within Stadium Kanker	44.4%	100.0%	60.0%
Total	Count	18	7	25	
	% within PD-L1	72.0%	28.0%	100.0%	
	% within Stadium Kanker	100.0%	100.0%	100.0%	


Chi-Square Tests

			Asymp. Sig. (2- sided)	Exact Sig.	Exact Sig.
Pearson Chi-Square	6.481 ^a	1	.011		
Continuity Correction ^b	4.373	1	.037		
Likelihood Ratio					
Fisher's Exact Test					
Linear-by-Linear Association	8.920	1	.003		


a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 2.80.

b. Computed only for a 2x2 table

Graph



Lampiran 3. Rekomendasi Persetujuan Etik



REKOMENDASI PERSETUJUAN ETIK

Nomor : 450/UN4.6.4.5.31/ PP36/ 2021

Tanggal: 12 Juli 2021

Dengan ini Menyatakan bahwa Protokol dan Dokumen yang Berhubungan Dengan Protokol berikut ini telah mendapatkan Persetujuan Etik :

No Protokol	UH21030118	No Sponsor Protokol	
Peneliti Utama	dr. Mila Habibasari	Sponsor	
Judul Peneliti	Ekspresi kadar PD-L1 sebagai penanda prognosis pada pasien karsinoma nasofaring		
No Versi Protokol	2	Tanggal Versi	6 Juli 2021
No Versi PSP	2	Tanggal Versi	6 Juli 2021
Tempat Penelitian	RS Dr. Wahidin Sudirohusodo Makassar		
Jenis Review	<input type="checkbox"/> Exempted <input checked="" type="checkbox"/> Expedited <input type="checkbox"/> Fullboard Tanggal	Masa Berlaku 12 Juli 2021 sampai 12 Juli 2022	Frekuensi review lanjutan
Ketua Komisi Etik Penelitian Kesehatan FKUH	Nama Prof.Dr.dr. Suryani As'ad, M.Sc.,Sp.GK (K)		
Sekretaris Komisi Etik Penelitian Kesehatan FKUH	Nama dr. Agussalim Bukhari, M.Med.,Ph.D.,Sp.GK (K)		

Kewajiban Peneliti Utama:

- Menyerahkan Amandemen Protokol untuk persetujuan sebelum di implementasikan
- Menyerahkan Laporan SAE ke Komisi Etik dalam 24 Jam dan dilengkapi dalam 7 hari dan Lapor SUSAR dalam 72 jam setelah Peneliti Utama menerima laporan
- Menyerahkan Laporan Kemajuan (progress report) setiap 6 bulan untuk penelitian resiko tinggi dan setiap setahun untuk penelitian resiko rendah
- Menyerahkan laporan akhir setelah Penelitian berakhir
- Melaporkan penyimpangan dari protokol yang disetujui (protocol deviation / violation)
- Mematuhi semua peraturan yang ditentukan

Lampiran 5. Dokumentasi Penelitian



