

DAFTAR PUSTAKA

1. Hooper N, Armstrong TJ. Hemorrhagic Shock. [Updated 2020 Nov 21]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK470382/>
2. Taghavi S, Askari R. Hypovolemic Shock. [Updated 2020 Jul 20]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK513297/>
3. Martel, M.-J. No. 115-Hemorrhagic Shock. *Journal of Obstetrics and Gynaecology Canada*. 2018. 40(12), e874–e882. doi:10.1016/j.jogc.2018.10.004
4. Abbas, Kun Arifi. Fluid Resuscitation in Trauma. *Indonesian Journal of Anesthesiology and Reanimation*. Volume 1 Number 2. 2019 : 52-57
5. Merrick, Claire et al. *Advanced Trauma Life Support*. 2018 . American College of Surgeons 10 th edition.
6. Rossaint R, Bouillon B, Cerny V, et al. The European guideline on management of major bleeding and coagulopathy following trauma: fourth edition. *Crit Care*. 2016;20:100. Published 2016 Apr 12. doi:10.1186/s13054-016-1265-x
7. Van der Mullen J, et al. Assessment of hypovolaemia in the critically ill. *Anaesthesiol Intensive Ther*. 2018;50(2):141-149. [[PubMed](#)] [[Reference list](#)]
8. Hernández, Glenn et al. “Statistical analysis plan for early goal-directed therapy using a physiological holistic view - the ANDROMEDA-SHOCK: a randomized controlled trial.” “Plano de análise estatística para o estudo do tratamento precoce baseado em metas com utilização de uma visão fisiológica holística - estudo ANDROMEDA-SHOCK: um estudo randomizado e controlado.” *Revista Brasileira de terapia intensiva* vol. 30,3 (2018): 253-263. doi:10.5935/0103-507X.20180041

9. Van Genderen, et al. *Early Peripheral Perfusion–guided Fluid Therapy in Patients with Septic Shock. American Journal of Respiratory and Critical Care Medicine.* 2015 *191*(4), 477–480.). doi:10.1164/rccm.201408-1575le
10. Castro R, Kattan E, Ferri G, et al. Effects of capillary refill time-vs. lactate-targeted fluid resuscitation on regional, microcirculatory and hypoxia-related perfusion parameters in septic shock: a randomized controlled trial. *Ann Intensive Care.* 2020;10(1):150. Published 2020 Nov 2. doi:10.1186/s13613-020-00767-4
11. Lauralee Sherwood. *Introduction to Human Physiology* 8th edition. 2013.
12. Cohen, Maya & Monaghan, Sean. (2021). Hemorrhagic shock and fluid dynamics. *Physiological Reports.* 9. 10.14814/phy2.14813.
13. Bonanno FG. Hemorrhagic shock: The "physiology approach". *J Emerg Trauma Shock.* 2012;5(4):285-295. doi:10.4103/0974-2700.102357
14. Krausz MM. Initial resuscitation of hemorrhagic shock. *World J Emerg Surg.* 2006;1:14. Published 2006 Apr 27. doi:10.1186/1749-7922-1-14
15. McGuire D, Gotlib A, King J. Capillary Refill Time. [Updated 2020 Apr 28]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK557753/>
16. Pickard A, Karlen W, Ansermino JM. Capillary refill time: is it still a useful clinical sign? *Anesth Analg.* 2011 Jul;113(1):120-3. [[PubMed](#)] [[Reference list](#)]
17. Liu C, Correia R, Ballaji H, Korposh S, Hayes-Gill B, Morgan S. Optical Fibre Sensor for Simultaneous Measurement of Capillary Refill Time and Contact Pressure. *Sensors (Basel).* 2020 Mar 03;20(5) [[PMC free article](#)] [[PubMed](#)] [[Reference list](#)]
18. Jacquet-Lagrèze, M., Bouhamri, N., Portran, P. *et al.* Capillary refill time variation induced by passive leg raising predicts capillary refill time response to volume expansion. *Crit Care* **23**, 281 (2019). <https://doi.org/10.1186/s13054-019-2560-0>
19. Pickard A, Karlen W, Ansermino JM. Capillary refill time: is it still a useful clinical sign?. *Anesthesia & Analgesia.* 2011 Jul 1;113(1):120-3

20. Andersen LW, Mackenhauer J, Roberts JC, Berg KM, Cocchi MN, Donnino MW. Etiology and therapeutic approach to elevated lactate levels. *Mayo Clin Proc.* 2013;88(10):1127-1140. doi:10.1016/j.mayocp.2013.06.012
21. Bakker J, Nijsten MW, Jansen TC. Clinical use of lactate monitoring in critically ill patients. *Ann Intensive Care.* 2013;3(1):12. Published 2013 May 10. doi:10.1186/2110-5820-3-12
22. Jevon P, Gallier H (2020) How to measure capillary refill time in patients who are acutely ill. *Nursing Times* [online]; 116: 8, 29-30
23. Susannah Fleming, Peter J Gill, Ann Van den Bruel and Matthew Thompson. *British Journal of General Practice* 2016; 66 (652): 587-588. DOI: <https://doi.org/10.3399/bjgp16X687925>
24. King, D., Morton, R., & Bevan, C. (2013). *How to use capillary refill time. Archives of Disease in Childhood - Education & Practice Edition, 99(3), 111–116.* doi:10.1136/archdischild-2013-305198
25. Paul L. Marino. Marino's The ICU Book 4th edition. 2014. Wolters Kluwer Health/Lippincott Williams & Wilkins
26. Réminiac F, Saint-Etienne C, Runge I, Ayé DY, Benzekri-Lefevre D, Mathonnet A, Boulain T: Apakah laktat vena sentral dan laktat arteri dapat dipertukarkan? Sebuah studi retrospektif manusia. *Anal anestesi.* 2012; 115:605–10
27. Zitek T, Skaggs ZD, Rahbar A, Patel J, Khan M: Apakah larutan Ringer laktat intravena meningkatkan serum laktat? *J Emerg Med.* 2018; 55:313–8
28. Smith ZR, Horng M, Rech MA: Obat-diinduksi hiperlaktatemia dan asidosis laktat: Sebuah tinjauan sistematis literatur. *Farmakoterapi.* 2019; 39:946–63
29. Bisarya R, Shaath D, Pirzad A, Satterwhite L, He J, Simpson SQ: Serum laktat buruk memprediksi saturasi oksigen vena sentral pada pasien sakit kritis: Sebuah studi kohort retrospektif. *J Perawatan Intensif.* 2019; 7:47
30. Pino, Richard M., Singh, Jaskaran. Appropriate Clinical Use of Lactate Measurements. *Anesthesiology* April 2021, Vol. 134, 637–644. <https://doi.org/10.1097/ALN.0000000000003655>

31. Uchida K, Nishimura T, Hagawa N, et al. The impact of early administration of vasopressor agents for the resuscitation of severe hemorrhagic shock following blunt trauma. *BMC Emerg Med.* 2020;20(1):26. Published 2020 Apr 16. doi:10.1186/s12873-020-00322-1
32. Kattan E, Hernández G. The role of peripheral perfusion markers and lactate in septic shock resuscitation. *J Intensive Med [Internet].* 2022;2(1):17–21. Available from: <https://www.sciencedirect.com/science/article/pii/S2667100X21000463>
33. Wise ES, Hocking KM, Polcz ME, Beilman GJ, Brophy CM, Sobey JH, Leisy PJ, Kiberenge RK, Alvis BD. Hemodynamic Parameters in the Assessment of Fluid Status in a Porcine Hemorrhage and Resuscitation Model. *Anesthesiology.* 2021 Apr 1;134(4):607-616. doi: 10.1097/ALN.0000000000003724. PMID: 33635950; PMCID: PMC7946734.
34. Castro R, Kattan E, Ferri G, Pairumani R, Valenzuela ED, Alegría L, et al. Effects of capillary refill time-vs. lactate-targeted fluid resuscitation on regional, microcirculatory and hypoxia-related perfusion parameters in septic shock: a randomized controlled trial. *Ann Intensive Care [Internet].* 2020;10(1):150. Available from: <https://doi.org/10.1186/s13613-020-00767-4>
35. Hernández G, et al. Effect of a Resuscitation Strategy Targeting Peripheral Perfusion Status vs Serum Lactate Levels on 28-Day Mortality Among Patients With Septic Shock: The ANDROMEDA-SHOCK Randomized Clinical Trial. *JAMA.* 2019 Feb 19;321(7):654-664. doi: 10.1001/jama.2019.0071. PMID: 30772908; PMCID: PMC6439620.
36. Jacquet-Lagrèze M, Bouhamri N, Portran P, Schweizer R, Baudin F, Lilot M, Fornier W, Fellahi JL. Capillary refill time variation induced by passive leg raising predicts capillary refill time response to volume expansion. *Crit Care.* 2019 Aug 16;23(1):281. doi: 10.1186/s13054-019-2560-0. PMID: 31420052; PMCID: PMC6697974.
37. Beloncle F, Meziani F, Lerolle N, Radermacher P, Asfar P. Does vasopressor therapy have an indication in hemorrhagic shock? *Ann Intensive Care.* 2013

May 22;3(1):13. doi: 10.1186/2110-5820-3-13. PMID: 23697682; PMCID: PMC3691630.

38. Hernandez G, Luengo C, Bruhn A, Kattan E, Friedman G, Ospina-Tascon GA, et al. When to stop septic shock resuscitation: clues from a dynamic perfusion monitoring. *Ann Intensive Care* 2014;4:30. doi:[10.1186/s13613-014-0030-z](https://doi.org/10.1186/s13613-014-0030-z).
39. Puskarich MA, Trzeciak S, Shapiro NI, et al. Whole blood lactate kinetics in patients undergoing quantitative resuscitation for severe sepsis and septic shock. *Chest*. 2013;143(6):1548-1553. doi:10.1378/chest.12-0878