

DAFTAR PUSTAKA

1. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, Bellomo R, Bernard GR, Chiche JD, Cooper-Smith CM, Hotchkiss RS, et al: The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016;315:801–10.
2. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving sepsis campaign: International guidelines for management of sepsis and septic shock: 2016. *Crit Care Med.* 2017;45(3):486–552.
3. Levy MM, Evans LE, Rhodes A. The surviving sepsis campaign bundle: 2018 update. *Intensive Care Med.* 2018;44(6):925–8.
4. Malbrain ML, Marik PE, Witters, Cordemans C, Kirkpatrick AW, Roberts DJ, Regenmortel NV. Fluid overload, de-resuscitation and outcomes in critically ill or injured patients: a systematic review with suggestions for clinical practice. *AIT Via Medica.* 2014;46:361–80.
5. Cordemans C, De laet I, Regenmortel NV, Schoonheydt K, Dits H, Huber W dkk. Fluid management in critically ill patients: the role of extravascular lung water, abdominal hypertension, capillary leak, and fluid balance. *Annals Intens Care.* 2012;2(Suppl 1):S1.
6. Backer DD, Cortes DO, Donadello K, Vincent JL. Pathophysiology of microcirculatory dysfunction and the pathogenesis of septic shock. *virulence.* 2014;5(1):73–9.
7. Granado RC, Mehta RL. Fluid overload in the ICU: evaluation and management. *BMC Nephrol.* 2016;17:1–9.
8. Strunden MS, Heckel K, Goetz AE, Reuter DA. Perioperative fluid and volume management: physiological basis, tools, and strategies. *Annals Og Intens Care.* 2011;1:1–8.
9. Stein A, Viera de Souza L, Belettina CB, Menegazzo WR, Viegas JR, Pereira EMC, dkk. Fluid overload and changes in serum creatinine after cardiac surgery: predictors of mortality and longer intensive care stay. A prospective cohort study. *Crit Care* 2012;16:1–9.

10. Rivers E, Nguyen B, Havstad S, Ressler J, Muzzin A, Knoblich B, dkk. Early goal directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med.* 2001;345:1368–77.
11. Cordemans C, De laet I, Regenmortel NV, Schoonheydt K, Dits H, Huber W, dkk. Fluid Management in critically ill patients: the role of extravascular lung water, abdominal hypertension, capillary leak, and fluid balance. *Annals Intens Care.* 2012;2(Suppl 1):1–12.
12. Hilton AK, Pellegrino VA, Scheinkestel CD. Avoiding common problems associated with intravenous fluid therapy. *MJA.* 2008;189(9):509–13.
13. Bouchard J, Soroko SB, Chertow GM, Himmelfarb J, Ikizler A, Paganini EP, dkk. Fluid accumulation, survival and recovery of kidney function in critically ill patients with acute kidney injury. *Kidney International.* 2009;76:422–7.
14. Gillespie RS, Seidel K, Symons JM. Effect of fluid overload and dose of replacement fluid on survival in hemofiltration. *Pediatr Nephrol.* 2004;19(12):1394–9.
15. Joen JS, Ji SM. Diagnostic value of procalcitonin and CRP in critically ill patients admitted with suspected sepsis. *J Dent Anesth Pain Med* 2015;15(3):135-140.
16. Ryu JA, Yang JH, Lee D et al. Clinical Usefulness of procalcitonin and C-Reactive Protein as outcome predictors in critically ill patients with severe sepsis and septic shock. *PLOS ONE journal* 2015; 1-12.
17. Xing-hai Chen, Yong-jie Yin, Jing-xiao Zhang. Sepsis and immune response. *World J Emerg Med.* 2011; 2:88–92.
18. Rizk NA, Kanafani ZA, Tabaja HZ, Kanj SS. Extended infusion of beta-lactam antibiotics: optimizing therapy in critically-ill patients in the era of antimicrobial resistance. *Expert Rev Anti Infect Ther.* Taylor & Francis; 2017;15(7):645–52. Available from: <https://doi.org/10.1080/14787210.2017.1348894>
19. Lien H, Pharm D. Extended Infusions of Beta-lactams in Pediatric Patients : The long and the short of it. 2017;23:1-23

20. Jang SM, Gharibian KN, Lewis SJ, Fissell WH, Tolwani AJ, Mueller BA. A Monte Carlo Simulation Approach for Beta-Lactam Dosing in Critically Ill Patients Receiving Prolonged Intermittent Renal Replacement Therapy. *J Clin Pharmacol*. 2018;58(10):1254–65.
21. Roberts JA, Kirkpatrick CMJ, Roberts MS, Robertson TA, Dalley AJ, Lipman J. Meropenem dosing in critically ill patients with sepsis and without renal dysfunction: Intermittent bolus versus continuous administration? Monte Carlo dosing simulations and subcutaneous tissue distribution. *J Antimicrob Chemother*. 2009;64(1):142–50.
22. Grupper M, Kuti JL, Nicolau DP. Continuous and prolonged intravenous β -lactam dosing: Implications for the clinical laboratory. *Clin Microbiol Rev*. 2016;29(4):750–72.
23. Hotchkiss R, Karl I. The pathophysiology and treatment of sepsis. *N Eng J Med*. 2003; 348:138-50.
24. Annane D, Bellissant E, Cavaillon JM. Septic shock. *Lancet* 2005; 365:63-78.
25. Acharya SP, Pradhan B, Marhatt MN. Application of the sequential organ failure assessment (SOFA) score in predicting outcome in ICU patients with SIRS. *Kathmandu University Medical Journal* 2007; 5(4): 475-83.