

## DAFTAR PUSTAKA

- [1] P. Bel *et al.*, “Clinical Nutrition ESPEN Prevalence and severity of malnutrition in hospitalized COVID-19 patients Doroth e,” no. xxxx, 2020, doi: 10.1016/j.clnesp.2020.09.018.
- [2] A. Academy, U. States, U. States, P. Services, and T. Force, “Important Information About Clinical Practice Guidelines : Key Tools for Improving Quality of Care and Patient Outcomes,” pp. 1–2, 2015, doi: 10.1111/wvn.12079.
- [3] P. Bel *et al.*, “Clinical Nutrition ESPEN Prevalence and severity of malnutrition in hospitalized COVID-19 patients Doroth e,” no. xxxx, 2020, doi: 10.1016/j.clnesp.2020.09.018.
- [4] D. K. Heyland, R. Dhaliwal, X. Jiang, and A. G. Day, “Identifying critically ill patients who benefit the most from nutrition therapy : the development and initial validation of a novel risk assessment tool,” 2011.
- [5] A. Rahman, R. M. Hasan, R. Agarwala, C. Martin, A. G. Day, and D. K. Heyland, “Identifying critically-ill patients who will benefit most from nutritional therapy : Further validation of the ‘ modified NUTRIC ’ nutritional risk assessment tool,” *Clinical Nutrition*, vol. 35, no. 1, pp. 158–162, 2016, doi: 10.1016/j.clnu.2015.01.015.
- [6] P. E. Wischmeyer and D. K. Heyland, “The Future of Critical Care Nutrition Therapy,” vol. 26, pp. 433–441, 2010, doi: 10.1016/j.ccc.2010.04.011.
- [7] L. A. Barker, B. S. Gout, and T. C. Crowe, “Hospital Malnutrition : Prevalence , Identification and Impact on Patients and the Healthcare System,” pp. 514–527, 2011, doi: 10.3390/ijerph8020514.
- [8] A. R. Blaser *et al.*, “Early enteral nutrition in critically ill patients : ESICM clinical practice guidelines,” *Intensive Care Med*, vol. 43, no. 3, pp. 380–398, 2017, doi: 10.1007/s00134-016-4665-0.
- [9] M. D. Rubinsky and A. P. Clark, “Early Enteral Nutrition in Critically Ill Patients,” no. October, 2012, doi: 10.1097/DCC.0b013e3182619944.
- [10] L. C. Robertson and M. Al-haddad, “Recognizing the critically ill patient,” *Anaesthesia and Intensive Care Medicine*, vol. 14, no. 1, pp. 11–14, 2013, doi: 10.1016/j.mpaic.2012.11.010.

- [11] R. S. Hotchkiss *et al.*, "HHS Public Access," pp. 1–47, 2017, doi: 10.1038/nrdp.2016.45.Sepsis.
- [12] R. S. Hotchkiss and I. E. Karl, "The Pathophysiology and Treatment of Sepsis," pp. 138–150, 2003.
- [13] J. C. Preiser, C. Ichai, J. C. Orban, and A. B. J. Groeneveld, "Metabolic response to the stress of critical illness," *British Journal of Anaesthesia*, vol. 113, no. 6. Elsevier Ltd, pp. 945–954, Dec. 01, 2014. doi: 10.1093/bja/aeu187.
- [14] K. Riskesdas, "Hasil Utama Riset Kesehata Dasar (RISKESDAS)," *J Phys A Math Theor*, vol. 44, no. 8, pp. 1–200, 2018, doi: 10.1088/1751-8113/44/8/085201.
- [15] "Competencies for Recognising and Responding to Acutely III Patients in Hospital".
- [16] M. P. Casaer and T. R. Ziegler, "Endocrine and metabolic considerations in critically ill patients 2 Nutritional support in critical illness and recovery," *Lancet Diabetes Endocrinol*, vol. 8587, no. 15, pp. 1–12, 2015, doi: 10.1016/S2213-8587(15)00222-3.
- [17] M. P. Casaer and T. R. Ziegler, "Endocrine and metabolic considerations in critically ill patients 2 Nutritional support in critical illness and recovery," *Lancet Diabetes Endocrinol*, vol. 8587, no. 15, pp. 1–12, 2015, doi: 10.1016/S2213-8587(15)00222-3.
- [18] P. Singer *et al.*, "ESPEN guideline on clinical nutrition in the intensive care unit," *Clinical Nutrition*, vol. 38, no. 1, pp. 48–79, 2019, doi: 10.1016/j.clnu.2018.08.037.
- [19] M. S. Siobal, J. E. Baltz, and J. Richardson, "A Guide to the Nutritional Assessment and Treatment of the Critically Ill Patient," 2021.
- [20] T. T. Maskoen, P. Dokter, and I. Care, *Panduan Nutrisi Klinis pada Pasien Sakit Kritis*. 2021.
- [21] L. A. Barker, B. S. Gout, and T. C. Crowe, "Hospital Malnutrition : Prevalence , Identification and Impact on Patients and the Healthcare System," pp. 514–527, 2011, doi: 10.3390/ijerph8020514.
- [22] D. K. Heyland, R. Dhaliwal, X. Jiang, and A. G. Day, "Identifying critically ill patients who benefit the most from nutrition therapy : the development and initial validation of a novel risk assessment tool," 2011.
- [23] N. M. Mehta *et al.*, *Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Pediatric Critically Ill Patient : Society of*

*Critical Care Medicine and American Society for Parenteral and Enteral Nutrition.* 2017. doi: 10.1177/0148607117711387.

- [24] M. S. Siobal, J. E. Baltz, and J. Richardson, “A Guide to the Nutritional Assessment and Treatment of the Critically Ill Patient,” 2021.
- [25] E. Reber, F. Gomes, M. F. Vasiloglou, and P. Schuetz, “Nutritional Risk Screening and Assessment,” pp. 1–19.
- [26] C. Machado, P. A. Defina, Y. Machado, E. Chinchilla, and E. Cuspineda, “It Is Imperative the use of Positive Air Pressure (CPAP) in Covid-19 Patients as Soon as the First Respiratory Symptoms Start,” *Frontiers Journal of Neurology and Neuroscience*, vol. 1, no. 1, pp. 1–14, 2020.
- [27] M. K. Alramly, M. S. Abdalrahim, and A. Khalil, “Validation of the modified NUTRIC score on critically ill Jordanian patients: A retrospective study,” *Nutr Health*, vol. 26, no. 3, pp. 225–229, 2020, doi: 10.1177/0260106020923832.
- [28] J. Vincent and J. Creteur, *Principles of Critical Care The Critically Ill Patient*, Third Edit. doi: 10.1016/B978-0-323-44942-7.00001-7.
- [29] J. Gil Cebrian, M. P. Bello Câmara, and R. Diaz-Alersi, “Apache II,” *Intensive Care Medicine*, vol. 13, no. 2. p. 143, 1987. doi: 10.1097/00003465-198603000-00013.
- [30] J. Marchetti, A. M. Dos Reis, A. F. Dos Santos, O. S. Franzosi, V. C. Luft, and T. Steemburgo, “O elevado risco nutricional está associado a desfechos desfavoráveis em pacientes internados na unidade de terapia intensiva,” *Rev Bras Ter Intensiva*, vol. 31, no. 3, pp. 326–332, 2019, doi: 10.5935/0103-507X.20190041.
- [31] B. Kucuk, S. Baltaci Ozen, G. M. Kocabeyoglu, N. M. Mutlu, E. Cakir, and I. Ozkocak Turan, “NUTRIC Score Is Not Superior to mNUTRIC Score in Prediction of Mortality of COVID-19 Patients,” *Int J Clin Pract*, vol. 2022, no. Cdc, pp. 1–9, 2022, doi: 10.1155/2022/1864776.
- [32] J. C.G., V. S., F. V., and H. S.G., “Mindfulness training affects attention--or is it attentional effort?,” *J Exp Psychol Gen*, vol. 141, no. 1, pp. 106–123, 2012.
- [33] A. Hill, G. Elke, and A. Weimann, “Nutrition in the intensive care unit—a narrative review,” *Nutrients*, vol. 13, no. 8, pp. 1–26, 2021, doi: 10.3390/nu13082851.
- [34] S. A. McClave *et al.*, “Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and

Enteral Nutrition (A.S.P.E.N.),” *Journal of Parenteral and Enteral Nutrition*, vol. 40, no. 2, pp. 159–211, 2016, doi: 10.1177/0148607115621863.

- [35] Y. M. Arabi *et al.*, “The intensive care medicine research agenda in nutrition and metabolism,” *Intensive Care Med*, vol. 43, no. 9, pp. 1239–1256, 2017, doi: 10.1007/s00134-017-4711-6.
- [36] Y. M. Arabi *et al.*, “The intensive care medicine research agenda in nutrition and metabolism,” *Intensive Care Med*, vol. 43, no. 9, pp. 1239–1256, 2017, doi: 10.1007/s00134-017-4711-6.
- [37] E. Estimation, “Journal of Parenteral and Enteral Nutrition,” 2013, doi: 10.1177/0148607113505868.
- [38] O. Zusman, M. Theilla, J. Cohen, I. Kagan, I. Bendavid, and P. Singer, “Resting energy expenditure , calorie and protein consumption in critically ill patients : a retrospective cohort study,” *Crit Care*, pp. 1–8, 2016, doi: 10.1186/s13054-016-1538-4.
- [39] P. J. M. Weijs, W. G. P. M. Looijaard, A. Beishuizen, A. R. J. Girbes, and H. M. O. Straaten, “Early high protein intake is associated with low mortality and energy overfeeding with high mortality in non-septic mechanically ventilated critically ill patients,” pp. 1–10, 2014, doi: 10.1186/s13054-014-0701-z.
- [40] D. C. Frankenfield and C. M. Ashcraft, “Estimating energy needs in nutrition support patients,” *Journal of Parenteral and Enteral Nutrition*, vol. 35, no. 5, pp. 563–570, 2011, doi: 10.1177/0148607111415859.
- [41] D. C. Frankenfield, A. Coleman, S. Alam, and R. N. Cooney, “Analysis of estimation methods for resting metabolic rate in critically ill adults,” *Journal of Parenteral and Enteral Nutrition*, vol. 33, no. 1, pp. 27–36, 2009, doi: 10.1177/0148607108322399.
- [42] E. Estimation, “Journal of Parenteral and Enteral Nutrition,” 2013, doi: 10.1177/0148607113505868.
- [43] T. Oshima *et al.*, “Indirect calorimetry in nutritional therapy. A position paper by the ICALIC study group,” *Clinical Nutrition*, vol. 36, no. 3, pp. 651–662, 2017, doi: 10.1016/j.clnu.2016.06.010.
- [44] J. F. Savard, C. Faisy, N. Lerolle, E. Guerot, J. L. Diehl, and J. Y. Fagon, “Validation of a predictive method for an accurate assessment of resting energy expenditure in medical mechanically ventilated patients,” *Crit Care Med*, vol. 36, no. 4, pp. 1175–1183, 2008, doi: 10.1097/CCM.0b013e3181691502.

- [45] M. J. Rennie, "Anabolic resistance in critically ill patients," *Crit Care Med*, vol. 37, no. SUPPL. 10, pp. 398–399, 2009, doi: 10.1097/CCM.0b013e3181b6ec1f.
- [46] A. MacDonald and L. Hildebrandt, "Comparison of formulaic equations to determine energy expenditure in the critically ill patient," *Nutrition*, vol. 19, no. 3, pp. 233–239, 2003, doi: 10.1016/S0899-9007(02)01033-X.
- [47] J. A. Wooley and H. C. Sax, "Indirect Calorimetry: Applications to Practice," *Nutrition in Clinical Practice*, vol. 18, no. 5, pp. 434–439, 2003, doi: 10.1177/0115426503018005434.
- [48] S. A. McClave, R. G. Martindale, and L. Kiraly, "The use of indirect calorimetry in the intensive care unit," *Curr Opin Clin Nutr Metab Care*, vol. 16, no. 2, pp. 202–208, 2013, doi: 10.1097/MCO.0b013e32835dbc54.
- [49] F. B. Cerra *et al.*, "Applied nutrition in ICU patients: A consensus statement of the American College of Chest Physicians," *Chest*, vol. 111, no. 3, pp. 769–778, 1997, doi: 10.1378/chest.111.3.769.
- [50] A. Raymond, H. Van Zanten, E. De Waele, and P. E. Wischmeyer, "Nutrition therapy and critical illness : practical guidance for the ICU , post-ICU , and long-term convalescence phases," pp. 1–10, 2019.
- [51] A. M. Deane *et al.*, "The Effects of Critical Illness on Intestinal Glucose Sensing, Transporters, and Absorption\*," vol. 2, pp. 57–65, doi: 10.1097/CCM.0b013e318298a8af.
- [52] A. Thorell *et al.*, "Intensive Insulin Treatment in Critically Ill Trauma Patients Normalizes Glucose by Reducing Endogenous Glucose Production," vol. 89, no. October, pp. 5382–5386, 2015, doi: 10.1210/jc.2004-1118.
- [53] "Effects of isoenergetic glucose-based or lipid-based parenteral nutrition on glucose metabolism, de novo lipogenesis, and respiratory gas exchanges in critically ill patients - PubMed."
- [54] Y. Han *et al.*, "The clinical and economic impact of the use of diabetes-specific enteral formula on ICU patients with type 2 diabetes," *Clinical Nutrition*, no. October, pp. 1–6, 2016, doi: 10.1016/j.clnu.2016.09.027.
- [55] A. R. H. Van Zanten, "Should We Increase Protein Delivery During Critical Illness ? Critically Ill Protein Catabolism in Critical Illness What Is the Best Protein Target in Critical Illness ?," 2016, doi: 10.1177/0148607115626905.
- [56] I. Bendavid, O. Zusman, I. Kagan, M. Theilla, and J. Cohen, "Early Administration of Protein in Critically Ill Patients : A Retrospective Cohort Study," pp. 1–10, doi: 10.3390/nu11010106.

- [57] P. J. M. Weijs *et al.*, “Low skeletal muscle area is a risk factor for mortality in mechanically ventilated critically ill patients,” pp. 1–7, 2014.
- [58] Z. A. Puthucheary *et al.*, “Acute Skeletal Muscle Wasting in Critical Illness,” vol. 310, no. 15, pp. 1591–1600, 2013, doi: 10.1001/jama.2013.278481.
- [59] P. Singer *et al.*, “ESPEN guideline on clinical nutrition in the intensive care unit,” *Clinical Nutrition*, vol. 38, no. 1, pp. 48–79, Feb. 2019, doi: 10.1016/j.clnu.2018.08.037.
- [60] J. Werner *et al.*, “Metabolic support in the critically ill: A consensus of 19,” *Critical Care*, vol. 23, no. 1. BioMed Central Ltd., Sep. 18, 2019. doi: 10.1186/s13054-019-2597-0.
- [61] G. Elke *et al.*, “Clinical Nutrition in Critical Care Medicine – Guideline of the German Society for Nutritional Medicine (DGEM),” *Clin Nutr ESPEN*, vol. 33, pp. 220–275, Oct. 2019, doi: 10.1016/j.clnesp.2019.05.002.
- [62] “Effect of Critical Illness on Triglyceride Absorption — Monash University.”
- [63] P. Green, M. Theilla, and P. Singer, “Lipid metabolism in critical illness,” 2016, doi: 10.1097/MCO.0000000000000253.
- [64] D. Chassard, B. Allaouchiche, J. C. Dumont, and C. Auboyer, “Glucose-lipid ratio is a determinant of nitrogen balance during total parenteral nutrition in critically ill patients: a prospective , randomized , multicenter blind trial with an intention-to-treat analysis,” pp. 1394–1400, 2005, doi: 10.1007/s00134-005-2771-5.
- [65] J. A. Barker, “Long-term Mortality Outcome Associated With Prolonged Admission to the ICU,” *Yearbook of Pulmonary Disease*, vol. 2007, pp. 338–339, 2007, doi: 10.1016/s8756-3452(08)70509-7.
- [66] D. A. Dongelmans *et al.*, “Linking of global intensive care (LOGIC): An international benchmarking in critical care initiative,” *J Crit Care*, vol. 60, pp. 305–310, 2020, doi: 10.1016/j.jcrc.2020.08.031.
- [67] C. Alberda *et al.*, “The relationship between nutritional intake and clinical outcomes in critically ill patients: Results of an international multicenter observational study,” *Intensive Care Med*, vol. 35, no. 10, pp. 1728–1737, 2009, doi: 10.1007/s00134-009-1567-4.
- [68] C. F. L. Couto, Â. Dariano, C. Texeira, C. H. Da Silva, A. B. Torbes, and G. Friedman, “Adequacy of enteral nutritional support in intensive care units does not affect the short- And long-term prognosis of mechanically ventilated patients: A pilot study,” *Rev Bras Ter Intensiva*, vol. 31, no. 1, pp. 34–38, 2019, doi: 10.5935/0103-507X.20190004.

- [69] Z. Y. Lee, M. Y. B. Nisak, and I. N. Airini, "Feeding adequacy among Critically Ill Patients in the Intensive Care Unit and Its Association with Clinical Outcomes: A Narrative Review," *Bangladesh Critical Care Journal*, vol. 5, no. 2, pp. 113–121, 2017, doi: 10.3329/bccj.v5i2.34390.
- [70] D. A. Gruenberg, W. Shelton, S. L. Rose, A. E. Rutter, S. Socaris, and G. McGee, "Factors influencing length of stay in the intensive care unit," *American Journal of Critical Care*, vol. 15, no. 5. pp. 502–509, Sep. 2006. doi: 10.4037/ajcc2006.15.5.502.
- [71] Rotter T, Kinsman L, J. El, and Willis J, "Clinical pathways: effects on professional practice, patient outcomes, length of stay and hospital costs (Review)," 2010. [Online]. Available: <http://www.thecochranelibrary.com>
- [72] K. B. Laupland, A. W. Kirkpatrick, J. B. Kortbeek, and D. J. Zuege, "Long-term mortality outcome associated with prolonged admission to the ICU," *Chest*, vol. 129, no. 4, pp. 954–959, 2006, doi: 10.1378/chest.129.4.954.
- [73] A. W. Ong *et al.*, "Characteristics and outcomes of trauma patients with ICU lengths of stay 30 days and greater: A seven-year retrospective study," *Crit Care*, vol. 13, no. 5, Sep. 2009, doi: 10.1186/cc8054.
- [74] A. Ingeman, G. Andersen, H. H. Hundborg, M. L. Svendsen, and S. P. Johnsen, "In-Hospital Medical Complications, Length of Stay, and Mortality Among Stroke Unit Patients," 2011, doi: 10.1161/STROKES.111.610881/-DC1.
- [75] J. A. Luker, J. Bernhardt, and K. A. Grimmer-Somers, "Demographic and stroke-related factors as predictors of quality of acute stroke care provided by allied health professionals," *J Multidiscip Healthc*, vol. 4, pp. 247–259, Jul. 2011, doi: 10.2147/JMDH.S22569.
- [76] D. K. Heyland, E. Konopad, T. W. Noseworthy, R. Johnston, and A. Gafni, "Is it 'worthwhile' to continue treating patients with a prolonged stay (>14 days) in the ICU? An economic evaluation," *Chest*, vol. 114, no. 1, pp. 192–198, 1998, doi: 10.1378/chest.114.1.192.
- [77] D. K. Heyland, R. Dhaliwal, X. Jiang, and A. G. Day, "Identifying critically ill patients who benefit the most from nutrition therapy: the development and initial validation of a novel risk assessment tool," 2011. [Online]. Available: <http://ccforum.com/content/15/6/R268>
- [78] F. G. Yanowsky-Escatell *et al.*, "Use of mNUTRIC-Score for Nutrition Risk Assessment and Prognosis Prediction in Critically Ill Patients with COVID-19: A Retrospective Observational Study," *Crit Care Res Pract*, vol. 2021, 2021, doi: 10.1155/2021/5866468.

- [79] M. C. De Vries, W. Koekkoek, M. H. Opdam, D. Van Blokland, and A. R. Van Zanten, “Nutritional assessment of critically ill patients: Validation of the modified NUTRIC score,” in *European Journal of Clinical Nutrition*, Nature Publishing Group, Mar. 2018, pp. 428–435. doi: 10.1038/s41430-017-0008-7.
- [80] A. Rahman, R. M. Hasan, R. Agarwala, C. Martin, A. G. Day, and D. K. Heyland, “Identifying critically-ill patients who will benefit most from nutritional therapy: Further validation of the ‘modified NUTRIC’ nutritional risk assessment tool,” *Clinical Nutrition*, vol. 35, no. 1, pp. 158–162, Feb. 2016, doi: 10.1016/j.clnu.2015.01.015.
- [81] D. H. Jeong *et al.*, “Comparison of accuracy of NUTRIC and modified NUTRIC scores in predicting 28-day mortality in patients with sepsis: A single center retrospective study,” *Nutrients*, vol. 10, no. 7, Jul. 2018, doi: 10.3390/nu10070911.
- [82] C. Y. Wang, P. K. Fu, W. C. Chao, W. N. Wang, C. H. Chen, and Y. C. Huang, “Full versus trophic feeds in critically ill adults with high and low nutritional risk scores: A randomized controlled trial,” *Nutrients*, vol. 12, no. 11, pp. 1–12, Nov. 2020, doi: 10.3390/nu12113518.
- [83] Y. Finkel, R. D. Baker, P. Rosenthal, and P. M. Sherman, “News and Views A Critical Perspective on Trophic Feeding,” 2004. [Online]. Available: <http://journals.lww.com/jpgn>
- [84] S. J. Peterson, L. McKeever, O. B. Lateef, S. Freels, G. Fantuzzi, and C. A. Braunschweig, “Combination of High-Calorie Delivery and Organ Failure Increases Mortality Among Patients With Acute Respiratory Distress Syndrome,” *Crit Care Med*, vol. 47, no. 1, pp. 69–75, Jan. 2019, doi: 10.1097/CCM.0000000000003476.
- [85] S. J. Peterson, O. B. Lateef, S. Freels, L. McKeever, G. Fantuzzi, and C. A. Braunschweig, “Early Exposure to Recommended Calorie Delivery in the Intensive Care Unit Is Associated With Increased Mortality in Patients With Acute Respiratory Distress Syndrome,” *Journal of Parenteral and Enteral Nutrition*, vol. 42, no. 4, pp. 739–747, May 2018, doi: 10.1177/0148607117713483.
- [86] T. W. Rice, S. Mogan, M. A. Hays, G. R. Bernard, G. L. Jensen, and A. P. Wheeler, “Randomized trial of initial trophic versus full-energy enteral nutrition in mechanically ventilated patients with acute respiratory failure,” in *Critical Care Medicine*, Lippincott Williams and Wilkins, 2011, pp. 967–974. doi: 10.1097/CCM.0b013e31820a905a.

- [87] N. M. Mehta *et al.*, "Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Pediatric Critically Ill Patient: Society of Critical Care Medicine and American Society for Parenteral and Enteral Nutrition," *Pediatric Critical Care Medicine*, vol. 18, no. 7, pp. 675–715, Jul. 2017, doi: 10.1097/PCC.0000000000001134.
- [88] D. Zhang, H. Li, Y. Li, and L. Qu, "Gut rest strategy and trophic feeding in the acute phase of critical illness with acute gastrointestinal injury," *Nutr Res Rev*, vol. 32, no. 2, pp. 176–182, Dec. 2019, doi: 10.1017/S0954422419000027.
- [89] Y. Arabi, S. Venkatesh, S. Haddad, A. Al Shimemeri, and S. Al Malik, "A prospective study of prolonged stay in the intensive care unit: predictors and impact on resource utilization," 2002.