

DAFTAR PUSTAKA

1. Patel SB, Kress JP. Sedation and analgesia in the mechanically ventilated patient. *Am J Respir Crit Care Med.* 2012;185(5):486–97.
2. Panahi Y, Dehcheshmeh HS, Mojtahedzadeh M, Joneidi-Jafari N, Johnston TP, Sahebkar A. Analgesic and sedative agents used in the intensive care unit: A review. *Journal of Cellular Biochemistry.* 2018;119(11), 8684–8693.
3. Ghodraty M, Pournajafian A, Niakan M, Totonchi GM, Mazhari F. Comparing the Effects of Sedation Induced By Fentanyl and Remifentanil in Patients Under Mechanical Ventilation. *Arak Med Univ J.* 2016;18(10103):73–9
4. Devlin JW, Skrobik Y, Gélinas C, Needham DM, Slooter AJC, Pandharipande PP, et al. Clinical practice guidelines for the prevention and management of pain, agitation/sedation, delirium, immobility, and sleep disruption in adult patients in the ICU. *Crit Care Med.* 2018;46:e825–73.
5. Narayanan M, Venkataraju A, Jennings J. Analgesia in intensive care: part 1. *BJA Education.* 2016, 16(2), 72–78.
6. Prajapat B, Sandhya AS, Chaudhry D, Esquinas AM. Opioids in intensive care unit (ICU): Causal agent for sleep apnea, unmasking the undiagnosed sleep apnea or both in the ICU. Oxford University Press. 2018, 12-19
7. Zhu Y, Wang Y, Du B, Xi X. Could remifentanil reduce duration of mechanical ventilation in comparison with other opioids for mechanically ventilated patients? A systematic review and meta-analysis. *Crit Care.* 2017;21(1).
8. Hyun DG, Huh JW, Hong SB, Koh Y, Lim CM. Iatrogenic opioid withdrawal syndrome in critically ill patients: A retrospective cohort study. *J Korean Med Sci.* 2020;35(15):1–12.
9. Öztürk D, Turan S, Ayik I, Karadeniz U, Soyal O, Coşar A, Özgök A, Erdemli Ö. Effects of remifentanil and fentanyl on postoperative pain and recovery in fast-track cardiac surgery. *GKDA Derg.* 2012, 18(4), 80–87.

10. Fountas A, Chai ST, Kourkouti C, Karavitaki N . Mechanisms of endocrinology: Endocrinology of opioids. In *European Journal of Endocrinology* BioScientifica Ltd. 2018,179(4), 183-196.
11. Azemati, S., Savaie, M., Khosravi, M. B., & Allahyari, E. (2013). *Combination of remifentanil with isoflurane or propofol: Effect on the surgical stress response* . *Acta Anaesthesiol Belg*. 2013;64(1):25-31
12. Tafur-Betancourt LA, Arévalo-Sánchez M, Lema-Flórez E. Effect on cough frequency and intensity during extubation of two plasma concentrations of remifentanil using TACAN: Randomised controlled clinical trial. *Colomb J Anesthesiol*. 2017;45(2):92–9.
13. Huttmann SE, Magnet FS, Karagiannidis C, Storre JH, Windisch, W. Quality of life and life satisfaction are severely impaired in patients with long-term invasive ventilation following ICU treatment and unsuccessful weaning. *Annals of Intensive Care*, 2018. 8(1).
14. Devabhakthuni S, Armahizer MJ, Dasta JF, Kane-Gill SL. Analgosedation : A Paradigm Shift in Intensive Care Unit Sedation Practice. *Annals of Pharmacotherapy* ,2012;46(4): 530–540.
15. Cuesta JM, Singer M. The stress response and critical illness: A review. *Critical Care Medicine*, 2012. 40(12), 3283–3289.
16. Al-sutari MM, Abdalrahim,MS, Hamdan-MansourAM, Ayasrah SM. Pain among mechanically ventilated patients in critical care units. In *Journal of Research in Medical Sciences*. J Res Med Sci. 2014;19(8):726-732.
17. Kotfis K, Zegan-Baraska M, Szydłowski L, Ukowski M, Ely EW. Methods of pain assessment in adult intensive care unit patients-Polish version of the CPOT (Critical Care Pain Observation Tool) and BPS (Behavioral Pain Scale). *Anaesthesiology Intensive Therapy* ; 49(1): 66–72.
18. Farooq F, Khan R, Ahmed A. Assessment of patient satisfaction with acute pain management service: Monitoring quality of care in clinical setting. *Indian J Anaesth*. 2016;60(4):248–52.

19. Havaladar AA. Pain Assessment in Intensive Care Unit: A Forgotten Entity or a Quality Indicator. *Indian Journal of Critical Care Medicine*. 2022; 26 (4): 419–420.
20. Severgnini P, Pelosi P, Contino E, Serafinelli E, Novario R, Chiaranda M. Accuracy of Critical Care Pain Observation Tool and Behavioral Pain Scale to assess pain in critically ill conscious and unconscious patients: Prospective, observational study. *Journal of Intensive Care*. 2016, 4(1).
21. Klompas M, Li L, Szumita P, et al. Associations Between Different Sedatives and Ventilator-Associated Events, Length of Stay, and Mortality in Patients Who Were Mechanically Ventilated. *Chest* 2016; 149:1373.
22. Muller JC, Kennard JW, Browne J S, Fecher AM, Hayward TZ . Hemodynamic monitoring in the intensive care unit. *Nutrition in Clinical Practice* 2012; 27 (3): 340–351.
23. Howell SJ. Consensus statements and expert guidance: interpret with care. *British Journal of Anaesthesia* 2019;22(6): 719–722).
24. Kwon YS, Jang JS, Hwang SM, Tark H, Kim JH, Lee JJ, Li Y. Effects of surgery start time on postoperative cortisol, inflammatory cytokines, and postoperative hospital day in hip surgery: Randomized controlled trial. *Medicine (United States)* 2019: 98(24).
25. Lovallo WR, Enoch MA, Acheson A, Cohoon AJ, Sorocco KH, Hodgkinson CA, Vincent AS, Glahn DC, Goldman D. (2015). Cortisol Stress Response in Men and Women Modulated Differentially by the Mu-Opioid Receptor Gene Polymorphism OPRM1 A118G. *Neuropsychopharmacology*, 2015; 40(11): 2546–2554.
26. Blum CA, Velly L, Brochet C, Ziegler F, Tavolacci MP, Hausfater P, Lvovschi VE. Relevance of cortisol and copeptin blood concentration changes in an experimental pain model. *Scientific Reports* 2022; 12(1).
27. Su WL, Chen YH, Chen CW, Yang SH, Su CL, Perng WC, Wu CP, Chen JH. Involuntary cough strength and extubation outcomes for patients in an ICU. *Chest*, 2010; 137(4): 777–782.

28. Dickinson RS, Morjaria JB, Wright CE, Morice AH. Is opiate action in cough due to sedation? *Therapeutic Advances in Chronic Disease* 2014; 5(5): 200–205.
29. Zetterlund P, Plos K, Bergbom I, Ringdal M. Memories from intensive care unit persist for several years-A longitudinal prospective multi-centre study. *Intensive and Critical Care Nursing* 2012; 28(3): 159–167.
30. Barr J, Fraser GL, Puntillo K, Ely EW, Gélinas C, Dasta JF, et al. Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. *Crit Care Med.* 2013;41:263–306
31. Devlin JW, Skrobik Y, Gélinas C, Needham DM, Slooter AJC, Pandharipande PP, et al. Clinical practice guidelines for the prevention and management of pain, agitation/sedation, delirium, immobility, and sleep disruption in adult patients in the ICU. *Crit Care Med.* 2018;46:e825–73.
32. Aoki Y, Kato H, Fujimura N, Suzuki Y, Sakuraya M, Doi M. Effects of fentanyl administration in mechanically ventilated patients in the intensive care unit: a systematic review and meta-analysis. *BMC Anesthesiology*, 2022; 22(1).
33. Tedders KM, McNorton KN, Edwin SB. Efficacy and safety of analgosedation with fentanyl compared with traditional sedation with propofol. *Pharmacotherapy* 2014; 34(6), 643–647.
34. Wilhelm W, Kreuer S. The place for short-acting opioids: special emphasis on remifentanil. *Crit Care.* 2008;12(3):S5.
35. Zhu Y, Wang Y, Du B, Xi X. Could remifentanil reduce duration of mechanical ventilation in comparison with other opioids for mechanically ventilated patients? A systematic review and meta-analysis. *Crit Care.* 2017;21:206
36. Pujol J, Martínez-Vilavella G, Gallart L, Blanco-Hinojo L, Pacreu S, Bonhomme V, et al. Effects of remifentanil on brain responses to noxious stimuli during deep propofol sedation. *British Journal of Anaesthesia* 2023 130(2), 330–338.
37. Yu EHY, Tran DHD, Lam SW, Irwin MG. Remifentanil tolerance and hyperalgesia: short-term gain, long-term pain? *Anaesthesia.* 2016;71:1347–62

38. Rozendaal FW, Spronk PE, Snellen FF, Schoen A, Zanten ARH, Foudraïne NA, Mulder PGH, Bakker J. Remifentanil-propofol analgo-sedation shortens duration of ventilation and length of ICU stay compared to a conventional regimen: A centre randomised, cross-over, open-label study in the Netherlands. *Intensive Care Medicine*, 2009 35(2), 291–298.
39. Dahaba AA, Grabner T, Rehak PH, List WF, Metzler H. Remifentanil versus morphine analgesia and sedation for mechanically ventilated critically ill patients: a randomized double blind study. *Anesthesiology* 2004;101(3):640-6.
40. Muellejans B, López A, Cross MH, Bonome C, Morrison L, Kirkham AJT. Remifentanil versus fentanyl for analgesia based sedation to provide patient comfort in the intensive care unit: a randomized, double-blind controlled trial *Critical Care (London, England)*, 2004.8(1).
41. Son HW, Lee JM, Park SH, Lee YJ, Oh JM, Hwang SK. Fentanyl versus Remifentanil for Cough Suppression and Recovery after Video-Assisted Thoracic Surgery. *J Chest Surg.* 2021;54(3):200–5.
42. Askin T, Karadeniz Ü, Ünver S, Boran E, Erdemli Ö. An efficacy comparison of fentanyl and remifentanil during off-pump coronary artery bypass graft surgery. *Turkish J Thorac Cardiovasc Surg.* 2013;21(3):683–90.
43. Ismaiel NM, Chankalal R, Zhou J, Henzler D. Using remifentanil in mechanically ventilated rats to provide continuous analgo-sedation. *J Am Assoc Lab Anim Sci.* 2012;51(1):58–62.
44. Poterman M, Kalmar AF, Buisman PL, Struys MMRF, Scheeren TWL. Improved haemodynamic stability and cerebral tissue oxygenation after induction of anaesthesia with sufentanil compared to remifentanil: A randomised controlled trial. *BMC Anesthesiol.* 2020;20(1):1–12.
45. Lee SK, Jeong MA, Sung JM, Yeon HJ, Chang JH, Lim H. Effect of remifentanil infusion on the hemodynamic response during induction of anesthesia in hypertensive and normotensive patients: a prospective observational study. *J Int Med Res.* 2019;47(12):6254–67.
46. Han JI. Should the clinical study on opioid-induced cough continue? *Korean J Anesthesiol.* 2011;60(6):391–2.

47. Lim JH, Ryu SJ, Lim YS. The incidence of cough induced by remifentanil during anesthetic induction was decreased by graded escalation of the remifentanil concentration. *Korean J Anesthesiol.* 2010;58(2):117–21.
48. Putro PWY, Nasihun T. Comparison in Efficacy between Fentanyl Continuous Infusion and Intrathecal Morphine for Pain After Caesarean Section Management. *Sains Med.* 2017;8(2):83–7.
49. Haroutounian S. Postoperative opioids, endocrine changes, and immunosuppression. *Schmerz.* 2018;32(5):374–80.
50. Winterhalter M, Brandl K, Rahe-Meyer N, Osthaus A, Hecker H, Hagl C, et al. Endocrine stress response and inflammatory activation during CABG surgery. A randomized trial comparing remifentanil infusion to intermittent fentanyl. *Eur J Anaesthesiol.* 2008;25(4):326–35.
51. Yuan Y, Liu L, Feng S, Ma Y, Wang Q. Effects of different remifentanil doses on the stress reaction and bis value of video laryngoscope-guided tracheal intubation. *Trop J Pharm Res.* 2018;17(10):2025–30.
52. Asakura A, Mihara T, Goto T. Does fentanyl or remifentanil provide better postoperative recovery after laparoscopic surgery? A randomized controlled trial. *BMC Anesthesiol.* 2018;18(1):1–8.
53. Çay M. The Effect of Cortisol Level Increasing Due to Stress in Healthy Young Individuals on Dynamic and Static Balance Scores. *North Clin Istanbul.* 2017;5(4):295–301.
54. Figueira JA, Sarafim-Silva BAM, Gonçalves GM, Aranha LN, Lopes FL, Corrente JE, et al. Predisposing factors for increased cortisol levels in oral cancer patients. *Compr Psychoneuroendocrinology.* 2022;9.
55. Haytural C, Aydnı B, Demir B, Bozkurt E, Parlak E, Dişibeyaz S, et al. Comparison of Propofol, Propofol-Remifentanil, and Propofol-Fentanyl Administrations with Each Other Used for the Sedation of Patients to Undergo ERCP. *Biomed Res Int.* 2015;2015.
56. Akcaboy ZN, Akcaboy EY, Albayrak D, Altinoren B, Dikmen B, Gogus N. Can remifentanil be a better choice than propofol for colonoscopy during monitored anesthesia care? *Acta Anaesthesiol Scand.* 2006;50(6):736–41.

57. Samantaray A, Rao MH. Effects of fentanyl on procedural pain and discomfort associated with central venous catheter insertion: A prospective, randomized, double-blind, placebo controlled trial. *Indian J Crit Care Med.* 2014;18(7):421–6.
58. Karol D, Weiniger CF. Update on Non-neuraxial Labor Analgesia. *Curr Anesthesiol Rep* [Internet]. 2021;11:348–54.