

5.1 Kesimpulan

Berdasarkan hasil dan pembahasan diatas, maka dapat diambil kesimpulan, yaitu: Tidak didapatkan adanya hubungan yang signifikan antara nilai NLR terhadap kesembuhan luka dengan korelasi negative dari NLR terhadap kesembuhan luka. Semakin tinggi nilai NLR maka semakin rendah prognosis kesembuhan suatu luka pada pasien trauma khususnya pasien dengan fraktur terbuka pada tibia begitu pula sebaliknya

5.2 Saran

Penelitian ini diharapkan dapat dijadikan sebagai batu loncatan dan referensi untuk penelitian-penelitian selanjutnya, khususnya mengenai hubungan antara indikator inflamasi lainnya terhadap kesembuhan luka pasien fraktur terbuka .

DAFTAR PUSTAKA

1. Maruyama Y, Inoue K, Mori K, Gorai K, Shimamoto R, Onitsuka T, et al. Neutrophil-lymphocyte ratio and platelet-lymphocyte ratio as predictors of wound healing failure in head and neck reconstruction. *Acta oto-laryngologica*. 2017;137(1):106–10.
2. Wilgus TA, Roy S, McDaniel JC. Neutrophils and wound repair: positive actions and negative reactions. *Advances in wound care*. 2013;2(7):379–88.

3. Muire PJ, Mangum LH, Wenke JC. Time Course of Immune Response and Immunomodulation During Normal and Delayed Healing of Musculoskeletal Wounds. *Front Immunol.* 2020; 11: 1056. Epub 2020/06/26. doi: 10.3389/fimmu.2020.01056. PubMed PMID: 32582170;
4. Raziyeva K, Kim Y, Zharkinbekov Z, Kassymbek K, Jimi S, Saparov A. Immunology of acute and chronic wound healing. *Biomolecules.* 2021;11(5):700.
5. Xia X, Li K, Wu R, Lv Q, Deng X, Fei Z, et al. Predictive value of neuron-specific enolase, neutrophil-to-lymphocyte-ratio and lymph node metastasis for distant metastasis in small cell lung cancer. *Clin Respir J.* 2020 Nov;14(11):1060–6.
6. Kovtun A, Messerer DA, Scharfffetter-Kochanek K, Huber-Lang M, Ignatius A. Neutrophils in tissue trauma of the skin, bone, and lung: two sides of the same coin. *Journal of immunology research.* 2018;2018.
7. Bastian OW, Kuijer A, Koenderman L, Stellato RK, van Solinge WW, Leenen LP, et al. Impaired bone healing in multitrauma patients is associated with altered leukocyte kinetics after major trauma. *Journal of Inflammation Research.* 2016;9:69.
8. Neumann MV, Strohm PC, Reising K, Zwingmann J, Hammer TO, Suedkamp NP. Complications after surgical management of distal lower leg fractures. *Scand J Trauma Resusc Emerg Med.* 2016 Dec 9;24(1):146.
9. Johnson BZ, Stevenson AW, Prèle CM, Fear MW, Wood FM. The role of IL-6 in skin fibrosis and cutaneous wound healing. *Biomedicines.* 2020;8(5):101.
10. Coates BA, McKenzie JA, Yoneda S, Silva MJ. Interleukin-6 (IL-6) deficiency enhances intramembranous osteogenesis following stress fracture in mice. *Bone.* 2021;143:115737.
11. Poletti FL, Macmull S, Mushtaq N, Mobasher R. Current concepts and principles in open tibial fractures—Part II Management and controversies. *MOJ Orthop Rheumatol.* 2017;8(2):00305.
12. Gustilo RB, Mendoza RM, Williams DN. Problems in the management of type III (severe) open fractures: a new classification of type III open fractures. *J Trauma.* 1984 Aug;24(8):742–6.
13. Muire, P. J., Mangum, L. H., & Wenke, J. C. (2020). Time course of immune response and immunomodulation during normal and delayed healing of musculoskeletal wounds. *Frontiers in Immunology*, 11. <https://doi.org/10.3389/fimmu.2020.01056>
14. Vatankhah, N., Jahangiri, Y., Landry, G. J., McLafferty, R. B., Alkayed, N. J., Moneta, G. L., & Azarbal, A. F. (2017). Predictive value of neutrophil-to-lymphocyte ratio in

diabetic wound healing. *Journal of Vascular Surgery*, 65(2), 478–483.
<https://doi.org/10.1016/j.jvs.2016.08.108>