

## DAFTAR PUSTAKA

- Affandi R, Sjafei DS, Rahardjo MF, & Sulistiono. Iktiologi. Suatu Pedoman Kerja Laboratorium. Departemen Pendidikan dan Kebudayaan. Direktorat Jenderal Pendidikan Tinggi. Pusat Antar Universitas Ilmu Hayat. Bogor: Institut Pertanian Bogor, 1992.
- Asiah, N., Sukendi, S., Junianto, J., Yustiati, A., & Windarti, W. (2019). Truss morfometrik dan karakter meristik ikan kelabau (*Osteochilus melanopleurus* Bleeker, 1852) dari tiga populasi di Sungai Kampar, Sungai Siak, dan Sungai Rokan, Provinsi Riau. *Jurnal Iktiologi Indonesia*, 19(2), 283–295. <http://jurnal-iktiologi.org/index.php/jii/article/view/462/pdf>.
- Cadrin, S. X., & Friedland, K. D. (1999). The utility of image processing techniques for morphometric analysis and stock identification. *Fisheries Research*, 43(1–3), 129–139. [https://doi.org/10.1016/S0165-7836\(99\)00070-3](https://doi.org/10.1016/S0165-7836(99)00070-3).
- Caivalcanti, M. J., Monteiro, L. R., & Lopes, P. R. D. (1999). Landmark-based morphometric analysis in selected species of serranid fishes (Perciformes: Teleostei). *Zoological Studies*, 38(3), 287–294.
- Chase, P.D. 2014. Meristics, pp. 171-184. In Cadrin, S.X., L.A. Kerr, & S. Mariani (eds.). 2014. Stock Identification Methods: Applications in Fishery Science. Second edition. Academic Press, London
- Davis TLO. (1982). Maturity and sexuality in barramundi, *Lates calcarifer* (Bloch). In the Northern Territory and south eastern Gulf of Carpentaria. Australia Jurnal of Marine and Fresh Research, 33:529-45.
- Effendie MI. 2002. *Biologi Perikanan*. Cetakan Kedua. Yogyakarta. Yayasan Pustaka Nusatama. 163 Halaman
- Elawa, A. 2004. Morphometric: Application in Biology and Paleontology. ISBN 3-540 21429-1 Springer-Verlag. Berlin. Heidelberg. New York.
- Elliott, N. G., Haskard, K., & Koslow, J. A. (1995). Morphometric analysis of orange roughy (*Hoplostethus atlanticus*) off the continental slope of southern Australia. *Journal of Fish Biology*, 46(2), 202–220. <https://doi.org/10.1111/j.1095-8649.1995.tb05962.x>.
- Fraser M.R, Nys R. de. (2005).The morphology and occurrence of jaw and operculum deformities in cultured barramundi (*Lates calcarifer*) larvae. Aquaculture. Vol. 250. Issue 1-2. Pages 496-503.
- Hair Jr., J.F., Anderson, R., Tatham, R., Black, W., 1996. Multivariate Data Analysis with Readings. Prentice Hall Inc., New Jersey.
- Hardianti, Q., Rusliadi, & Mulyadi. (2016). Effect Of Feeding Made With Different Composition On Growth and Survival Seeds Of Barramundi (*Lates calcarifer* , Bloch ). *Jurnal Online Mahasiswa*, 1–10.
- Hartaty, H., & Setyadji, B. (2016). Parameter Populasi Ikan Tongkol Krai (*Auxis thazard*) Di Perairan Sibolga Dan Sekitarnya. *Jurnal Bawal Widya Riset*

- Perikanan Tangkap, 8(3), 183–190.
- Haryati. (2020). Pemanfaatan Tepung Keong Mas yang Difermentasi Enzim Papain Terhadap Tingkat Kecernaan, Sintasan dan Pertumbuhan Ikan Kakap Putih (*Lates calcarifer*). In *Skripsi*. Fakultas Pertanian. Universitas Muhammadiyah Makassar
- Irmawati, Alimuddin, & Tassakka, A. C. M. A. . (2019). *Dokumen Kebijakan Budidaya Ikan Kakap Putih (Lates calcarifer Bloch, 1790) Berbasis Ekosistem*. December. <https://doi.org/10.13140/RG.2.2.36149.22243>.
- Irmawati, I., Tassakka, A. C. M. A., Alimuddin, A., Nadiarti, N., Husain, A. A. A., Umar, M. T., & Parawansa, B. S. (2020). Identifikasi stok ikan kakap putih (*Lates calcarifer* bloch, 1790) menggunakan karakter morfometrik. *Jurnal IPTEKS Pemanfaatan Sumberdaya Perikanan*, 7(13).
- Irmawati, I., Umar, M. T., Ambo Ala Husain, A., Citra Malina, A., Nurdin Kadir, N., & Alimuddin, A. (2020). Distribution and characteristics of Asian seabass (*Lates calcarifer* Bloch, 1790) in South Sulawesi. *IOP Conference Series: Earth and Environmental Science*, 564(1). <https://doi.org/10.1088/1755-1315/564/1/012011>.
- Irmawati I, Malina AC, Alimuddin A, Kadriah IAK. 2021. Budidaya Ikan Kakap Putih Tinjauan Kelayakan di Keramba Jaring Apung Dan Tambak Tradisional. Makassar. Nas Media Pustaka.
- Iwatsuki, Y. (2013). Review of the *Acanthopagrus latus* complex (Perciformes: Sparidae) with descriptions of three new species from the Indo-West Pacific Ocean. *Journal of Fish Biology* 83: 64–95.
- Jasmine, Rohit, P., Abdussamad, E. M., Koya, K. P. S., Joshi, K. K., Kemparaju, S., Prakasan, D., Elayathu, M. N. K., & Sebastine, M. (2013). Biology and Fishery of The Bullet Tuna. *Auxis rochei*(Risso, 1810) in Indian Waters. Indian. *Journal Fish*, 60(2), 13-20.
- Jerry, D. R. (2013). *Biology and Culture of Asian Seabass*. CRC Press.
- Kristanto, A. H. (2022). Domestikasi Ikan Air Tawar Asli Indonesia Mendukung Produksi Perikanan. Penerbit:BRIN
- Mayunara. (1994). Beberapa Tipe Dan Teori Hermaprodit Pada Ikan Laut. *Oseana*, 19(1), 21–31.
- Mohaddasi, M., Shabanipour, N., & Abdolmaleki, S. (2013). Morphometric variation among four populations of Shemaya (*Alburnus chalcooides*) in the south of Caspian Sea using truss network. *The Journal of Basic & Applied Zoology*, 66(2), 87–92. <https://doi.org/10.1016/j.jobaz.2013.09.001>.
- Mulyono M. 2011. *Budidaya Ikan Kakap Putih (Lates Calcarifer Bloch)*. Jakarta. STP Press.
- Pethiyagoda, R., & Gill, A. C. (2012). Description of two new species of sea bass (Teleostei: Latidae: *Lates*) from Myanmar and Sri Lanka. *Zootaxa*, 16(3314), 1–16. <https://doi.org/10.11164/zootaxa.3314.1.1>.

- Rawat, S., Benakappa, S., Kumar, J., Naik, K., Pandey, G., & Pema, C. W. (2017). Identification of fish stocks based on Truss Morphometric : A review. *Journal of Fisheries and Life Sciences*, 2(1), 9–14.
- Ridho, M. R., & Patriono, E. (2016). Aspek Reproduksi Ikan Kakap Putih (*Lates calcarifer* Bloch) di Perairan Terusan Dalam Kawasan Taman Nasional Sembilang Pesisir Kabupaten Banyuasin. *Jurnal Penelitian Sains*, 18, 1–7.
- Saleky, D., & Dailami, M. (2021). Konservasi Genetik Ikan Kakap Putih ( *Lates calcarifer* Bloch, 1790 ) melalui Pendekatan DNA Barcoding dan Analisis Filogenetik di Sungai Kumbe Merauke Papua. *Jurnal Kelautan Tropis*, 24(2), 141–150.
- Sari, N. F. (2021). Keragaman Genetik dan Abnormalitas Ikan Kakap Putih ( *Lates calcarifer* Bloch, 1790) Tipe Liar dan Hasil Domestikasi. *Skripsi*, FIKP, Universitas Hasanuddin.
- Strauss, R. E., & Bond, C. E. (2016). *Taxonomic Methods: Morphology* In *Methods for Fish Biology*. 109–140.
- Sulistiono, M. Rizki. 2013. Distribusi dan Penyebaran Ikan Kakap Putih. Study aquaculture. [wordpress.com](http://wordpress.com).
- Takács, P., Vitál, Z., Ferincz, Á., & Staszny, Á. (2016). Power and Subjectivity of Different Fish Morphometric Analysis Methods. *Plos One*, 1–16. <https://doi.org/10.1371/journal.pone.0157890>.
- Tarwiyah. 2001. Pedoman Teknis Penanggulangan Penyakit Ikan Budidaya Laut. Direktorat Jenderal Perikanan. Departemen Pertanian. Jakarta
- Tukan, Y. Y. 2022. Studi Morfometrik dan Meristik Ikan Bawah Putih Ekor Pendek (*Pampus chinensis*) Ikan Bawah Putih Ekor Panjang (*Pampus argenteus*) dan Ikan Bawah Hitam (*Parastromateus niger*) Hasil Tangkapan Trawl di Perairan Pulau Bunya. Skripsi, Fakultas Perikanan dan Ilmu Kelautan, Universitas Borneo Tarakan. Tarakan
- Turan, C. (1999). A Note on The Examination of Morphometric Differentiation Among Fish Populations : The Truss System. *Journal of Zoology*, 23(3), 259–263.
- Turan, C. (2004). Stock identification of Mediterranean horse mackerel (*Trachurus mediterraneus*) using morphometric and meristic characters. *Journal of Marine Science*, 781, 774–781. <https://doi.org/10.1016/j.icesjms.2004.05.001>.
- Turan, C., Oral, M., Öztürk, B., & Düzungüneş, E. (2006). Morphometric and meristic variation between stocks of Bluefish (*Pomatomus saltatrix*) in the Black, Marmara, Aegean and northeastern Mediterranean Seas. *Fisheries Research*, 79(1–2), 139–147. <https://doi.org/10.1016/j.fishres.2006.01.015>.
- Utari, S. P. S. D., Panjaitan, F. C. A., & Temaja, I. M. B. B. Morphometric Measurement and Yield of Barramundi Fish Fillet (*Lates calcarifer*) at UD. Damen, Bali. *Berkala Perikanan Terubuk*, 51(3), 1990-1998.
- Wang, C. M., Lo, L. C., Zhu, Z. Y., Pang, H. Y., Liu, H. M., Tan, J., Lim, H. S., Chou, R., Orban, L., & Yue, G. H. (2011). Mapping QTL for an Adaptive Trait : The

- Length of Caudal Fin in *Lates calcarifer* Mapping QTL for an Adaptive Trait : The Length of Caudal Fin in *Lates calcarifer*. *Mar Biotechnol*, 13, 74–82.  
<https://doi.org/10.1007/s10126-010-9271-5>.
- Wardle, C. S., Videler, J. J., & Altringham, J. D. (1995). Tuning In To Fish Swimming Waves: Body Form, Swimming Mode And Muscle Function. *Journal of Experimental Biology*, 198, 1629–1636.
- Wilkinson, L., Blank, G., Gruber, C., 1996. Desktop Data Analysis with SYSTAT. Prentice Hall, Englewood Cliffs, NJ.
- Zairin, M. Jr. 2003. Endokrinologi dan Perannya Bagi Masa Depan Perikanan Indonesia. Orasi Ilmiah Guru Besar FPIK IPB. Bogor.

## **LAMPIRAN**

**Lampiran 1.** Hasil pengukuran ikan kakap putih (*Lates calcarifer* Bloch, 1790) di Tambak Desa Langnga Kecamatan Mattirosompe Kabupaten Pinrang

No	1	2	3	4	5	6	7	8	9	10
SL	49.5	45	46.7	34.5	46.3	34.5	32.1	43.8	43.7	41
A1-A2	14.5	13.8	13.6	10.06	14.09	10.09	10.445	13.79	13.53	13.4
A1-A3	18.8	17.95	17.5	13.4	20.3	12.9	12.75	17.4	15.7	16.3
A1-A5	15.1	14.25	14	10.3	15.9	10.7	10.7	14.45	14	14.55
A1-A4	14.82	14.1	14.2	10.38	14.41	10.41	10.765	14.11	13.85	13.72
A1-A6	3.31	3.27	4.23	2.415	3.565	2.41	2.275	4.89	2.86	4.88
A6-A4	10.65	10.2	9.88	7.22	9.65	7.11	7.585	11.33	9.77	9.98
A6-A7	Ki	1.66	1.8	1.75	1.37	1.9	1.49	1.475	1.79	1.72
	Ka	1.83	1.82	1.75	1.57	1.9	1.49	1.4	1.79	1.93
A8-A9	Ki	1.49	1.66	1.59	1.52	1.8	1.12	1.345	1.5	1.44
	Ka	1.57	1.6	1.68	1.49	1.73	1.48	1.37	1.49	1.58
A1-A10	7.55	6.75	6.66	5.05	6.99	5.71	4.5	6.73	6.465	6.32
A1-A11	Ki	6.22	6.415	6.15	4.37	6.325	4.66	4.535	6.3	5.94
	Ka	6.49	6.44	6.38	4.17	6.33	4.62	4.735	6.44	6.13
B3-B12	20.65	20.9	19.1	14.52	19.7	14.33	14.7	18	20.15	19.35
B3-B5	14.71	14.33	13.16	10.59	14.1	10.7	11	12.94	14.4	13.84
B3-B13	21.35	20.75	19.6	15.4	22.5	15.1	15.5	19.4	20.9	19.65
B3-B14	23.85	23.5	26.5	16.8	23.1	17.2	15.7	21.5	23.5	22.25
B12-B5	25.5	25.7	23.9	19	24.5	18.3	18.1	22.5	23.5	23
B12-B13	11	11.37	10.8	8.45	11.55	8.1	8.5	9.33	10.7	10.44
B12-B14	7.55	8.28	7.4	5.58	7.95	5.55	4.5	6.61	7.7	7.16
B5-B13	19.5	18.6	17.75	14.15	17.55	13.7	11.8	17.55	16.9	16.15
B3	14.22	14.03	13.14	10.36	13.92	10.09	10.22	12.27	14.02	13.74
B4	6.465	6.24	6.135	4.8	6.145	4.3	4.6	5.33	5.64	6.14
B13-B14	5.96	5.55	6.3	4.12	6.24	4.27	3.9	5.6	5.14	5.88
C1-C15	32.3	30.9	30.4	22.9	30.85	23	22.55	30.8	29.4	28.8
C1-C13	34.4	32.7	31.9	24.9	32.7	24.3	23.6	32	34.5	30.5
B12-D16	9.16	7.35	7.8	4.78	7.6	4.66	5.8	7.16	7.41	6.44
B14-D17	7.97	6.6	6.6	4.1	7.5	3.55	4.75	5.83	5.92	6.38
B12-D17	11.7	10.72	10.2	6.49	10.35	6.77	7.65	9.38	10.95	10.56
B14-D16	11.2	10.59	10.1	7.11	10.1	6.27	7.4	9.16	9.55	8.89
D16-D17	7.49	7.8	7	5.16	6.8	4.72	5.35	6.77	6.97	6.92
D18-D19	7.02	7.55	7.55	5.6	7.93	5.59	5.22	6.89	7.33	7.09
D20-D21	10.52	11.03	11.04	9.1	11.43	9.09	8.72	10.39	10.83	10.59

**Lampiran 2.** Hasil pengukuran ikan kakap putih (*Lates calcarifer* Bloch, 1790) di Sungai Salarang Kabupaten Maros

No	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
SL	71.4	35	32	35	34.5	34.5	41.5	39.2	41.9	45.5	27.3	29.5	27.5	27	26	21.7	23	27.5	29.5	30	
A1-A2	31	14.2	13.5	12.9	11.9	12.1	14.9	15.2	16	15	16	12.2	11.2	12.9	14.9	6.5	6.6	8	9	10	
A1-A3	36.3	17.3	16	16	15.6	15.1	18	18.8	18.5	18.6	18	15.6	14.9	16	18	10.3	10.2	12	14.5	15	
A1-A5	27.5	12.2	12.7	12.6	12.6	12.5	15.8	14.2	14.5	14.4	11.2	10.8	12.1	12.2	15.2	8.4	8.1	9.2	10.9	12.4	
A1-A4	42.55	31.845	32.06	32.63	32.43	32.14	39.35	38.25	14.5	13.25	10.5	11.4	10.5	11	38.25	6.9	7.2	9.2	12.8	13	
A1-A6	6	3.9	3.9	3	2.8	2.5	3	2.95	3.2	2.9	2.1	2.4	2.5	3.8	3	2.6	2.6	2.4	3	3.3	
A6-A4	19	8.9	9.2	8.4	9	9	10.5	10.45	10.4	10.4	13	13.6	9	9.2	10.2	5.5	6	5.9	10	9	
A6-A7	Ki	2.5	1.9	1.8	1.4	1.4	1.2	1.7	1.4	1.4	1.3	1.2	1.2	1.2	1.4	0.8	0.9	1.2	1.3	1.4	
	Ka	2.4	1.9	1.9	1.5	1.4	1.3	1.6	1.2	1.5	1.5	1.3	1	1.2	1.2	1.3	0.8	0.8	1.2	1.4	1.3
A8-A9	Ki	2.4	1.3	1.3	1.2	1.7	1	1.5	1.1	1.2	1.1	1	1.1	1.4	1.3	1.2	0.7	0.7	1.2	1.4	1.4
	Ka	2.75	1.15	1.4	1.3	1.1	1.2	1.1	1.55	1.2	1.2	1.1	1	1.2	1.2	1.4	0.7	0.8	1.1	1.2	1.3
A1-A10	10.7	3.35	3.2	3.3	3.4	4.2	6.1	5.9	4.1	6.1	5.7	5.7	5.1	4.3	4	2.3	2.4	3.2	3.4	3.3	
A1-A11	Ki	9.5	5.6	4.3	5	4.9	4.8	5.5	5.5	6.3	5.7	4.5	4.5	4.5	4.5	5.4	3.4	3.1	3.2	4.2	5
	Ka	10	5.75	4.9	5	5.1	4.9	6	5.6	6.5	5.9	4.9	4.8	4.5	4.8	5.2	3.2	3.2	3.1	4.3	5.2
B3-B12	33	13	13.7	14.9	13.9	13.9	17.4	17.6	18.5	21.1	14	15.1	14.2	14.3	16	7.4	9.2	12.4	13.6	14.3	
B3-B5	23	11.5	10.2	11.2	10.7	10.8	12.2	13.4	12.3	13.1	10	10.2	10.5	11.2	13.2	6.9	7.1	10.2	12	12.5	
B3-B13	33.5	13.7	13.9	15.6	14.9	14.9	17.5	18.2	18.3	19.4	13.2	14.7	14.4	14.2	17.6	9.2	10	11.2	14.2	14	
B3-B14	36	14.5	14.9	16.9	16	16.4	20	20.2	20.3	22	15	17.2	15.8	15	19.2	11.2	11.5	11	16.2	17.5	
B12-B5	34.5	17.6	17.3	18.5	17.1	17.8	21.6	22.4	22.2	24.7	17	18.6	17.8	16.2	23.2	12	12.1	12.5	22.9	18	
B12-B13	20.8	8.8	7.2	9.7	8	7.9	9.8	9.1	10	10	7.4	8.1	8.4	7.8	8.6	5.4	5.5	7.8	9	9.2	
B12-B14	36	14.5	14.9	16.9	16	16.4	20	20.2	20.3	22	15	17.2	15.8	15	19.2	11.2	11.5	11	16.2	17.5	
B5-B13	33	11.5	11.7	11.7	11.7	13.4	13.3	15.2	16	15.6	16.9	11.6	13.4	12.4	11.2	15.8	10.2	11	10.4	12.2	13
B3	22.5	10.1	10.5	11.4	11.5	10.5	11.9	12.7	13.2	12.5	9.8	10.5	9.6	9	11.8	5	6.45	7.8	10.2	11	
B4	8.5	4.45	4.3	4.02	4.7	4.8	5.9	5.1	5.1	5.5	4.65	4.95	4.8	4.9	4.85	2.9	3.2	3.4	4.6	5.2	
B13-B14	9.4	5.1	5.1	5.1	4.9	4.9	6.3	6.3	6.3	6.5	3.7	5.5	4.1	3.8	5.8	2.2	2.7	3.2	5.2	5.8	
C1-C15	55	23.8	24.2	24.2	24.1	23.9	29.3	28.6	28.6	29.5	21.9	24.7	23	24	26.6	16.5	17.9	23.5	24	25.8	
C1-C13	57	25	25.5	25.7	25.4	25.2	31	29.8	30.1	32.7	23.1	24.7	23.9	25	28.2	17.2	18.8	22	24	25.2	
B12-D16	11.5	6.6	5	4.9	4.8	5	6.9	4.8	6.3	4.6	4.2	5.5	4.5	4.5	4.6	2.6	3.2	4.2	4.8	5.2	
B14-D17	12	5.9	4.6	4.7	4.7	4.3	5.6	4.4	5.2	5.2	4.7	4.8	4.6	4.8	4	2.2	2.2	3.9	4.2	4.6	
B12-D17	14	8.1	7	7.5	7.8	7.1	9.2	7.6	8.2	6.8	6.3	7.5	6.6	6	7.2	4.7	5	5.8	7.8	8	
B14-D16	16.5	9	7	7.3	7.2	7	9	7.2	8.6	9.2	6.2	7.2	7	7	7.8	4.2	4.5	5	8.2	8	
D16-D17	10.5	6	5	5.2	5.2	5	6.4	6.9	6.5	6.1	4.4	4.9	5	5	5.6	3.2	3.9	4.5	5	5.6	
D18-D19	10.2	6.6	7	5.5	5.3	5.5	6.1	6.2	6.2	6.3	5.2	6	5.7	6.2	6	4.2	4.2	4.8	7	7.2	
D20-D21	13.7	10.1	10.5	9	8.8	9.4	9.6	9.7	10.2	9.8	8.9	9.5	9.2	9.7	9.2	8.5	8.1	8.7	11.2	11.4	

**Lampiran 3.** Hasil perhitungan pengukuran panjang baku ikan kakp putih (*Lates calcarifer* Bloch, 1790) di Tambak Desa Langnga Kecamatan Mattirosompe Kabupaten Pinrang dan di Sungai Salarang Kabupaten Maros

Descriptives				
Lokasi			Statistic	Std. Error
PB Maros		Mean	33.9750	2.43569
		95% Confidence Interval for	Lower Bound	28.8770
		Mean	Upper Bound	39.0730
		5% Trimmed Mean	32.5778	
		Median	31.0000	
		Variance	118.651	
		Std. Deviation	10.89272	
		Minimum	21.70	
		Maximum	71.40	
		Range	49.70	
Pinrang		Interquartile Range	10.80	
		Skewness	2.258	.512
		Kurtosis	6.916	.992
		Mean	41.7100	1.89335
		95% Confidence Interval for	Lower Bound	37.4269
		Mean	Upper Bound	45.9931
		5% Trimmed Mean	41.8111	
		Median	43.7500	
		Variance	35.848	
		Std. Deviation	5.98729	

**Lampiran 4.** Struktur matrix ikan kakap putih (*Lates calcarifer* Bloch, 1790) di Tambak Desa Langnga Kecamatan Mattirosompe Kabupaten Pinrang dan di Sungai Salarang Kabupaten Maros

Structure Matrix

	Function
	1
B12-D17 <sup>a</sup>	-.208
D18-D19 <sup>a</sup>	-.191
D20-D21 <sup>a</sup>	-.186
D16-D17 <sup>a</sup>	-.182
B3-B5 <sup>a</sup>	-.157
A1-A6 <sup>a</sup>	-.142
B3-B14 <sup>a</sup>	-.127
A8-A9 Ka <sup>a</sup>	-.118
B4	-.118
B3-B12 <sup>a</sup>	-.113
B3 <sup>a</sup>	-.111
A6-A7 Ki <sup>a</sup>	-.107
A1-A11 Ki <sup>a</sup>	-.105
B3-B13	-.104
B12-B5 <sup>a</sup>	-.098
C1-C13 <sup>a</sup>	-.096
A6-A7 Ka <sup>a</sup>	-.094
A1-A5 <sup>a</sup>	-.093
A8-A9 Ki <sup>a</sup>	-.089
B12-B13	-.083
B14-D17	-.078
A1-A11 Ka <sup>a</sup>	-.075
C1-C15 <sup>a</sup>	-.068
B12-B14	.068
A1-A4 <sup>a</sup>	-.055
A1-A2	-.048
B14-D16 <sup>a</sup>	-.045
A1-A3 <sup>a</sup>	-.043
B12-D16 <sup>a</sup>	-.041
A1-A10 <sup>a</sup>	-.026
B5-B13 <sup>a</sup>	-.017
A6-A4 <sup>a</sup>	-.008
B13-B14 <sup>a</sup>	.004

