

## **sDAFTAR PUSTAKA**

- Smith-Bindman R, Lipson J, Marcus R, Kim K P, Mahesh N, Gould R, de Gonzalez A B and Miglioretti D L 2009 Arch Intern Med. 169(22) 2078–2086
- Linet M, Kim K and Rajaraman P 2009 Pediatr Radiol. 39(S1) S4–S26
- Power S P, Moloney F, Twomey M, James K, O'Connor O J and Maher M M 2016 World J Radiol. 8(12) 902–915
- Toori A J, Shabestani-Monfared A, Deeband M R, Abdi R and Nabahati M 20ss15 J Biomed Phys Eng. 5(4) 177–184
- American Association of Physicist in Medicine (AAPM) 2011 AAPM Report 204.
- Kopp M., Loewe T., Wuest W. et al 2020 Sci Rep 10, 9475
- Internasioanl commission on radiation PROTECTION. *Managing patient dose in computed tomography*. ICRP publication 87. Ann ICRP, 2000;33:4
- Toth, T. L. *Dose Reduction Opportunities For Ctscanner.Pediatric Radiology*. 32 (4): 261-267,2002
- C. Anam, F. Haryanto, R. Widita, I. Arif, andG. Dougherty, “*The evaluation of the effective diameter (Deff) calculation and its impact on the size-specificdose estimate (SSDE)*, ” Atom Indonesia,43, 55-60 (2017)

