

## REFERENSI

1. Canales C, Elsayes A, Yeh DD, Belcher D, Nakayama A, McCarthy CM, et al. Nutrition Risk in Critically Ill Versus the Nutritional Risk Screening 2002: Are They Comparable for Assessing Risk of Malnutrition in Critically Ill Patients? *Journal of Parenteral and Enteral Nutrition.* 2019 Jan 1;43(1):81–7.
2. Rattanachaiwong S, Zribi B, Kagan I, Theilla M, Heching M, Singer P. Comparison of nutritional screening and diagnostic tools in diagnosis of severe malnutrition in critically ill patients. *Clinical Nutrition.* 2020 Nov 1;39(11):3419–25.
3. C. O. Schell et al., “The global need for essential emergency and critical care,” *Crit. Care*, vol. 22, no. 1, pp. 1–5, 2018, doi: 10.1186/s13054-018-2219-2
4. Sharma K, Mogensen KM, Robinson MK. Pathophysiology of Critical Illness and Role of Nutrition. Vol. 34, *Nutrition in Clinical Practice*. John Wiley and Sons Inc; 2019. p. 12–22.
5. Griffith DM, Lewis S, Rossi AG, Rennie J, Salisbury L, Merriweather JL, et al. Systemic inflammation after critical illness: relationship with physical recovery and exploration of potential mechanisms. *Thorax.* 2016 Sep;71(9):820–9.
6. Cox CE. Persistent Systemic Inflammation in Chronic Critical Illness. *Respir Care.* 2012 Jun;57(6):859–66.
7. Brummel NE, Hughes CG, Thompson JL, Jackson JC, Pandharipande P, McNeil JB, et al. Inflammation and Coagulation during Critical Illness and Long-Term Cognitive Impairment and Disability. *Am J Respir Crit Care Med.* 2021 Mar 15;203(6):699–706.
8. Wendel Garcia PD, Fumeaux T, Guerci P, Heuberger DM, Montomoli J, Roche-Campo F, et al. Prognostic factors associated with mortality risk and disease progression in 639 critically ill patients with COVID-19 in Europe: trial report of the international RISC-19-ICU prospective observational cohort. *EClinicalMedicine.* 2020 Aug;25:100449.



9. Cox CE. Persistent Systemic Inflammation in Chronic Critical Illness. *Respir Care.* 2012 Jun;57(6):859–66.
10. Altay S, Gürdoğan M, Keskin M, Kardaş F, Çakır B. The Inflammation-Based Glasgow Prognostic Score as a Prognostic Factor in Patients with Intensive Cardiovascular Care Unit. *Medicina (B Aires).* 2019 May 15;55(5):139.
11. Howard R, Kanetsky PA, Egan KM. Exploring the prognostic value of the neutrophil-to-lymphocyte ratio in cancer. *Sci Rep.* 2019 Dec 1;9(1).
12. Salciccioli JD, Marshall DC, Pimentel MAF, Santos MD, Pollard T, Celi AA, et al. The association between the neutrophil-to-lymphocyte ratio and mortality in critical illness: An observational cohort study. *Crit Care.* 2015 Jan 19;19(1).
13. Wu X, Luo Q, Su Z, Li Y, Wang H, Liu Q, et al. Neutrophil-to-lymphocyte ratio as a predictor of mortality in intensive care unit patients: A retrospective analysis of the Medical Information Mart for Intensive Care III Database. *BMJ Open.* 2021 Nov 11;11(11).
14. Al-Mazedi MS, Rajan R, Al-Jarallah M, Dashti R, Al Saber A, Pan J, et al. Neutrophil to lymphocyte ratio and in-hospital mortality among patients with SARS-CoV-2: A retrospective study. *Annals of Medicine and Surgery.* 2022 Oct 1;82.
15. Ali HS, Ananthegowda DC, Ebrahim EMA, Kannappilly N, Al Wraidiat M, Mohamed AS, et al. Neutrophil-to-lymphocyte ratio as a predictor of clinical outcomes in critically ill COVID-19 patients: A retrospective observational study. *Health Sci Rep.* 2022 Sep 1;5(5).
16. Tekinalp A. The Association Between Lymphocyte/Neutrophil Ratio and Clinical Course in Intensive Care Patients. *Eurasian Journal of Medical Investigation.* 2018;
17. Lu Y, Ren C, Jiang J. The relationship between prognostic nutritional index and all-cause mortality in critically ill patients: A retrospective study. *Int J Clin Med.* 2021;14:3619–26.



18. Cheng YL, Sung SH, Cheng HM, Hsu PF, Chao- ;, Guo Y, et al. Prognostic Nutritional Index and the Risk of Mortality in Patients With Acute Heart Failure. Available from: <http://ahajournals.org>
19. Huang Y, Zhang Q, Li P, Chen M, Wang R, Hu J, et al. The prognostic nutritional index predicts all-cause mortality in critically ill patients with acute myocardial infarction. *BMC Cardiovasc Disord.* 2023 Dec 1;23(1).
20. Li T, Qi M, Dong G, Li X, Xu Z, Wei Y, et al. Clinical Value of Prognostic Nutritional Index in Prediction of the Presence and Severity of Neonatal Sepsis. *J Inflamm Res.* 2021;14:7181–90.
21. Al-Shami I, Hourani HMA, Alkhatib B. The use of prognostic nutritional index (PNI) and selected inflammatory indicators for predicting malnutrition in COVID-19 patients: A retrospective study. *J Infect Public Health.* 2023 Feb 1;16(2):280–5.
22. Wang ZH, Lin YW, Wei XB, Li F, Liao XL, Yuan HQ, et al. Predictive Value of Prognostic Nutritional Index on COVID-19 Severity. *Front Nutr.* 2021 Jan 14;7.
23. Oh TK, Song IA, Lee JH. Clinical usefulness of C-reactive protein to albumin ratio in predicting 30-day mortality in critically ill patients: A retrospective analysis. *Sci Rep.* 2018 Dec 1;8(1).
24. Park JE, Chung KS, Song JH, Kim SY, Kim EY, Jung JY, et al. The C-reactive protein/albumin ratio as a predictor of mortality in critically ill patients. *J Clin Med.* 2018 Oct 1;7(10).
25. Mohamed SA, Elhawary R. C-Reactive Protein/Albumin Ratio as an Independent Predictor of Mortality in Critically Ill Pediatric Patients. *Journal of Child Science.* 2020 Jan 1;10(1):E1–11.
26. Gyawali P, Shrestha H, Pant V, Risal P, Gautam S. C-reactive protein to albumin ratio among patients admitted to intensive care unit of a tertiary care hospital: A descriptive cross-sectional study. *Journal of the Nepal Medical Association.* 2021 Dec 11;59(244):1247–51.



27. Buonacera A., Stancanelli B., Colaci M., Malatino L. Neutrophil to Lymphocyte Ratio: An Emerging Marker of the Relationships between the Immune System and Diseases. *Int J Mol Sci.* 2022 Mar 26;23(7):3636.
28. Drăgoescu A.N., Pădureanu V., Stănculescu A.D., Chiuțu L.C., Tomescu P., Geormăneanu C., Pădureanu R., Iovănescu V.F., Ungureanu B.S., Pănuș A., et al. Neutrophil to Lymphocyte Ratio (NLR)—A Useful Tool for the Prognosis of Sepsis in the ICU. *Biomedicines.* 2021;10:75. doi: 10.3390/biomedicines10010075.
29. Gurol G., Ciftci I.H., Terzi H.A., Atasoy A.R., Ozbek A., Koroglu M. Are There Standardized Cutoff Values for Neutrophil-Lymphocyte Ratios in Bacteremia or Sepsis? *J. Microbiol. Biotechnol.* 2015;25:521–525. doi: 10.4014/jmb.1408.08060.
30. Liu S., Wang X., She F., Zhang W., Liu H., Zhao X. Effects of Neutrophil-to-Lymphocyte Ratio Combined with Interleukin-6 in Predicting 28-Day Mortality in Patients with Sepsis. *Front. Immunol.* 2021;12:639735.
31. Jang J.Y., Yoo G., Lee T., Uh Y., Kim J. Identification of the robust predictor for sepsis based on clustering analysis. *Sci. Rep.* 2022;12:2336.
32. Lee H., Kim I., Kang B.H., Um S.-J. Prognostic value of serial neutrophil-to-lymphocyte ratio measurements in hospitalized community-acquired pneumonia. *PLoS ONE.* 2021;16:e0250067.
33. Shah N., Parikh V., Patel N., Patel N., Badheka A., Deshmukh A., Rathod A., Lafferty J. Neutrophil lymphocyte ratio significantly improves the Framingham risk score in prediction of coronary heart disease mortality: Insights from the National Health and Nutrition Examination Survey-III. *Int. J. Cardiol.* 2014;171:390–397.
34. Sun X., Luo L., Zhao X., Ye P., Du R. The neutrophil-to-lymphocyte ratio on admission is a good predictor for all-cause mortality in hypertensive
35. Arvanitakis K., Mitroulis I., Germanidis G. Tumor-Associated Neutrophils in Hepatocellular Carcinoma Pathogenesis, Prognosis, and Therapy. *Cancers.* 2021;13:2899. doi: 10.3390/cancers13122899.



36. Silberman S., Abu-Yunis U., Tauber R., Shavit L., Grenader T., Fink D., Bitran D., Merin O. Neutrophil-Lymphocyte Ratio: Prognostic Impact in Heart Surgery. Early Outcomes and Late Survival. *Ann. Thorac. Surg.* 2018;105:581–586. doi: 10.1016/j.athoracsur.2017.07.033.
37. Hajibandeh S., Hajibandeh S., Hobbs N., Mansour M. Neutrophil-to-lymphocyte ratio predicts acute appendicitis and distinguishes between complicated and uncomplicated appendicitis: A systematic review and meta-analysis. *Am. J. Surg.* 2020;219:154–163.
38. Yang L, Gao C, He Y, Wang X, Yang L, Guo S, Chen J, He S, Sun Y, Gao Y, Guo Q. The Neutrophil-to-Lymphocyte Ratio is Associated with the Requirement and the Duration of Invasive Mechanical Ventilation in Acute Respiratory Distress Syndrome Patients: A Retrospective Study. *Can Respir J.* 2022 Jul 16;2022:1581038.
39. Gracia Lizbeth Velazquez, Obed Isaí Aguilera Olvera, Gladis Lucy Arcos, Jose Camilo Muñoz Chaves, German Antonio Aguirre-Gomez, Villalobos Silva Jose Antonio, Correlation between neutrophil-lymphocyte ratio and mechanical ventilation in acute respiratory distress syndrome, *Journal of Critical Care*, Volume 81, 2024.
40. Sun, W., Luo, Z., Jin, J., Cao, Z., & Ma, Y. (2021). The Neutrophil/Lymphocyte Ratio Could Predict Noninvasive Mechanical Ventilation Failure in Patients with Acute Exacerbation of Chronic Obstructive Pulmonary Disease: A Retrospective Observational Study. *International Journal of Chronic Obstructive Pulmonary Disease*, 16, 2267–2277.
41. Chen L, Bai P, Kong X, Huang S, Wang Z, Wang X, et al. Prognostic Nutritional Index (PNI) in Patients With Breast Cancer Treated With Neoadjuvant Chemotherapy as a Useful Prognostic Indicator. *Front Cell Dev Biol.* 2021;9:656741.
42. Kang, S. H. *et al.* Prevalence and prognostic significance of malnutrition in patients with acute coronary syndrome treated with percutaneous coronary intervention. *Medicine (Baltimore)* **101**, e30100 (2022).



43. Zhang, H. *et al.* The predictive value of a preoperative systemic immune-inflammation index and prognostic nutritional index in patients with esophageal squamous cell carcinoma. *J. Cell. Physiol.* **234**, 1794–1802 (2019)
44. Qiu, C. *et al.* Evaluation of prognostic nutritional index in patients undergoing radical surgery with nonsmall cell lung cancer. *Nutr. Cancer* **67**, 741–747 (2015)
45. Lu Y, Ren C, Jiang J. The Relationship Between Prognostic Nutritional Index and All-Cause Mortality in Critically Ill Patients: A Retrospective Study. *Int J Gen Med.* 2021 Jul 17;14:3619-3626.
46. Wany M, Helby J, Nordestgaard BG, Birgens H, Bojesen SE. Lymphopenia and risk of infection and infection-related death in 98,344 individuals from a prospective Danish population-based study. *PLoS Med.* 2018 Nov;15(11):e1002685
47. Baker JR, Donnelly LE. Leukocyte function in COPD: clinical relevance and potential for drug therapy. *Int J Chron Obstruct Pulmon Dis.* 2021;16:2227–2242.
48. Divo MJ, Celli BR, Poblador-Plou B, Calderon-Larranaga A, de Torres JP, Gimeno-Feliu LA, et al. Chronic Obstructive Pulmonary Disease (COPD) as a disease of early aging: evidence from the EpiChron Cohort. *PLoS ONE.* 2018;13(2):e0193143.
49. Wu X, Luo Q, Su Z, Li Y, Wang H, Liu Q, et al. Neutrophil-to-lymphocyte ratio as a predictor of mortality in intensive care unit patients: a retrospective analysis of the Medical Information Mart for Intensive Care III Database. *BMJ Open.* 2021;11(11):e053548.
50. Hu Y, Long H, Cao Y, Guo Y. Prognostic value of lymphocyte count for in-hospital mortality in patients with severe AECOPD. *BMC Pulm Med.* 2022 Oct 5;22(1):376.
51. Gutierrez-Rodriguez, R., Aguilar-Alonso, E., Arias-Verdu, M. *et al.* Lymphopenia assessment in ICU patients and relationship with mortality. *ICMx* **3** (Suppl 1), A340 (2015)



52. Adrie, C., Lugosi, M., Sonnevile, R. *et al.* Persistent lymphopenia is a risk factor for ICU-acquired infections and for death in ICU patients with sustained hypotension at admission. *Ann. Intensive Care* 7, 30 (2017)
53. Fang Y, Zheng T, Zhang C. Prognostic Role of the C-Reactive Protein / Albumin Ratio in Patients With Gynecological Cancers: A Meta-Analysis. *Front Oncol.* 2021;11:737155.
54. Wei X.L., Wang F.H., Zhang D.S., Qiu M.Z., Ren C., Jin Y., Zhou Y.X., Wang D.S., He M.M., Bai L., et al. A novel inflammation-based prognostic score in esophageal squamous cell carcinoma: The C-reactive protein/albumin ratio. *BMC Cancer.* 2015;15:350–361.
55. Kinoshita A., Onoda H., Imai N., Iwaku A., Oishi M., Tanaka K., Fushiyama N., Koike K., Nishino H., Matsushima M. The C-reactive protein/albumin ratio, a novel inflammation-based prognostic score, predicts outcomes in patients with hepatocellular carcinoma. *Ann. Surg. Oncol.* 2015;22:803–810.
56. Wu M., Guo J., Guo L., Zuo Q. The C-reactive protein/albumin ratio predicts overall survival of patients with advanced pancreatic cancer. *Tumour Biol.* 2016;37:12525–12533.
57. Park JE, Chung KS, Song JH, Kim SY, Kim EY, Jung JY, Kang YA, Park MS, Kim YS, Chang J, Leem AY. The C-Reactive Protein/Albumin Ratio as a Predictor of Mortality in Critically Ill Patients. *J Clin Med.* 2018 Oct 8;7(10):333.
58. Ranzani O.T., Zampieri F.G., Forte D.N., Azevedo L.C., Park M. C-reactive protein/albumin ratio predicts 90-day mortality of septic patients. *PLoS ONE.* 2013;8.



## Lampiran 1. Rekomendasi Etik



KEMENTERIAN PENDIDIKAN, KEBUDAYAAN, RISET DAN TEKNOLOGI  
 UNIVERSITAS HASANUDDIN FAKULTAS KEDOKTERAN  
 KOMITE ETIK PENELITIAN UNIVERSITAS HASANUDDIN  
 RSUP Dr. WAHIDIN HIDAYAH MAKASSAR  
 Sekretariat : Lantai 2 Gedung Laboratorium Terpadu  
 AL PERINTAH KEPERENCANAAN KAMPUS TAHUNALAMBAR KM. 30 MAKASSAR 90243.  
 Contact Person: dr. Aqilahin Suharto, MM, PhD, SpMK, Telp. 0911 2178181, Fax. 041-561403



### REKOMENDASI PERSETUJUAN ETIK

Nomor : 169/UN4.6.4.5.33/1936/2023

Tanggal : 19 Desember 2023

Dengan ini Mewysatakan bahwa Protokol dan Dokumen yang Berhubungan Dengan Protokol berikut ini telah mendapatkan Persetujuan Etik :

No Protokol	URG23110838	No Sponsor	
Peneliti Utama	dr. Yella Wullur	Sponsor	
Judul Penelitian	Analisa Parameter Laboratorium Gizi (NUL, PNL, CRP, Albunin Ratio) Terhadap Status Klinis Pasien ICU		
No Verisi Protokol	I	Tanggal Verisi	8 Nopember 2023
No Verisi PSP		Tanggal Verisi	
Tempat Penelitian	RS Wahidin Sudirohusodo Makassar		
Jenis Review	<input type="checkbox"/> Exempted <input checked="" type="checkbox"/> Expedited <input type="checkbox"/> Fullboard Tanggal	Masa Berlaku 19 Desember 2023 sampai 19 Desember 2024	Frekuensi review lanjutan
Ketua KEP Universitas Hasanuddin	Nama Prof. dr. Muhib Nurrozi, MM, PhD, SpMK, Subsp. Bakti(K)	Tanda tangan	
Sekretaris KEP Universitas Hasanuddin	Nama dr. Firdaus Hanifid, PhD, SpMK(K)	Tanda tangan	

Ketentuan Penelitian Utama:

- Mengembalikan atau mendukung Protokol untuk persetujuan sebelum di implementasikan.
- Mengembalikan Laporan SAR ke Komite Etik dalam 24 jam dan diberi waktu 7 hari dan Lajur SEIAK dalam 72 jam sejak Peneliti Utama menerima laporan.
- Mengembalikan Laporan Komittee (progress report) setiap 6 bulan untuk posisi risiko tinggi dan setiap setahun untuk penelitian risiko rendah.
- Mengembalikan laporan akhir setelah Penelitian berakhir.
- Melaporkan perubahan dari protokol yang diterapkan (protocol deviation / violation).
- Memantau semua peraturan yang ditentukan.

