

The Influence of Fixed Orthodontic Treatment on Tooth Discoloration among Dental Students in Makassar, Indonesia

Puspitasari Yustisia¹, Mohammad Dharma Utama², Bachtiar, Rachmi³, Harun Achmad⁴, Ilmianti⁵, Selviani, Yusrini⁶, Eva Novawati⁷, Fachira Jamila Rusdi⁸

^{1,7}Department of Orthodontics, Faculty of Dentistry, Moslem University of Indonesia

²Department of Prosthodontics, Faculty of Dentistry, Hasanuddin University

³Department of Periodontics, Faculty of Dentistry, Moslem University of Indonesia

⁴Department of Pedodontics, Faculty of Dentistry, Hasanuddin University

⁵ Department of Dental Public Health, Faculty of Dentistry, Moslem University of Indonesia

⁶ Department of Oral Medicine, Faculty of Dentistry, Moslem University of Indonesia

⁷Student of Dentistry Study, Faculty of Dentistry, Moslem University of Indonesia

*Corresponding Author: harunachmader@gmail.com

ABSTRACT

The aims of fixed orthodontic treatment is to improve mastication function, aesthetics, prevent tissue damage and restore the normal function of oral cavity. Fixed orthodontic treatment could led poor oral hygiene for the patient because it could be a good place for residual food, led plaque accumulation under the appliance. Furthermore, plaque accumulation could led stain on tooth surface and tooth discoloration could develop. This research was to investigate the influence of fixed orthodontic treatment on tooth discoloration among Dental Students in Makassar, Indonesia. The research was conducting as an analytic observational with cross-sectional study design Sample of this research were 1st and 2nd year of dental students of Faculty of Dentistry, Universitas Muslim Indonesia in Makassar, Indonesia who were under fixed orthodontic treatment. Samples were divided into 2 groups, < 1 year and \geq 2 years of fixed orthodontic treatment. Intraoral photograph from frontal view were taken, and outline of the labial surfaces of all the incisors were enlarged to scale (magnification \times 4) and each tooth face divided into 4-mm squares. Labial surface of the incisors were drawn by the examiner on to the grid system using photoshop cc 2020. Data was statistically analyzed using Fisher exact test. The p-value of this research was less than 0.05, therefore the results of this research are statistically significant. Hypothesis was accepted and showed that there was an influence of fixed orthodontic treatment on tooth discoloration among dental students in Makassar, Indonesia.

Keywords: fixed orthodontic treatment, tooth discoloration, grid system

Introduction

The main aim of orthodontic treatment is to correct malocclusion, in order, whenever possible, to achieve functionally appropriate occlusion and optimum dental and facial aesthetics. "Orthodontic treatment need" can be defined as the degree to which a person needs orthodontic treatment because of certain features of his or her malocclusion, the functional, dental health or aesthetic impairment it occasions and the negative psychological and social repercussions to which it gives rise.^[1]

The use of orthodontic devices today, both fixed and removable, has been widely used by the society, adults, and especially students. Amalia (2009) stated that patients with fixed orthodontic had less

knowledge about the treatment. The patients were not aware of the side effects of using fixed orthodontics, and those related to poor oral hygiene can worsen the root resorption and bone resorption, which can slow down the treatment. Lack of knowledge about the side effects of orthodontic treatment will cause the patients to be unaware of the side effects caused by the use of orthodontics devices. This will complicate the oral hygiene procedure in orthodontic patients. ^{[2],[3],[4]}

The benefits of a medical intervention must outweigh any potential harm. Apart from its benefits, orthodontic treatment can cause certain complications. During clinical orthodontic procedures, the use of various methods, devices, and materials can cause unwanted side effects, both local (e.g., tooth discoloration, decalcification, root resorption, and periodontal complications) and systemic (e.g., allergic reactions and crossinfections).^[5]

Fixed orthodontic treatment can be a place for plaque accumulation due to increased biofilm formation after insertion. A research conducted by Jayanti (2017) mentioned that there was a relationship between the duration of the fixed orthodontic treatment and periodontal treatment needs. The effects of periodontal disease associated with fixed orthodontic use were gingivitis, periodontitis, gingival recession or hypertrophy, and alveolar bone loss. Periodontal complications that may appear during orthodontic treatment are gingivitis, periodontitis, gingival recession or hypertrophy, alveolar bone loss, dehiscences, fenestrations, interdental folds, and dark triangles. ^{[6],[7],[8],[9]}

Plaque also attacks the hard tissue of the teeth. This corresponds with the research conducted by Un-Bong (2017) stating that there was clinical development of dental discoloration during orthodontic treatment. Besides causing periodontal disease, plaque formed from fixed orthodontic devices could also leave colored marks on the surface of the tooth. This statement was supported by the results of research conducted by Jufri (2017), stating that there was a relationship between fixed orthodontic treatment and dental discoloration.^{[10],[11]}

Color alterations after orthodontic treatment are the result of several factors. More severe alterations occur when chemically cured resins are used as bonding materials rather than light cured composites. The resin tags cannot be removed by cleaning procedures without altering the enamel's surface. Irreversible changes occur to the enamel's surface morphology, rugosity, and texture, with negative consequences on its reflective properties, luminosity, and optical perception. Evidence shows that the adhesive resins used for bracket bonding do not present good color stability with time. Ultraviolet light and corrosion products from the orthodontic appliance combined with food dyes induce color alterations, with a tendency to move toward the yellow tones. Orthodontic forces induce variations in pulp vascularization. It is a possible factor for the premature aging of teeth, and it is an endogenous factor for discoloration too. In situations where white spots and lesions are present, even if remineralization occurs,

<http://annalsofrscb.ro> 10721

the outcome may be different from the initial enamel structure. This is because of the differences in the minerals in the treated enamel surface compared to the untreated enamel.^{[12],[13],[14],[15],[16],[17],[18]}

Leftover food, ultraviolet light, and corrosion products that come from fixed orthodontic devices affect discoloration with the tendency for teeth to turn into the yellowish color. According to the theory the orthodontic treatment period varies according to the difficulty of the case. The average of estimated treatment time is 2 years but in fact the treatment time is often 50% longer than the predicted time, and it usually occurs in adolescents.^{[19],[20]}

Based on the above, the researcher intends to conduct a research on “The Influence of Fixed Orthodontic Treatment on Tooth Discoloration among Dental Students in Makassar, Indonesia”

Research Methods

This research used observational analytic study design with cross-sectional study. This research was conducted at FKG-UMI Makassar in October-November 2019. The samples used were 23 people using fixed orthodontics in the upper and lower jaws with a purposive sampling method in accordance with the criteria determined by the writers. The data were processed by using SPSS version 25 with the use of fisher exact test.

Research Result

A research has been conducted about the Influence of fixed orthodontic treatment on tooth discoloration among Dental students in Makassar, Indonesia. The data obtained from this study consisted of 23 samples of fixed orthodontic treatment users. The data obtained are as follows:

Table 5.1.1 Average Length of Fixed Orthodontic Treatment

Duration of Treatment (month)	Frequency	Percentage
> 12	15	65.2
≤ 12	8	34.8
Total	23	100

Table 5.1.1 shows the average length of fixed orthodontic treatment of 23 samples. There are 15 users of fixed orthodontic treatment with treatment duration >12 months (65.2%) and 8 users of fixed orthodontic treatment with treatment duration ≤12 months (34.8%).

Table 5.1.2 Degrees of Dental Discoloration

Dental Discoloration	Frequency	Percentage
0	9	39.1
1	12	52.2
2	2	8.7
3	0	0
4	0	0
Total	23	100

Table 5.1.2 shows the degrees of dental discoloration of 23 samples. 12 samples (52.2%) have the discoloration degree of 1, and 2 samples (8.7%) have the discoloration degree of 2.

Table 5.1.3 The Influence of Orthodontic Treatment on Dental Discoloration

Duration of Treatment (month)	Dental Discoloration						Total	P-value	
	0		1		2				
	N	%	n	%	n	%			
> 12	3	20	10	66.7	2	13.3	15	100	0.033
≤ 12	6	75	2	25	0	0	8	100	
Total	9	39.1	12	52.2	2	8.7	23	100	

Table 5.1.3 shows that out of 15 samples with treatment duration >12 months, 10 samples (66.7%) are in discoloration degrees 1, and 2 samples (13.3%) are in discoloration degrees 2. Meanwhile, out of 8 samples with treatment duration \leq 12 months, 6 samples (75%) are in the discoloration degree 0, and 2 samples (25%) are in discoloration degree 1.

From table 5.1.3, the P-value obtained from statistical result fisher exact Test is 0.033, or less than 0.05 which means that there is a significant influence between the duration of fixed orthodontic treatment with dental discoloration.



Figure 1. An example of tooth discoloration on maxillary and mandibular incisors.

Discussion

The results of this study with P-value obtained from statistical result fisher exact test equal to 0.033 indicate that there is a significant influence to the results of this research. Thus, the alternative hypothesis (H_a) is accepted and the null hypothesis (H_0) is rejected. The highest level of dental discoloration is those with treatment duration >12 months with discoloration degree 1 (table 5.1.3). This correlates with Jufri's research (2017), which stated that there was a significant relationship between fixed orthodontic treatment and dental discoloration.

In table 5.1.1, it is found that the average length of fixed orthodontic treatment >12 months is used by 15 people (65.2%). Theoretically, the duration of fixed orthodontic treatment varies according to the difficulty of this case. The estimated average treatment period is 2 years but in reality it might take 50% longer than expected that usually occurs in adolescents and may be longer in adults due to more difficult cases. This is supported by the research conducted by Yovela (2009) at Faculty of Dentistry, Universitas Indonesia which showed that adult patients needed longer periods of treatment.^{[21],[22]}

In fact, the question reveals a strong desire, especially by adult patients, for shorter treatment, since the anti-aesthetic look provided by orthodontic brackets in addition to longer correction time are the major factors responsible for demotivating patients to have treatment begin.^[23]

In the literature, no consensus has been reached about orthodontic treatment time. A recent systematic review revealed mean treatment time with fixed appliances of 19.9 months. However, there was significant variation among studies (with mean values ranging from 14 to 33 months), and the quality of treatment outcomes was not assessed. Whenever cases were assessed under the American Board of Orthodontics (ABO) standards, one-phase orthodontic treatment mean time was 24.6 months.^{[24],[25],[26]}

On the other hand, orthodontic treatment meantime seems to be beyond patients' expectation. When asked about how long they would like treatment to last, 40.8% of adolescent patients answered less than 6 months, while 33.2% of them answered between 6 and 12 months. Among adult patients, 42.9% answered between 6 and 12 months, while 26.5% answered between 12 and 18 months.^[27]

Treatment duration would be almost 25 months if there were no missed appointments, no repairs were required, and only 1 instructor supervised the treatment. For any missed appointment treatment duration increased 2.67 months; for every orthodontic appliance issue/breakage treatment time increased 1.14 months; and for each different instructor it increased 6.6 months.^[28]

Extremely long treatment time has been associated with greater susceptibility to iatrogenesis, which in turn are associated with orthodontic appliances. This is the case of root resorption, white spots, carious lesions, and gingival inflammation.^[29]

Table 5.1.2 regarding the degree of dental discoloration shows that it is on the degree of discoloration 1 in which 0.01% to 25% of the area is discolored. This result corresponds with Amalia's research (2009) which stated that the level of patient knowledge about fixed orthodontic treatment is categorized as low. The patients were unaware of the side effects from the use of fixed orthodontics,

especially those related to poor oral hygiene that can worsen the root resorption and bone resorption, which can slow down the treatment. Lack of knowledge about the side effects of orthodontic treatment will cause the patients to be unaware of the side effects caused by the use of orthodontics devices. This will complicate the oral hygiene procedure in orthodontic patients. Fixed orthodontic treatment can be a place for plaque accumulation due to increased biofilm formation after insertion. A research conducted by Jayanti (2017) mentioned that there was a relationship between the duration of the fixed orthodontic treatment and periodontal treatment needs. The effects of periodontal disease associated with fixed orthodontic use were gingivitis, periodontitis, gingival recession or hypertrophy, and alveolar bone loss. Plaque is the most important factor in the initiation, development, and recurrence of periodontal disease.

[2],[3],[4],[6],[30]

A research conducted by Sepideh Arab, et. al. in (2016) found that the intricate design of fixed orthodontic devices affects oral hygiene by influencing several factors, such as changes in salivary pH. Changes in salivary pH in the oral cavity cause demineralization in the enamel layer of the tooth which increases susceptibility to dental caries. Discoloration after fixed orthodontic treatment is influenced by multifactorial. Leftover food, ultraviolet light, and corrosion products that come from fixed orthodontic devices affect discoloration with the tendency for teeth to turn yellowish. Some of the variables causing dental discoloration are directly related to orthodontic treatment techniques. ^{[11],[19],[31],[33],[34]}

The frequency of dental discoloration can be much higher depending on the severity of the malocclusion, and fixed orthodontic devices have greater effects than removable orthodontic devices due to the use of resin material used to attach the brackets to the enamel. A research conducted by Huang et al (2013) stated that there were few data related to the incidence of dental discoloration. A research conducted by Un-Bong (2017) showed that there was a clinical development on dental discoloration during the orthodontic treatment. Besides causing periodontal disease, plaque formed from fixed orthodontic devices could also leave color marks on the surface of the tooth. ^{[10],[19],[31],[32],[34]}

A research conducted by Sandison (1981) and Fleming (1982) showed that enamel discoloration was influenced by enamel bonding and debonding procedures. The change in color of the enamel could originate from the post-debonding resin removal protocol, and the penetration of the resin tag into the enamel structure at depths reaching 50 mm. Impregnation of the resin into the enamel structure could not be reversed by debonding and cleaning procedures, and enamel discoloration could occur by direct absorption of food coloring and products from corrosion in orthodontic devices. ^{[35],[36]}

The relationship between dental discoloration and fixed orthodontic treatment is still controversial. Some researchers have concluded that performing bonding and debonding procedures do not seem to have a significant effect on dental enamel color. Other researchers have shown that dental discoloration will occur when using this procedure. This systematic review is to evaluate the dental discoloration on enamel after the use of orthodontic resin and its cleaning procedures, and it is an interesting thing to learn. ^[36]

According to Rammadhan et al. (2016), one of the efforts to prevent and overcome dental health problems was through the dental health education approach. Dental health education delivered to someone is expected to be able to change his/her behavior and gain more knowledge about individual dental health from unhealthy behavior towards healthy behavior. Good dental health and oral hygiene are also influenced by the role of the dentist in providing motivation and instruction during fixed orthodontic treatment. Dentists need to provide proper motivation and instruction from the beginning to the end of fixed orthodontic treatment so that the patients are more concerned about the health of their teeth and mouth. In this study, however, the degree of dental discoloration of most students at Faculty of Dentistry, Universitas Muslim Indonesia is generally good because of the influence of knowledge acquired while undergoing their studies in the dentistry program. This can also increase students' knowledge about dental care and dental discoloration, especially when having orthodontic treatment. ^{[37],[38], [39], [40], [41], [42], [43]}

Conclusion

Based on the research that has been done, it can be concluded that there is a significant influence between the duration of fixed orthodontic treatment on dental discoloration among dental students in Makassar, Indonesia.

REFERENCES

- [1] Carlos Bellot-Arcís, José María Montiel-Company and José Manuel Almerich-Silla. Orthodontic treatment need: an epidemiological approach. Stomatology Department, University of Valencia Spain. 2012
- [2] Herwanda, et al. Knowledge of teenager with years 15-17 old in Senior High School 4 Banda Aceh city in using fixed orthodontic. Journal of Syiah Kuala Dentistry Society. 2016; 1(1) :79 – 80
- [3] Mararu, Wahyu P, et al. Dental hygiene status of patient using fixed orthodontic in Senior High School 7 Manado. Journal e-GiGi (eG). 2017; 5(2): 159-165
- [4] Amalia M. Knowledge of patient with side effect orthodontic treatment in Dental Hospital of Dentistry Faculty of Indonesia University. Jakarta, Indonesia University, Thesis. 2009
- [5] Graber T, Eliades T, Athanasiou AE. Risk Management in Orthodontics: Experts Guide to Malpractice. Quintessence Publishing Co, Inc, Chicago. 2004
- [6] Jayanti, Triana Amaliah, et al. The Relationship between duration of fixed orthodontic treatment with periodontal status and treatment needs among students in The Faculty Of Dentistry Universitas Muslim Indonesia Makassar In 2017. Dentino (Jur. Ked. Gigi). 2018; 3(1): 85 – 90
- [7] Dannan A. An update on periodontic-orthodontic interrelationships. J Indian Soc Periodontol. 2010;14(1):66-71.
- [8] Klukowska M, Bader A, Erbe C, Bellamy P, White DJ, Anastasia MK, Wehrbein H. Plaque levels of patients with fixed orthodontic appliances measured by digital plaque image analysis. Am J Orthod Dentofacial Orthop. 2011;139(5):463-70.
- [9] Thilander B, Rygh P, Reitan K. Tissue reactions in orthodontics. in Orthodontics: current principles and techniques, T. Graber, et al. eds., 2005. St Louis. Mosby;145-219.
- [10] Un-Bong Baik, et al. Teeth discoloration during orthodontic treatment. The Korean Journal of Orthodontics. 2017; 47(5): 334-339
- [11] Jufri, Yuli Prihastuti. Relation of orthodontic fixed with discoloration teeth. Dentistry Faculty of Hasanuddin University Makassar. Thesis. 2017
- [12] Fjeld M, Øgaard B. Scanning electron microscopic evaluation of enamel surfaces exposed to 3 orthodontic bonding systems. Am J Orthod Dentofacial Orthop. 2006;130(5):575-81.
- [13] Øgaard B, Fjeld M. The enamel surface and bonding in orthodontics. seminars in orthodontics. 2010;16(1):37-48.
- [14] Santamaria M Jr, Milagres D, Stuani AS, Stuani MB, Ruellas AC. Initial changes in pulpal microvasculature during orthodontic tooth movement: a stereological study. Eur J Orthod 2006; 28(3):217-20.
- [15] Faltermeier, A., Rosentritt, M., Reicheneder, C. & Behr, M. Discolouration of orthodontic adhesives caused by food dyes and ultraviolet light. European Journal of Orthodontics. 2008; 30(1): 89-93
- [16] Lestari, Nurasisa, et al. Relation time using fixed orthodontic with plaque accumulation and saliva's pH in student of dentistry faculty of Moslem University of Indonesia in 2017. As-Syifaa. 2018; 10(01): 126-133
- [17] Al Maaitah EF, Adeyemi AA, Higham SM, Pender N, Harrison JE. Factors affecting demineralization during orthodontic treatment: a post-hoc analysis of RCT recruits. Am J Orthod Dentofacial Orthop. 2011;139(2):181-91
- [18] Danesh G, Hellak A, Lippold C, Ziebur T, Schafer E. Enamel surfaces following interproximal reduction with different methods. Angle Orthod. 2007;77(6):1004-10.
- [19] Yamaguchi M, Kasai K. The effects of orthodontic mechanics on the dental pulp. Seminars Orthod. 2007;13:272–80.
- [20] Ramazanzadeh BA, Sahhafian AA, Mohtasham N, Hassanzadeh N, Jahanbin A, Shakeri MT. Histological changes in human dental pulp following application of intrusive and extrusive orthodontic forces. J Oral Sci. 2009;51(1):109-15.
- [21] Guttenberg, S.A. Cosmesis of the mouth, face and jaw. Willey-Blackwell, USA. 2012: 39
- [22] Yovela. Krinawati. Management of protrusion anterior teeth maxilla anterior with abnormal periodontal pada in <http://annalsofrscb.ro>

- adult patient. *Indonesian Journal of Dentistry*. 2009; 16(1): 25,30.
- [23] Pacheco-Pereira C, Pereira JR, Dick BD, Perez A, Flores-Mir C. Factors associated with patient and parent satisfaction after orthodontic treatment: a systematic review. *Am J Orthod Dentofacial Orthop*. 2015;148(4):652-9.
- [24] Tsichlaki A, Chin SY, Pandis N, Fleming PS. How long does treatment with fixed orthodontic appliances last? A systematic review. *Am J Orthod Dentofacial Orthop*. 2016;149(3):308-18.
- [25] Aljehani D, Baeshen HA. Effectiveness of the american board of orthodontics discrepancy index in predicting treatment time. *J Contemp Dent Pract*. 2018;19(6):647-50.
- [26] Papageorgiou SN, Hochli D, Eliades T. Outcomes of comprehensive fixed appliance orthodontic treatment: a systematic review with meta-analysis and methodological overview. *Korean J Orthod*. 2017;47(6):401-13
- [27] Uribe F, Padala S, Allareddy V, Nanda R. Patients', parents', and orthodontists' perceptions of the need for and costs of additional procedures to reduce treatment time. *Am J Orthod Dentofacial Orthop*. 2014;145:65-73.
- [28] C6mo, & Zapata-Nore6a, Oscar & Barbosa-Liz, D & Carvajal, Alvaro & Plaza, Sonia. Factors related to Orthodontic Treatment Duration for a Graduate Orthodontics Program. 2017. 17-25.
- [29] Pinto AS, Alves LS, Maltz M, Susin C, Zenkner JEA. Does the duration of fixed orthodontic treatment affect caries activity among adolescents and young adults? *Caries Res*. 2018;52(6):463-7
- [30] Puspitasari, Yustisia, et al. The relation between duration of fixed orthodontic treatment and oral hygiene status among students of Faculty of Dentistry Universitas Muslim Indonesia 2017. *Dentino (Jur. Ked. Gigi)*. 2018; 3(2): 162 – 167
- [31] Preoteasa, Cristina Teodora, et al. Risk and complication associated with orthodontic treatment. University of Medicine and Pharmacy, Bucharest, Romania. 2012
- [32] Huang GJ, et al. Effectiveness of MI paste plus and Prev-Dent fluoride varnish for treatment of white spot lesions: A randomized controlled trial. *Am J Orthod Dentofacial Orthop*. 2013; 143: 31-41.
- [33] Arab S, et al. Effect of fixed orthodontic treatment on saliva flow, pH and, microbial count. 2016; 3(1): 18-19.
- [34] Carillo LE, et al. Effect of Orthodontic treatment on saliva plaque, and the levels of *Streptococcus Mutans* and *Lactobacillus*. 2010; 15(6): 924-929.
- [35] Sandison RM. Tooth surface appearance after debonding. *Br J Orthod*. 1998; 8: 199–201.
- [36] Chen, Qiushuo et al. Influence of orthodontic treatment with fixed appliances on enamel color: A systematic review. *BMC Oral Health*. 2015; 15(31)
- [37] Ramadhan A, et al. Relation level of knowledge dental health to number of dental caries in Junior High School 1 Marabahan. *Dentino Journal*. 2016; 1(2).
- [38] Mantiri SC, et al. Status of hygiene and caries status of student who using fixed orthodontic. *Dental Journal*. 2013; 1(1).
- [39] Achmad H, Djais AJ, Petrenko EG, Larisa V, Putra AP. 3-d printing as a tool for applying biotechnologies in modern medicine. *International Journal of Pharmaceutical Research*, 2020. 12(4), pp. 3454-3463.
- [40] Achmad H, Djais AI, Jannah M, Huldani, Putra AP. Antibacterial chitosan of milkfish scales (*Chanos chanos*) on bacteria *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans*. *Systematic Review in Pharmacy*, 2020. 11(6), pp. 836-841.
- [41] Achmad H, Djais AI, Syahrir S, Fitri A, Ramadhany YF. A literature us regarding the use of herbal medicines in pediatric dentistry. *International Journal of Pharmaceutical Research*. 2020. 12, PP. 881-897.
- [42] Achmad H, Djais AI, Syahrir S, Fitri A, Ramadhany YF. Impact Covid-19 in pediatric dentistry: A literature review. *International Journal of Pharmaceutical Research*, 2020. 12, p. 830-840.
- [43] Djais AI, Achmad H, Dewiayu D, Sukmana BI, Huldani. Effect of Combination of Demineralization Freeze Dentin Matrix (DFDDM/0 and *Moringa oleifera* lam osteoprotegerin (OPG) and receptor activator of nuclear factor kappa B ligand (RANKL) as a marker of bone remodeling. *Systematic Reviews in Pharmacy*. 2020. 11(6), pp. 771-779.

