

DAFTAR PUSTAKA

- Ahsanuddin. 2021. *Adverse Events Associated with Intranasal Sprays: An Analysis of the Food and Drug Administration Database and Literature Review. Annals of Otology, Rhinology & Laryngology*, 000348942110072. doi:10.1177/00034894211007222
- Anatomi dan fisiologi lidah (internet). Available from. [http:// digilib .uinsby.ac.id/15869/10/Bab%207.pdf](http://digilib.uinsby.ac.id/15869/10/Bab%207.pdf)
- Altin F, Cingi C, Uzun T, Bal C.2020. Olfactory and gustatory abnormalities in COVID-19 cases. *Eur Arch Otorhinolaryngol*; 277:2775–81
- Baek Y.J., Lee Y.J., Yoon J.S., et al.2021. Duration of culturable SARS-CoV-2 within different specimens among mild and severe COVID-19 patients: a longitudinal study. *J Infect* doi: <https://doi.org/doi:10.1016/j.jinf.2021.04.025>.
- Baig AM.2020. Neurological manifestations in COVID19 caused by SARSCoV 2. *CNS neuroscience & therapeutics*. ;26(5):499.
- Bastier, P. et al. 2015. Nasal irrigation : From empiricism to evidence-based medicine . A review', *European Annals of Otorhinolaryngology, Head and Neck diseases*. Elsevier Masson SAS, 132(5), pp. 281–285. doi: 10.1016/j.anorl.2015.08.001.
- Bastier, lechot, bordenave, et al. 2015. Nasal Irrigation: from empiricism to evidence-base medicine a review. *Eur Ann Otorhinolaryngol Head Neck Dis* Nov;132(5):281-5

- Butowt R, von Bartheld CS, 2020. Anosmia in COVID-19: underlying mechanisms and assessment of an olfactory route to brain infection. *Neuroscientist* 2020 Sep 11 DOI: 10.1177/1073858420956905
- Boscolo-Rizzo P, Borsetto D, Fabbris C, Spinato G, Frezza D, Menegaldo A, et al. Evolution of altered sense of smell or taste in patients with mildly symptomatic COVID-19. *JAMA Otolaryngology-Head & Neck Surgery* 2020;146(8):729-32.
- Boyce J M. 2006. Effects of ageing on smell and taste. *Postgrad Med J* ;82:239–241. doi: 10.1136/pgmj.2005.039453
- Cartabuke R, Andherson B, Elmaraghy at al. 2019. Hemodynamic and pharmacokinetic analysis of oxmetazoline use during nasal surgery in children . *Laryngoscope*, 129:2775–2781. <https://doi.org/10.1002/lary.27760>
- Carolin A. et al (2021) Analysis of *Cycle Threshold* values in SARS-CoV-2-PCR in a long-term study. *Journal of Clinical Virology* 138 104791. <https://doi.org/10.1016/j.jcv.2021.104791>
- Chary E, Carsuzaa F, Trijolet J-P, et al. 2020. Prevalence and recovery from olfactory and gustatory dysfunctions in COVID-19 infection: a prospective multicenter study. *American Journal of Rhinology & Allergy*;34(5):686-93. DOI: 10.1177/1945892420930954

- Cooper KW, Brann DH, Farruggia MC, et al. 2020.COVID-19 and the chemical senses: Supporting players take center stage. *Neuron*; 107(2):219–33.
- COVID-19.go.id (internet), 2021. Data sebaran COVID-19. Available from : <http://Covid19.go.id>
- Ganong WF. 2001. Smell and taste. In *Review of medical physiology*. 20th ed. San Fransisco: Medical Publishing Division;. p. 340-7.
- Gerkin RC, et al. 2020. The best COVID-19 predictor is recent smell loss: a cross-sectional study. [https://www.medrxiv.org/ content/ 10.1101/ 2020.07.22.20157263v2](https://www.medrxiv.org/content/10.1101/2020.07.22.20157263v2). DOI: 10.1101/2020.07.22.20157263
- Guzainuer A.et al, 2023. A follow up study of *Cycle Threshold* values of SARS-CoV-2 in Hunan Province, China. *Infectious Disease Modelling* 8 203-211. <https://doi.org/10.1016/j.idm.2023.01.004> 2468-0427/
- Guan W, Ni Z, Hu Y, Liang W et al. 2020. Clinical Characteristics Of Coronavirus Disease 2019 in china. *N Engl J Med* ;38;1708-20. Doi: 10.1056/NEJMoa200032
- Heidari F, Karimi E, Firouzifar M, et al. 2020. Anosmia as a prominent symptom of COVID-19 infection. *Rhinology* ;58(3):302-3.
- Higgnis T, Athur, et al. 2020. Intranasal Antiviral Drug Delivery and Coronavirus Disease 2019 (COVID-19): A State of the Art Review. *Otolaryngology– Head and Neck Surgery*, Vol. 163(4) 682–694

- Hopkins C, Lechien JR, Saussez S. 2020. More than ACE2? NRP1 may play a central role in the underlying pathophysiological mechanism of olfactory dysfunction in COVID-19 and its association with enhanced survival. *Medical Hypotheses*. Nov 20 . DOI: 10.1016/j.mehy.2020.110406
- Huriyati E, Nelvia T. Et al. 2014. Gangguan Fungsi Penghidu dan Pemeriksaannya. [Http: //jurnal.fk.unand.ac.id](http://jurnal.fk.unand.ac.id).
- Lechien JR, Chiesa-Estomba CM, De Siati DR, et al. 2020. Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *European Archives of Oto-rhinolaryngology* ;277(8):2251-61. DOI: 10.1007/s00405-020-05965-1
- Luigi A. Vaira, MD ; Giovanni Salzano, et al. 2020. Anosmia and Ageusia: Common Findings in COVID-19 Patients. *Laryngoscope*, 00:1. DOI: 10.1002/lary.28692
- Mahraeen E, Karimi A, Barzegary A. 2020. Predictors of mortality in patients with COVID-19—a systematic review *Eur J Integr Med* Dec;40:101226. [http: //doi:10.1016/j.eujim.101226](http://doi:10.1016/j.eujim.101226)
- Matthew M, Nadine C, et al. 2021. Structural Basis of SARS Cov 2 Omicron Immune Evasion and Receptor Engagement. *Science*
- Martina et al. 2023. Nasal cast deposition for xylo- and oxymetazoline formulations using two different nasal pumps *Rhinology Online*, Vol 6: 11 - 17, [http://doi.org /10.4193/RHINOL/23.002](http://doi.org/10.4193/RHINOL/23.002)

- Meng X, Deng Y, Dai Z, Meng Z. 2020. COVID-19 and anosmia: A review based on up-to-date knowledge. *Am J Otolaryngol.* ;41(5):102581
- Mueller CA, Grasinger E, Naka A, 2006. A self administered odor identification test procedure using the "sniffin sticks". *Chem Senses* ; 31: 595-98.
- Meunier N, Briand L, Jacquin A, et al. 2020. COVID-19-Induced Smell and Taste Impairments: Putative Impact on Physiology. *Front.physiol.* [https://doi.org/ 10.3389/fphys.625110](https://doi.org/10.3389/fphys.625110)
- Neighbors, et al. 2023. Intranasal corticosteroid and oxymetazoline for chronic rhinitis: a systematic review. *J Laryngol Otol* 2022;136:8–16. <https://doi.org/10.1017/S0022215121003364>
- O'Byrne L, et al. 2022. Interventions for the treatment of persistent post-COVID-19 olfactory dysfunction. *Cochrane Database of Systematic Reviews* , Issue 9. Art. No.: CD013876. DOI: 10.1002/14651858.CD013876.pub3
- Okosua. et al .2020. Smell and Taste Dysfunction in Patients With COVID-19: A Systematic Review and Meta-analysis. *Elsevie.* 2020;95(8):1621-1631
<https://doi.org/10.1016/j.mayocp.2020.05.030>
- Onghyun, et al. 2020. .Prevalence and Duration of Acute Loss of Smell or Taste in COVID-19 Patients. *J Korean Med Sci.*11;35(18):e174
<https://doi.org/10.3346/jkms.2020.35.e174>

- Retno SM, Mir'ah Z, et al. 2016. Penggunaan terkini oksimetazolin pada praktik klinik. ORLI Vol. 46 No. 2.
- Rothan H A, 2020. Byrareddy S N. The Epidemiology And Pathogenesis Of Coronavirus Disease (C0vid 19) Outbreak. Journal Of Autoimmunity, <http://doi.org/10.1016/j.jaut.2020.102433>
- Rouby C, Danguin TT, Vigouroux M, et all. 2011. The Lyon clinical olfactory test: Validation and measurement of hiposmia and anosmia in healthy and diseased population. International Journal of otolaryngology; 23:1-9
- Rachana, R., Santhi, T. 2019. Efficacy of Saline Nasal Irrigation in Chronic Rhinosinusitis 7, 71–77
- Rajamani S. et al. 2022. A cross-sectional study to assess the prevalence of anosmia and ageusia and its association with disease severity among COVID-19 affected patients in Salem, Tamil Nadu. J Family Med Prim Care; 11:2340-4.ebsite: www.jfmprc.com DOI: 10.4103/jfmprc.jfmprc_1308_21
- Richard C,et al. 2021. Topical Nasal Decongestant Oxymetazoline: Safety Considerations for Perioperative Pediatric Use. From The American Academy Of Pediatrics. *Pediatrics* 148 (5): e2021054271. <https://doi.org/10.1542/peds.2021-054271>
- Salim S, Qurraisha A. 2021. Omicron SARS-CoV-2 variant: a new chapter in the COVID-19 pandemic . www.thelancet.com Vol 398, [https://doi.org/10.1016/S0140-6736\(21\)02758-6](https://doi.org/10.1016/S0140-6736(21)02758-6)

- Stephen W, Sarah J, Kashif J, et al. 2021. Clinical efficacy of nitric oxide *nasal spray* (NONS) for the treatment of mild COVID-19 infection. Letters to the Editor/ Journal of Infection 83 237–279.
- Saussez S, Lechien JR, Hopkins C. 2020. Anosmia: an evolution of our understanding of its importance in COVID-19 and what questions remain to be answered. European Archives of Oto rhino-laryngology. Sep 9. DOI: 10.1007/s00405-020-06285-0
- Thongngarm T, Assanasen P, Pradubpongsa P, Tantilipikorn P. The effectiveness of oxymetazoline plus intranasal steroid in the treatment of chronic rhinitis: a randomised controlled trial. Asian Pac J Allergy Immunol. 2016;34(1):30-37.
- Vaira, 2020. Objective evaluation of anosmia and ageusia in COVID-19 patients: Single-center experience on 72 cases
- Vallecillo MVS, 2012. Fraire ME, Cagnani CB, Zernotti ME. Olfactory dysfunction in patient with cronic rhinosinusitis. International journal of otolaryngology; Article ID 32206: 1-5.
- Von Bartheld, Butowt R, 2020. Anosmia in COVID-19: Underlying Mechanisms and Assessment of an Olfactory Route to Brain Infection. Sagepub.com/journal .
<http://Doi:10.1177/1073858420956905>
- Winther B, buchert D, et al. 2010. Decreased rhinovirus shedding after intranasal oxymetazoline application in adults with induced colds

compared with intranasal saline. American Journal of Rhinology & Allergy. Vol. 24, No. 5

Yerikala R, Abhirami B, Sri G, et al. 2018. Formulation and evaluation of oxymetazoline hydrochloride nasal gels. Journal of Drug Delivery & Therapeutics; 8(6):49-57

Yan H et al. 2020. alSelf-reported olfactory loss associates with outpatient clinical course in COVID-19 International Forum of Allergy & Rhinology, Vol. 10, No. 7,doi.org/10.1002/alr.22592

Zubair AS, McAlpine LS, Gardin T, et al. 2020. Neuropathogenesis and neurologic manifestations of the coronaviruses in the age of coronavirus disease 2019: A review. JAMA Neurol;77(8):1018-24

Zachary M. 2020. A primer on viral-associated olfactory loss in the era of COVID-19. Internat Forum Allerg Rhinol ;10(7):814-20