

DAFTAR PUSTAKA

1. Kirkaldy-Willis WH, Farfan HF. Instability of the lumbar spine. Clin Orthop Relat Res. 1982;165.
2. Biely S, Smith SS, Silfies SP. Clinical Instability of the Lumbar Spine : Diagnosis and Intervention. Analysis. 2006;18(3).
3. Sunderland G, Foster M, Dheerendra S, Pillay R. Patient-Reported Outcomes Following Lumbar Decompression Surgery: A Review of 2699 Cases. Glob Spine J. 2021;11(2).
4. Regev GJ, Leor G, Ankori R, Hochberg U, Ofir D, Khashan M, et al. Long-term pain characteristics and management following minimally invasive spinal decompression and open laminectomy and fusion for spinal stenosis. Med. 2021;57(10).
5. Chou R, Gordon DB, De Leon-Casasola OA, Rosenberg JM, Bickler S, Brennan T, et al. Management of postoperative pain: A clinical practice guideline from the American pain society, the American society of regional anesthesia and pain medicine, and the American society of anesthesiologists' committee on regional anesthesia, executive committee, and administrative council. J Pain. 2016;17(2).
6. St-Jacques B, Ma W. Role of prostaglandin E2 in the synthesis of the pro-inflammatory cytokine interleukin-6 in primary sensory neurons: an in vivo and in vitro study. J Neurochem. 2011 Sep;118(5):841-54.

7. Sudiarta IPG, Wiargitha IK, Mahadewa TGB. Perbedaan nilai Neutrophil Lymphocyte Ratio (NLR) terhadap pemeriksaan kultur darah dalam mendiagnosis sepsis pada pasien peritonitis di RSUP Sanglah, Bali, Indonesia. *Intisari Sains Medis*. 2020;11(1):165.
8. Paliogiannis P, Deidda S, Maslyankov S, Paycheva T, Farag A, Mashhour A, et al. Blood cell count indexes as predictors of anastomotic leakage in elective colorectal surgery: A multicenter study on 1432 patients. *World J Surg Oncol*. 2020;18(1):1–8.
9. Hajibandeh S, Hajibandeh S, Hobbs N, Mansour M. Neutrophil-to-lymphocyte ratio predicts acute appendicitis and distinguishes between complicated and uncomplicated appendicitis: A systematic review and meta-analysis. *Am J Surg [Internet]*. 2020;219(1):154–63.
10. Kuchálik J, Magnuson A, Tina E, Gupta A. Does local infiltration analgesia reduce perioperative inflammation following total hip arthroplasty? A randomized, double-blind study. *BMC Anesthesiol*. 2017;17(1):1–9.
11. Asghar A, Saqib U, Aijaz S, Hussain K, Bukhari S, Hayat A. Utility of Neutrophil-to-Lymphocyte Ratio, Platelets-to-Lymphocyte Ratio and Call Scire for Prognosis Assessment in COVID-19 Patients. 2020;19(2):590–6.
12. Jimeno S, Ventura PS, Castellano JM, García-Adasme SI, Miranda M, Touza P, et al. Prognostic implications of neutrophil-lymphocyte ratio in COVID-19. *Eur J Clin Invest*. 2021;51(1):1–9.
13. Djordjevic D, Rondovic G, Surbatovic M, Stanojevic I, Udovicic I, Andjelic

- T, et al. Neutrophil-to-Lymphocyte Ratio, Monocyte-to-Lymphocyte Ratio, Platelet-to-Lymphocyte Ratio, and Mean Platelet Volume-to-Platelet Count Ratio as Biomarkers in Critically Ill and Injured Patients: Which Ratio to Choose to Predict Outcome and Nature of Bacte. *Mediators Inflamm.* 2018;2018.
14. Perkins NM, Tracey DJ. Hyperalgesia due to nerve injury: Role of neutrophils. *Neuroscience.* 2000;101(3):745–57.
 15. Butterfield T a, Best TM, Merrick M a. The Dual Roles of Neutrophils and. *J Athl Train.* 2006;41(4):457–65.
 16. Cassuto J, Sinclair R, Bonderovic M. Anti-inflammatory properties of local anesthetics and their present and potential clinical implications. *Acta Anaesthesiol Scand.* 2006;50(3):265–82.
 17. Wang L, Liang S, Chen H, Xu Y, Wang Y. The effects of epidural anaesthesia and analgesia on T lymphocytes differentiation markers and cytokines in patients after gastric cancer resection. *BMC Anesthesiol.* 2019;19(1):1–7.
 18. Kourilovitch M, Galarza–Maldonado C. Could a simple biomarker as neutrophil-to-lymphocyte ratio reflect complex processes orchestrated by neutrophils? *J Transl Autoimmun* [Internet]. 2022;(xxxx):100159.
 19. Daoudia M., Decruynaere C., Le Polain de Waroux B., Thonnard J.L., Plaghki L., Forget P. Biological inflammatory markers mediate the effect of preoperative pain-related behaviours on postoperative analgesics

- requirements. *BMC Anesthesiol.* 2015;15:183.
20. Turgut H.C., Alkan M., Atac M.S. Neutrophil lymphocyte ratio predicts postoperative pain after orthognathic surgery. *Niger J Clin Pract.* 2017;20(10):1242–1245.
 21. Shell WE, Pavlik S, Roth B, Silver M, Breitstein ML, May L, Silver D. Reduction in Pain and Inflammation Associated With Chronic Low Back Pain With the Use of the Medical Food Theramine. *Am J Ther.* 2016 Nov/Dec;23(6):e1353-e1362.
 22. de Queiroz BZ, Pereira DS, Lopes RA, Felício DC, Silva JP, Rosa NM, Dias JM, Dias RC, Lustosa LP, Pereira LS. Association Between the Plasma Levels of Mediators of Inflammation With Pain and Disability in the Elderly With Acute Low Back Pain: Data From the Back Complaints in the Elders (BACE)-Brazil Study. *Spine (Phila Pa 1976).* 2016 Feb;41(3):197-203.
 23. Licciardone JC, Kearns CM, Hodge LM, Bergamini MV. Associations of cytokine concentrations with key osteopathic lesions and clinical outcomes in patients with nonspecific chronic low back pain: results from the OSTEOPATHIC Trial. *J Am Osteopath Assoc.* 2012 Sep;112(9):596-605.
 24. Sebba A. Pain: A review of interleukin-6 and its roles in the pain of rheumatoid arthritis. *Open Access Rheumatol Res Rev.* 2021;13:31–43
 25. Zhou YQ, Liu Z, Liu ZH, Chen SP, Li M, Shahveranov A, Ye DW, Tian YK. Interleukin-6: an emerging regulator of pathological pain. *J Neuroinflammation.* 2016 Jun 7;13(1):141.

26. Hindman BJ, Palecek JP, Posner KL, Traynelis VC, Lee LA, Sawin PD. Cervical spinal cord, root, and bony spine injuries: A closed claims analysis. *Anesthesiology* 2011;114:782-95.
27. Paul L, Adrian M.N. 2005. Approach and Considerations Regarding the Patient with Spinal Injury. www.elsevier.com/locate/injury. Australia. Pp : 2-11. 15th Oct 2017
28. Prolo LM, oklund SA, Zawadzki N, Desai M, Polo DJ. Uninstrumented posterior lumbar interbody fusion: have technological advances in stabilizing the lumbar spine truly improved outcomes? *World neurosurg.* 2018;115:503-4.
29. Macres SM, Moore PG, Fishman SM. Acute pain management. In: Barash PG, Cullen BF, Stoelting RK, Cahalan MK, Stock MC, Orthega R, Sharar SR, Holt NF. *Clinical anesthesia*, edisi8. 2017. Wolters Kluwer : Philadelphia. p. 3919-32.
30. Lavand'homme P. From preemptive to preventive analgesia: time to reconsider the role of perioperative peripheral nerve block? *Reg Anesth Pain med.* 2011;36(1)4-6.
31. Wahdah SW, Radwan TA, Mohammed MM, Mohamed AA, Salama AK. Effect of bilateral ultrasound-guided erector spinae blocks on postoperative pain and opioid use after lumbar spine surgery: A prospective randomized controlled trial. *Egypt J Anaesth.* 2020;37(1):100-6.
32. Altıparmak B, Tker MK, Uysal AI, Demirbilek SG. Comparison of the

efficacy of erector spinae plane block performed with different concentrations of bupivacaine on postoperative analgesia after mastectomy surgery: randomized, prospective, double blinded trial. *BMC Anesthesiol.* 2019;19(1):31.

33. Forero M, Adhikary SD, Lopez H, Tsui C, Chin KJ, . The Erector Spinae Plane Block A Novel Analgesic Technique in Thoracic Neuropathic Pain. *Reg Anesth Pain Med.* 2016;41:621-7.
34. Chin KJ, Adkhikary SD, Forero M. Erector spinae plane (ESP) block: a new paradigm in regional anesthesia and analgesia. *Curr Anesthesiol Rep.* 2019;9:271-80.
35. Apibunyopas Y, Ho B,. Ultrasound-guided erector spinae plane (ESP) block: A novel intervention for mechanical back pain in the emergency department. *CJEM.* 2019;21(2):302-5.
36. Butterworth JF, Mackey DC, Wasnick JD. Local anesthetic. in: Butterworth JF, Mackey DC, Wasnick JD, eds. *Morgan & Mikhail's Clinical anesthesiology*, edisi 6. McGraw Hill : New York. p 452-471.
37. Waelkens P, Alsabbagh E, Sauter A, Joshi GP, Beloeil H; PROSPECT Working group** of the European Society of Regional Anaesthesia and Pain therapy (ESRA). Pain management after complex spine surgery: A systematic review and procedure-specific postoperative pain management recommendations. *Eur J Anaesthesiol.* 2021 Sep 1;38(9):985-994.
38. Cumming KI, Naguib MA. Opioid agonist and antagonist. In: Flood P,

- Rathmell JP, Shafer S, eds. Pharmacology and physiology in anesthetic practice. edisi 5. Wolters Kluwer : Philadelphia. p217-50.
39. Daniel M, Weiskopf RB, Noorani M. Fentanyl augments the blockade of the sympathetic response to incision (MAC-BAR) produced by desflurane and isoflurane : desflurane and isoflurane MAC-BAR without and with fentanyl. *Anesthesiology*. 1998;88:43–49.
 40. Rosales C. Neutrophil: A cell with many roles in inflammation or several cell types? *Front Physiol*. 2018;9(FEB):1–17.
 41. Yang AP, Liu JP, Tao WQ, Li HM. The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients. *Int Immunopharmacol*. 2020;84(April):106504.
 42. Hwang SY, Shin TG, Jo IJ, Jeon K, Suh GY, Lee TR, et al. Neutrophil-to-lymphocyte ratio as a prognostic marker in critically-ill septic patients. *Am J Emerg Med* 2017 Feb;35(2):234-239.
 43. Hamza M, Dionne RA. Mechanisms of non-opioid analgesics beyond cyclooxygenase enzyme inhibition. *Curr Mol Pharmacol*. 2009 Jan;2(1):1-14.
 44. Canbolat N, Buget MI, Sivriköz N, Altun D, Kucukay S. Relação entre a proporção neutrófilo/linfócito e a dor pós-operatória em artroplastia total de joelho e quadril [The relationship between neutrophil to lymphocyte ratio and postoperative pain in total knee and hip arthroplasty]. *Braz J Anesthesiol*. 2019 Jan-Feb;69(1):42-47.

45. Öner K, Okutan AE, Ayas MS, Paksoy AE, Polat F. Predicting postoperative pain with neutrophil/ lymphocyte ratio after arthroscopic rotator cuff repair. Asia Pac J Sports Med Arthrosc Rehabil Technol. 2020 Mar 17;20:24-27
46. Akilli NB, Yortanlı M, Mutlu H, et al. Prognostic importance of neutrophil-lymphocyte ratio in critically ill patients: short- and long-term outcomes. Am J Emerg Med. 2014;32: 1476-80.
47. KÖLÜKÇÜ V, KAHVECİ M. Predictive Role of Neutrophil/Lymphocyte Ratio for Postoperative Pain. Gaziosmanpaşa Tıp Dergisi. 2022;14(2):110-