

Daftar Pustaka

- Adhikari, K., Bhandari, S., Aryal, K., Mahato, M., & Shrestha, J. (2021). Effect of different levels of nitrogen on growth and yield of hybrid maize (*Zea mays* L.) varieties. *Journal of Agriculture and Natural Resources*, 4(2), 48–62. <https://doi.org/10.3126/janr.v4i2.33656>
- Amanullah, Iqbal, A., Ali, A., Fahad, S., & Parmar, B. (2016). Nitrogen source and rate management improve maize productivity of smallholders under semiarid climates. *Frontiers in Plant Science*, 7(NOVEMBER2016), 1–9. <https://doi.org/10.3389/fpls.2016.01773>
- Agunbiade, V. F., & Babalola, O. O. (2024). Drought stress amelioration attributes of plant-associated microbiome on agricultural plants. *Bioinformatics and Biology Insights*, 18, 11779322241233442. <https://doi.org/10.1177/11779322241233442>
- Agyare, W. A., B. S. Freduah, E. Ofori, D. S. Kpongor, and B. O. A. (2013). Field and modelled maize (*Zea Mays*) response to water stress at different growth stages. *Glob. J. Biol.* 2, 68–75.
- Ahmad, N., Malagoli, M., Wirtz, M. (2016). Drought stress in maize causes differential acclimation responses of glutathione and sulfur metabolism in leaves and roots. *BMC Plant Biol.* 16, 247. Available at: <https://doi.org/10.1186/s12870-016-0940-z>.
- Arzu, K., Onder, O., Bilir, O., and Kosar, F. (2018). Application of multivariate statistical analysis for breeding strategies of spring safflower (*Carthamus tinctorius* L.). *Turkish J. F. Crop.* 23, 12–19.
- Assefa, Y., Prasad, P. V. V., Carter, P., Hinds, M., Bhalla, G., Schon, R., ... & Ciampitti, I. A. (2018). A new insight into corn yield: Trends from 1987 through 2015. *Agronomy Journal*, 110(4), 1793-1804.
- Astuti K., O. R. Prasetyo, I N. Khasanah, 2021. Analisis Produktivitas Jagung dan Kedelai di Indonesia 2020 (Hasil Survei Ubinan), Badan Pusat Statistik/BPS-Statistics Indonesia
- Azrai, M., Aqil, M. Andayani, NN..Efendi, R., Suarni, Suwardi, Jihad, M., Zainuddin ,B., Salim, Bahtiar, Muliadi, A., Yasin ,M.,

- Hannan ,MFI., Rahman, Syam, A. (2024). Optimizing ensembles machine learning, genetic algorithms, and multivariate modeling for enhanced prediction of maize yield and stress tolerance index. *Front. Sustain. Food Syst* 8. Available at: <https://doi.org/10.3389/fsufs.2024.1334421>.
- Azrai, M., Aqil, M., Efendi, R., Andayani, N. N., Makkulawu, A. T., Iriany, R. N., et al. (2023). A comparative study on single and multiple trait selections of equatorial grown maize hybrids. *Front. Sustain. Food Syst* 7. doi: 10.3389/fsufs.2023.1185102.
- Balbua, M.G., H.T. Osman, E.E. Kandil, T. Javed, S. F. L. (2022). Determination of morpho-physiological and yield traits of maize inbred lines (*Zea mays* L.) under optimal and drought stress conditions. *Front. Plant Sci.* 13, 1–17. doi: 10.3389/fpls.2022.959203.
- Bandyopadhyay N., C. Bhuiyan, A. K. S. (2020). Drought mitigation: Critical analysis and proposal for a new drought policy with special reference to Gujarat. *Prog. Disaster Sci.* 5. Available at: <https://doi.org/10.1016/j.pdisas.2019.100049>.
- Bänziger, M., Edmeades, G. O., Beck, D., and Bellon, M. (2000). Breeding for Drought and Nitrogen Stress Tolerance in Maize: From Theory to Practice. *Mex. D.F. CIMMYT*, 68.
- Bänziger, M., Edmeades, G. O., & Lafitte, H. R. (1999). Selection for drought tolerance increases maize yields over a range of nitrogen levels. *Crop Science*, 39(4), 1035-1040.
- Black, C. A. (1976). *Soil-Plant Relationships*. John Wiley & Sons, New York.
- Burgess, A. J., Retkute, R., Herman, T., & Murchie, E. H. (2017). Exploring relationships between canopy architecture, light distribution, and photosynthesis in contrasting rice genotypes using 3D canopy reconstruction. *Frontiers in Plant Science*, 8(May), 1–15. <https://doi.org/10.3389/fpls.2017.00734>
- Bolaños, J., and Edmeades, G. O. (1996). The importance of the anthesis-silking interval in breeding for drought tolerance in tropical maize. In: *Developing Drought and Low N-Tolerant Maize. (Eds.). *Proc. a Symp. March 25-29, CIMMYT, EL Batán, Mex. D.F., Mex.*

- Cairns, J. E., Hellin, J., Sonder, K., Araus, J. L., MacRobert, J. F., Thierfelder, C., & Prasanna, B. M. (2012). Adapting maize production to climate change in sub-Saharan Africa. *Food Security*, 4, 399-417.
- Chauhan J, Srivastava JP, Singhal RK, Soufan W, Dadarwal BK, Mishra UN, Anuragi H, Rahman MA, Sakran MI, Brestic M, Zivcak M, S. M. and S. A. (2022). Alterations of Oxidative Stress Indicators, Antioxidant Enzymes, Soluble Sugars, and Amino Acids in Mustard [*Brassica juncea* (L.) Czern and Coss.] in Response to Varying Sowing Time, and Field Temperature. *Front. Plant Sci.* 13, 875009. doi: 10.3389/fpls.2022.875009.
- Chukwudi, U.P., F.R. Kutu, and S. M. (2021). Heat stress effect on the grain yield of three drought-tolerant maize varieties under varying growth conditions. *Plants* 10, 1–15. doi: 10.3390/plants10081532.
- Deribe, H. (2024). Review on Effects of Drought Stress on Maize Growth, Yield and Its Management Strategies. *Commun. Soil Sci. Plant Anal.*, 1–21. Available at: <https://doi.org/10.1080/00103624.2024.2404663>.
- De Araujo Rufino, C., Fernandes-Vieira, J., Martín-Gil, J., Abreu Júnior, J. D. S., Tavares, L. C., Fernandes-Correa, M., & Martín-Ramos, P. (2018). Water stress influence on the vegetative period yield components of different maize genotypes. **Agronomy*, 8*(8), 151. <https://doi.org/10.3390/agronomy8080151>
- Duvick, D. N. (2005). The Contribution of Breeding to Yield Advances in Maize (*Zea mays* L.). *Advances in Agronomy*, 86, 83-145.
- Dodig, D., Božinović, S., Nikolić, A., Zorić, M., Vančetović, J., Ignjatović-Micić, D., Delić, N., Weigelt-Fischer, K., Altmann, T., & Junker, A. (2021). Dynamics of maize vegetative growth and drought adaptability using image-based phenotyping under controlled conditions. *Frontiers in Plant Science*, 12, 652116. <https://doi.org/10.3389/fpls.2021.652116>
- Edmeades, G., Lafitte, H. R., Balanos, J., Chapman, S., Banziger, M., & Deutsch, J. (1994). Developing Maize Program Special Report. CIMMYT. D.F.Mexico.
- Efendi, R. and M. Azrai. 2010. Tanggap genotipe jagung terhadap cekaman kekeringan: Peranan akar. *Jurnal Penelitian*

Pertanian Tanaman Pangan 29(1):1-10

- Efendi, R., Slamet Bambang, P., Arif Subechan, M., Aqil, M., & Azrai, M. (2021). Combining ability of S3 maize inbred lines and related contributing traits for high yield under high population density. *IOP Conference Series: Earth and Environmental Science*, 911(1). <https://doi.org/10.1088/1755-1315/911/1/012009>
- Fan, P., Ming, B., Anten, N. P. R., Evers, J. B., Li, Y., Li, S., & Xie, R. (2022). Plastic response of leaf traits to N deficiency in field-grown maize. *AoB PLANTS*, 14(6), 1–10. <https://doi.org/10.1093/aobpla/plac053>
- Gbegbelegbe S, Chikoye D, Alene A, K.-B. S. and C. G. (2024). Strategic Foresight analysis of droughts in southern Africa and implications for food security. *Front. Sustain. Food Syst.* 7, 1159901. doi: 10.3389/fsufs.2023.1159901.
- Ghosh, M., Swain, D. K., Jha, M. K., Tewari, V. K., & Bohra, A. (2020). Optimizing chlorophyll meter (SPAD) reading to allow efficient nitrogen use in rice and wheat under rice-wheat cropping system in eastern India. *Plant Production Science*, 23(3), 270–285. <https://doi.org/10.1080/1343943X.2020.1717970>
- Gouesnard, B., A. Zanetto, and C. W. (2016). Identification of adaptation traits to drought in collections of maize landraces from southern Europe and temperate regions. *Euphytica* 209, 565–584. doi: 10.1007/s10681-015-1624-8.
- Gupta, N. K., Gupta, S., Singh, J., Garg, N. K., Saha, D., Singhal, R. K. (2022). On-farm hydro and nutri-priming increases yield of rainfed pearl millet through physio-biochemical adjustments and anti-oxidative defense mechanism. *PLoS One* 17, e0265325. doi: 10.1371/journal.pone.0265325.
- Halliday, D.J. and M.E. Trenkel. 1992. IFA World Fertilizer Use Manual. International Fertilizer Industry Association, Paris.
- Hammer, G. L., Dong, Z., McLean, G., Doherty, A., Messina, C., Schussler, J., ... & Cooper, M. (2009). Can changes in canopy and/or root system architecture explain historical maize yield trends in the U.S. corn belt? *Crop Science*, 49(1), 299-312.

- Herlinda, S., Suharjo, R., Elbi Sinaga, M., UGFwazi, F., Suwandi, S. 2021. First report of occurrence of corn and rice strains of fall armyworm, *Spodoptera frugiperda* in South Sumatra, Indonesia and its damage in maize. *Journal of the Saudi Society of Agricultural Sciences*. <https://doi.org/10.1016/j.jssas.2021.11.003>
- Hessl, A. E., Anchukaitis, K. J., Jelsema, C., Cook, B., Byambasuren, O., Leland, C. (2018). Past and future drought in Mongolia. *Sci. Adv.* 4, e1701832–e1701832. doi: 10.1126/sciadv.1701832.
- Huang, Y., Zhu, Y., & Xu, X. (2022). *Yield and trait stability of maize genotypes under nitrogen and planting density stresses in tropical environments*. *Theoretical and Applied Genetics**, 135, 823-835.
- Iriany, R.N., A.M. Takdir, M.H.G. Yasin, and M.J. Mejaya. 2007. Maize genotypes tolerance to drought stress. *Penelitian Pertanian Tanaman Pangan* 26(3):156 - 160
- Jackson, W.A. and Volk, R.J. (1992), Nitrate and ammonium uptake by maize: adaptation during relief from nitrogen suppression. *New Phytologist*, 122: 439-446. <https://doi.org/10.1111/j.1469-8137.1992.tb00071.x>
- Jansen, C., Zhang, Y., Liu, H., Gonzalez-Portilla, P. J., Lauter, N., Kumar, B., Truccillo-Silva, I., Martin, J. P. S., Lee, M., Simcox, K., Schussler, J., Dhugga, K., & Lübberstedt, T. (2015). Genetic and agronomic assessment of cob traits in corn under low and normal nitrogen management conditions. *Theoretical and Applied Genetics*, 128(7), 1231–1242. <https://doi.org/10.1007/s00122-015-2486-0>
- Jafari, F., Wang, B., Wang, H., & Zou, J. (2024). Breeding maize of ideal plant architecture for high-density planting tolerance through modulating shade avoidance response and beyond. *Journal of Integrative Plant Biology*, 66(5), 849–864. <https://doi.org/10.1111/jipb.13603>
- Jhu, M. Y., & Nakayama, H. (2024). Dancing in the sun: maize azimuthal canopy re-orientation for efficient light capture. *Plant Cell*, 36(5), 1568–1569. <https://doi.org/10.1093/plcell/koae026>
- Jones, B. (1998). *Plant Nutrition Manual*. Crc Press, Boston.

- Kalqutny, S. H., Pakki, S., Muis, A. 2020. The potential use of dna based molecular techniques in the study of maize downy mildew. *Jurnal Ilmu dan Teknologi Pertanian Agrosaintek*. 4(1): 17–27.
- Kandel, B. P. (2020). Spad value varies with age and leaf of maize plant and its relationship with grain yield. *BMC Res. Notes* 13, 13–16. doi: 10.1186/s13104-020-05324-7.
- Kementan, 2021: Inilah 10 Provinsi Produsen Jagung Terbesar Indonesia. <https://www.pertanian.go.id/home/?show=news&act=view&id=4639>
- Khatibi, A., S. Omrani, A. Omrani, S.H. Shojaei, S. M. N. M. (2022). Response of Maize Hybrids in Drought-Stress Using Drought Tolerance Indices. *Water (Switzerland)* 14, 1–10. doi: 10.3390/w14071012.
- Kira, O.; Nguy-Robertson, A.L.; Arkebauer, T.J.; Linker, R.; Gitelson, A. A. (2016). Informative spectral bands for remote green LAI estimation in C3 and C4. *Crop. Agric. Meteorol.*, 243–249.
- Leverne, L., & Krieger-Liszkay, A. (2021). Moderate drought stress stabilizes the primary quinone acceptor QA and the secondary quinone acceptor QB in photosystem II. **Physiologia Plantarum*, 171*(2), 260–267. <https://doi.org/10.1111/ppl.13286>
- Lindsay, W.L. (1979) *Chemical Equilibrium in Soils*. John Wiley & Sons, New York.
- Liu, T., Song, F., Liu, S., Zhu, X. 2011. Canopy structure, light interception, and photosynthetic characteristics under different narrow-wide planting patterns in maize at silking stage. *Spanish Journal of Agricultural Research*. 9(4), 1249-1261.
- Liu, Z., Gao, J., Gao, F., Liu, P., Zhao, B., & Zhang, J. (2018). Photosynthetic characteristics and chloroplast ultrastructure of summer maize response to different nitrogen supplies. *Frontiers in Plant Science*, 9(May), 1–13. <https://doi.org/10.3389/fpls.2018.00576>
- Liu, N., Li, L., Li, H., Liu, Z., Lu, Y., & Shao, L. (2023). Selecting maize cultivars to regulate canopy structure and light interception for high yield. *Agronomy Journal*, 115(2), 770–780.

<https://doi.org/10.1002/agj2.21278>

- Liu, G., Yang, Y., Liu, W., Guo, X., Xie, R., Ming, B., Xue, J., Zhang, G., Li, R., Wang, K., Hou, P., & Li, S. (2022). Optimized canopy structure improves maize grain yield and resource use efficiency. *Food and Energy Security*, 11(2), 1–11. <https://doi.org/10.1002/fes3.375>
- Lopez, C., Torres, E., & Gomez, R. (2020). *Genotype-environment interactions in maize yield under combined water and nutrient stress across multiple locations*. *Crop Science*, 60(1), 45-53.
- Mantilla-Perez, M. B., & Salas Fernandez, M. G. (2017). Differential manipulation of leaf angle throughout the canopy: Current status and prospects. *Journal of Experimental Botany*, 68(21–22), 5699–5717. <https://doi.org/10.1093/jxb/erx378>
- Marliyanti, L., Syukur, M., Widodo, W. 2014. Daya hasil 15 galur cabai IPB dan ketahanannya terhadap penyakit antraknosa yang disebabkan oleh *Colletotrichum acutatum*. *AGH Online Journal*. 1(1): 7-13.
- Monneveux, P., Sanchez, C., Beck, D., & Edmeades, G. O. (2006). Drought tolerance improvement in tropical maize source populations: Evidence of progress. *Crop Science*, 46(1), 180-191.
- Monneveux, P., Zaidi, P. H., & Sanchez, C. (2005). Population density and low nitrogen affects yield-associated traits in tropical maize. *Crop Science*, 45(2), 535–545. <https://doi.org/10.2135/cropsci2005.0535>
- Moradi, H., Akbari, G. A., Khorasani, S. K., & Ramshini, H. A. (2012). Evaluation of drought tolerance in corn (*Zea mays* L.) new hybrids with using stress tolerance indices. *Eur. J. Sustain. Dev.* 1, 543. Available at: <https://doi.org/10.14207/ejsd.2012.v1n3p543> .
- Moustakas, M., Sperdouli, I., & Moustaka, J. (2022). Early drought stress warning in plants: Color pictures of photosystem II photochemistry. **Climate*, 10*(11), 179. <https://doi.org/10.3390/cli10110179>
- Magorokosho C., P. K. V. and T. P. (2003). Selection for drought tolerance in two tropical maize populations. *African Crop Sci. J.* 11, 151–161. Available at: <https://www.ajol.info/index.php/acsj/article/view/27566>.
- Mu, X., Chen, Q., Chen, F., Yuan, L., & Mi, G. (2016). Within-leaf nitrogen allocation in adaptation to low nitrogen supply in maize during grain-filling stage. *Frontiers in Plant Science*,

7(MAY2016), 1–11. <https://doi.org/10.3389/fpls.2016.00699>

- Olivoto T., Lucio A., Silva J., Sari B., D. M. (2019). Mean performance and stability in multi-environment trials ii: selection based on multiple traits. *Agron. Journal*, 111(6) 111, 2961–2969. Available at: <https://doi.org/10.2134/agronj2019.03.0221>.
- Panikkai, S., Nurmalinga, R., Mulatsih, S., Purwati, P. 2017. Analisis ketersediaan jagung nasional menuju pencapaian swasembada dengan pendekatan model dinamik. *Informatika Pertanian*, Vol. 26 (1): 41 – 48
- Pettigrew, W. T. (2008). Peran fisiologis nitrogen pada jagung. **Ilmu Tanaman**, 48(1), 35-43.
- Pusdatin, 2022. Analisis kinerja perdagangan jagung. Pusat Data dan Informasi, Kementerian Pertanian.
- Pusdatin, 2021. Analisis kinerja perdagangan jagung. Pusat Data dan Informasi, Kementerian Pertanian.
- Qian, L. (2014). Effect of low nitrogen stress on different low nitrogen tolerance maize cultivars seedling stage growth and physiological characteristics. *Acta Pratacultural Science*. <https://consensus.app/papers/effect-of-low-nitrogen-stress-on-different-low-nitrogen-qian/c435d3df6571516cbb21c7ac78c623d/>
- Radford, P. J. (1967). Growth analysis formulae-their use and abuse 1. *Crop Sci.* 7, 171–175. doi: 10.2135/cropsci1967.0011183X000700030001x.
- Rana, L., Banerjee, H., Mazumdar, D., Sarkar, S., Ray, K., Garai, S., Nayek, J., & Kumar, M. (2021). Determination of Principal Yield Attributing Traits of Hybrid Maize (*Zea mays* L.) Using Multivariate Analysis. *International Journal of Bio-Resource and Stress Management*. <https://doi.org/10.23910/1.2021.2366>
- Rorie, R., Purcell, L., Mozaffari, M., Marsh, M., & Longer, D. (2011). Association of “Greenness” in Corn with Yield and Leaf Nitrogen Concentration. *Agronomy Journal*, 103, 529. <https://doi.org/10.2134/agronj2010.0296>
- Rossi S., Chapman C., H. B. (2020). Suppression of heat-induced

- leaf senescence by γ -aminobutyric acid, proline, and ammonium nitrate through regulation of chlorophyll degradation in creeping bentgrass. *Environ. Exp. Bot.* 177, 104116. doi: 10.1016/j.envexpbot.2020.104116.
- Rossini, F., Vear, F., & Echarte, L. (2011). Breeding maize for density tolerance: Possible exploitation of indirect selection tools. *Crop Science*, 51(4), 1274-1282.
- Saad-Allah, K. M., Nessem, A. A., Ebrahim, M. K. H., and Gad, D. (2022). Evaluation of drought tolerance of five maize genotypes by virtue of physiological and molecular responses. *Agronomy* 12, 59. doi: 10.3390/agronomy12010059.
- Sairam, M., Maitra, S., Sahoo, U., Sagar, L., & Krishna, T. G. (2023). Evaluation of precision nutrient tools and nutrient optimization in maize (*Zea mays* L.) for enhancement of growth, productivity and nutrient use efficiency. *Research on Crops*, 24(4), 666–677. <https://doi.org/10.31830/2348-7542.2023.ROC-1016>
- Setiawan, K., Basri, M. 2017. An Analysis of Efficiency the Production of Commodities Maize in Belu, East Nusa Tenggara, Indonesia. *IOSR Journal of Environmental Science, Toxicology and Food Technology (IOSR-JESTFT)*,. 11(10): 64–69.
- Silva, A. N. da, Schoninger, E. L., Trivelin, P. C. O., Dourado-Neto, D., Pinto, V. M., & Reichardt, K. (2016). Maize Response to Nitrogen: Timing, Leaf Variables and Grain Yield. *Journal of Agricultural Science*, 9(1), 85. <https://doi.org/10.5539/jas.v9n1p85>
- Smakhtin, V. and Schipper, E. L. (2008). Droughts: The Impact of Semantics and Perceptions. *Water Policy* 1, 131–143. Available at: <https://doi.org/10.2166/wp.2008.036>.
- Smith, J.D., Brown, K., & Johnson, R. (2019). *Effects of plant density on yield and resource use efficiency in maize hybrids under low nitrogen conditions*. **Field Crops Research**, 232, 90-98.
- Song, H., Y. Li, L. Zhou, Z. Xu, and G. Z. (2018). Maize leaf functional responses to drought episode and rewetting. *Agric. For. Meteorol.* 249, 57–70. doi: 10.1016/j.agrformet.2017.11.023.
- Sobiech, A., Tomkowiak, A., Nowak, B., Bocianowski, J., Wolko, Ł., Sychała, J. 2022. Associative and physical mapping of

markers related to fusarium in maize resistance, obtained by next-generation sequencing (NGS). *Int. J. Mol. Sci.* 23: 6105. <https://doi.org/10.3390/ijms23116105>

Sobir, Syukur, M. 2012. *Genetika Tanaman*. IPB Press, Bogor, Indonesia.

Su, Z., J. Zhao, T.H. Marek, K. Liu, M. T. H. (2022). Drought tolerant maize hybrids have higher yields and lower water use under drought conditions at a regional scale. *Agric. Water Manag.* 274, 107978. doi: 10.1016/j.agwat.2022.107978.

Subekti, N.A., Syafruddin., Efendi, R., Sunarti, S. 2008. *Morfologi Tanaman dan Fase Pertumbuhan Jagung*. Balai Penelitian Tanaman Serealia, Maros, Indonesia.

Sulaiman, A.A., Kariyasa, K., Hoerudin, Subagyo, K., Suwandi, Bahar, F.A. 2017. *Cara cepat swasembada jagung*. Sekretariat Jendral Kementrian Pertanian RI, Jakarta, Indonesia.

Suwardi dan M. Azrai. 2013. Pengaruh cekaman kekeringan genotipe jagung terhadap karakter hasil dan komponen hasil. p.149-157. *Seminar Nasional Serealia. Meningkatkan Peran Peneliti Serealia Menuju Pertanian Berkelanjutan*. Maros, 18 Juni 2013.

Syukur, M., Sujiprihati, S., Yunianti, R. 2015. *Teknik Pemuliaan Tanaman: Edisi Revisi*. Penebar Swadaya, Depok. Indonesia.

Tian, J., Wang, C., Xia, J., Wu, L., Xu, G., Wu, W., Li, D., Qin, W., Han, X., Chen, Q., Jin, W., & Tian, F. (2019). Teosinte ligule allele narrows plant architecture and enhances high-density maize yields. *Science*, 365(6454), 658–664. <https://doi.org/10.1126/science.aax5482>

Tokatlidis, I. S., & Koutroubas, S. D. (2004). A review of maize hybrids' dependence on high plant populations and its implications for crop yield stability. *Field Crops Research*, 88(2-3), 103-114.

Udo, E. F., Ajala, S. O., & Olaniyan, A. B. (2017). Physiological and morphological changes associated with recurrent selection for low nitrogen tolerance in maize. *Euphytica*, 213(7), 1–13. <https://doi.org/10.1007/s10681-017-1928-y>

- Vacaro, E., Neto, J.F.B., Pegoraro, D.G., Nuss, C.N. and Conceicao., L.D.H. 2002. Combining ability of twelve maize populations. *Pesq. Agropec. Bras.*, 37 (1), 67–72.
- Wang, X., Zhao, J., & Guo, Z. (2018). *Adaptive responses of maize genotypes to nitrogen and density stress in subtropical climates*. *Plant Science*, 269, 26-34.
- Wkdw, I., Wkhlu, G., Surfhvvhv, Y., Oljkw, D. U. H., Zdwhu, K., Qxwulwlyh, D. Q. G., & Ffruglj, P. (2007). The indication of nitrogen deficiency in maize growing using SPAD-502 chlorophyll meter. *Cereal Research Communications*, 35(2), 7–10.
- WU, Y. wei, LI, Q., JIN, R., CHEN, W., LIU, X. lin, KONG, F. lei, KE, Y. pei, SHI, H. chun, & YUAN, J. chao. (2019). Effect of low-nitrogen stress on photosynthesis and chlorophyll fluorescence characteristics of maize cultivars with different low-nitrogen tolerances. *Journal of Integrative Agriculture*, 18(6), 1246–1256. [https://doi.org/10.1016/S2095-3119\(18\)62030-1](https://doi.org/10.1016/S2095-3119(18)62030-1)
- Xue, J., Zhao, Y., Gou, L., Shi, Z., Yao, M., Zhang, W. 2016. “How High Plant Density of Maize Affects Basal Internode Development and Strength Formation.” *Crop Science*, 56(6): 3295–3306.
- Yan, Weikai; Hunt, L.A. (2001). Interpretation of Genotype × Environment Interaction for Winter Wheat Yield in Ontario. *Crop Science*, 41(1), 19–. doi:10.2135/cropsci2001.41119x
- Yang, W., Duan, L., Chen, G., Xiong, L., & Liu, Q. (2017). Plant phenomics and high-throughput phenotyping: accelerating rice functional genomics using multidisciplinary technologies. *Current Opinion in Plant Biology*, 37, 63-73.
- Yang, P., Liu, Z., Zhao, Y., Cheng, Y., Li, J., Ning, J., Yang, Y., Huang, J. 2020. “Comparative Study of Vegetative and Reproductive Growth of Different Tea Varieties Response to Different Fluoride Concentrations Stress.” *Plant Physiology and Biochemistry* 154: 419–428.
- Yuan L., Pu R., Zhang J., Wang J., Yang H. (2016). Using high spatial resolution satellite imagery for mapping powdery mildew at a regional scale. *Precision Agric.* 17, 332–348. 10.1007/s11119-015-9421-x

- Zainuddin, B., and Aqil, M. (2021). Analysis of the relationship between leaf color spectrum and soil plant analysis development. *IOP Conf. Ser. Earth Environ. Sci.* 911. doi: 10.1088/1755-1315/911/1/012045.
- Zhang, L., Wang, F., & Zhao, Y. (2021). *Plant density and nitrogen effects on maize phenology and grain yield under contrasting environmental conditions*. *Agronomy Journal*, 113(5), 1674-1685.
- Zainuddin, B., Syam, E., Azrai, M., Musa, Y., Efendi, R., Priyanto, S. B., Andayani, N. N., Zainuddin, B., Syam, E., Azrai, M., Musa, Y., Efendi, R., Priyanto, S. B., & Andayani, N. N. (2024). Analysis of Plant Ideotype and Yield in Hybrid Maize under Varied Population Densities. 55(7).