

DAFTAR PUSTAKA

1. Asadi-Amoli, F. *et al.* (2015) 'Comparison of Direct Immunofluorescence and Giemsa Staining in Chlamydia trachomatis Follicular Conjunctivitis', *Zahedan Journal of Research in Medical Sciences*. doi: 10.17795/zjrms-2181.
2. BOWEN, V. *et al.* (2017) 'Gonorrhea', *Current Epidemiology Reports*, 4, pp. 1–10. Chan, P. A. *et al.* (2012) 'Recommendations For the Diagnosis of Neisseria gonorrhoeae and Chlamydia trachomatis, Including Extra-Genital Sites', *Med Health R I*, 95(8), pp. 252–4.
3. Cook, R., Hutchison, S. and Ostergaard, L. (2005) 'Systematic review: noninvasive testing for Chlamydia trachomatis and Neisseria gonorrhoeae', *AnnIntern Med*, 142, pp. 914–25.
4. Cosentino, L., Landers, D. and Hillier, S. (2003) 'Detection of Chlamydia trachomatis and Neisseria gonorrhoeae by strand displacement amplification and relevance of the amplification control for use with vaginal swab specimens.', *J ClinMicrobiol*, 41(8), pp. 3592–6.
5. Creighton, S. (2014) 'Gonorrhoea', *BMJ Clin Evid*, 2014, p. 1604.
6. Dela, H. *et al.* (2019) 'Risk factors associated with gonorrhea and chlamydia transmission in selected health facilities in Ghana', *BMC Infect Dis*, 19, p. 425. Dielissen, P. W., Teunissen, D. A. and Lagro-Janssen, A. L. (2013) 'Chlamydia prevalence in the general population: is there a sex difference? a systematic review', *BMC Infect Dis*, 13(534), pp. 1–12.
7. Dubbink, J. H. and Bos, M. (2015) 'Microbiological Characteristics of Chlamydia trachomatis and Neisseria gonorrhoeae Infections in South African Women', *Journal of Clinical Microbiology*, 54(1), pp. 1–10.
8. Hill, S. A., Masters, T. L. and Wachter, J. (2016) 'Gonorrhea - an evolving disease of the new millennium', *Microb Cell*, 3(9), pp. 371–89.
9. Jaschek, G. *et al.* (1993) 'Direct detection of Chlamydia trachomatis in urine specimens from symptomatic and asymptomatic men by using a rapid polymerase chain reaction assay', *Journal of Clinical Microbiology*. doi:10.1128/jcm.31.5.1209-1212.1993.

10. Knox, J. *et al.* (2002) 'Evaluation of self-collected samples in contrast to practitioner-collected samples for detection of chlamydia trachomatis, neisseriagonorrhoeae, and trichomonas vaginalis by polymerase chain reaction among women living in remote areas', *Sexually Transmitted Diseases*. doi: 10.1097/00007435-200211000-00006.
11. Korownyk, C., Kraut, R. Y. and Kolber, M. R. (2018a) 'Vaginal self-swabs for chlamydia and gonorrhea', *Can Fam Physician*, 64(6), p. 448.
12. Korownyk, C., Kraut, R. Y. and Kolber, M. R. (2018b) 'Vaginal self-swabs for chlamydia and gonorrhea', *Canadian Family Physician*.
13. Kresno, S. (2010) 'Polymerase Chain Reaction', in Kresno, S. B. (ed.) *Imunologi: Diagnosis dan Prosedur Laboratorium. Ed 5*. Jakarta: Badan Penerbit Fakultas Kedokteran Universitas Indonesia, pp. 524–7.
14. Lanjouw, E. *et al.* (2015) '2015 European guideline on the management of Chlamydia trachomatis infections', *Int J STD AIDS*, 1(2), pp. 1–16.
15. Meyer, T. (2016) 'Diagnostic Procedures to Detect Chlamydia trachomatis Infections', *Microorganisms*. doi: 10.3390/microorganisms4030025.
16. Meyer, T. and Buder, S. (2020) 'The laboratory diagnosis of neisseria gonorrhoeae: Current testing and future demands', *Pathogens*, 9(2), p. 91. doi: 10.3390/pathogens9020091.
17. Omer, H. M. H. *et al.* (2020) 'Cytological and molecular screening of Chlamydia trachomatis in infertile women attending a maternity teaching hospital in Gezira State, Sudan: a cross-sectional study', *F1000Research*. doi: 10.12688/f1000research.23490.1.
18. Papp, J. R. *et al.* (2014) 'Recommendations for the laboratory-based detection of Chlamydia trachomatis and Neisseria gonorrhoeae - 2014', *MMWR Recommendations and Reports*.
19. Philip A. Chan *et al.* (2016) 'Extragenital Infections Caused by Chlamydia trachomatis and Neisseria gonorrhoeae: A Review of the Literature', *Infectious Diseases in Obstetrics and Gynecology*, 1(2), pp. 1–17.
20. Piszczek, J. and Jean, R. S. (2015) 'Gonorrhea', *Can Pharm J*, 148(2), pp. 82–89. Van Der Pol, B. *et al.* (2020) 'Evaluation of the Performance of a Point-

- of-Care Test for Chlamydia and Gonorrhoea’, *JAMA Network Open*. doi: 10.1001/jamanetworkopen.2020.4819.
21. Quillin, S. J. and Seifert, H. S. (2018) ‘Neisseria gonorrhoeae host-adaptation and pathogenesis’, *Nat Rev Microbiol*, 16(4), pp. 226–240.
 22. Rodrigues, J. F. *et al.* (2016) ‘Diarrheogenic Escherichia coli’, *Brazilian Journal of Microbiology*. Sociedade Brasileira de Microbiologia, 47, pp. 3–30. doi: 10.1016/j.bjm.2016.10.015.
 23. Rönn, M. *et al.* (2019) ‘Evaluation of the performance of nucleic acid amplification tests (NAATs) in detection of chlamydia and gonorrhoea infection in vaginal specimens relative to patient infection status: a systematic review.’, *BMJ Open*, 9(1), p. e022510.
 24. Rosen, T. (2012) ‘Gonorrhoea, Mycoplasma, and vaginosis.’, in *Fitzpatrick’s dermatology in general medicine*. New York: Mc Graw-Hill, pp. 2514–2519.
 - Rowley, L. N. J. *et al.* (2015) ‘Global Estimates of the Prevalence and Incidence of Four Curable Sexually Transmitted Infections in 2012 Based on Systematic Review and Global Reporting’, *PLoS One*, 10(12), p. e0143304.
 25. Schachter, J. (2001) ‘NAATs to diagnose Chlamydia trachomatis genital infection: A promise still unfulfilled’, *Expert Review of Molecular Diagnostics*. doi: 10.1586/14737159.1.2.137.
 26. Schachter, J. *et al.* (2005) ‘Vaginal swabs are the specimens of choice when screening for Chlamydia trachomatis and Neisseria gonorrhoeae: Results from a multicenter evaluation of the APTIMA assays for both infections’, in *Sexually Transmitted Diseases*. doi: 10.1097/01.olq.0000190092.59482.96.
 27. Thomas R. Frieden and Jaffe, H. W. (2015) ‘Sexually Transmitted Diseases Treatment Guidelines, 2015’, *MMWR Recomm Rep*, 24, pp. 55–57.
 28. Thorley, N. and Radcliffe, K. (2015) ‘The performance and clinical utility of cervical microscopy for the diagnosis of gonorrhoea in women in the era of the NAAT’, *International Journal of STD and AIDS*. doi: 10.1177/0956462414551233.
 29. Walker, C. K. and Sweet, R. L. (2011) ‘Gonorrhoea infection in women: prevalence, effects, screening, and management’, *Int J Womens Health*, 3, pp.

197–206.

30. Wilcox, M. H. *et al.* (2000) ‘Combined cervical swab and urine specimens for PCR diagnosis of genital *Chlamydia trachomatis* infection’, *Sexually Transmitted Infections*. doi: 10.1136/sti.76

LAMPIRAN

Tabel 1. Hasil pemeriksaan sampel serviks pada pasien gonore

			Gonorrhoea		
			Terdiagnosis	Bukan	Total
			GO	GO	
PCR	Sampel	Positif	12	0	12
	serviks	Negatif	5	59	64
Total			17	59	76

Sensitivitas = $(a/n1) \times 100\% = 12/17 \times 100\% = 70.5\%$

Spesifisitas = $(d/n2) \times 100\% = 59/59 \times 100\% = 100\%$

Nilai prediksi positif = $(a/n3) \times 100\% = 12/12 = 100\%$

Nilai prediksi negatif = $(d/n4) \times 100\% = 59/64 \times 100\% = 92.1\%$

Akurasi = $a+d / N \times 100\% = 16+59/ 76 \times 100\% = 93.4 \%$

Tabel 2. Hasil pemeriksaan sampel vagina pada pasien gonore

			Gonorrhoea		
			Terdiagnosis GO	Bukan	Total
				GO	
PCR	Sampel	Positif	16	0	16
	dinding vagina	Negatif	1	59	60
Total			17	59	76

Sensitivitas = $(a/n1) \times 100\% = 16/17 \times 100\% = 94.1\%$

Spesifisitas = $(d/n2) \times 100\% = 59/59 \times 100\% = 100\%$

Nilai prediksi positif = $(a/n3) \times 100\% = 16/16 = 100\%$

Nilai prediksi negatif = $(d/n4) \times 100\% = 59/60 \times 100\% = 98.3\%$

Akurasi = $a+d / N \times 100\% = 16+59/ 76 \times 100\% = 98.6 \%$

Tabel 3. Hasil pemeriksaan sampel vagina pada pasien chlamydia

Chlamydia			Chlamydia		
			Terdiagnosis	Bukan	Total
			chalmydia	chlamydia	
PCR	Sampel	Positif	31	0	31
	dinding vagina	Negatif	3	42	45
Total			34	42	76

$$\text{Sensitivitas} = (a/n1) \times 100\% = 91.1\%$$

$$\text{Spesifisitas} = (d/n2) \times 100\% = 44/44 \times 100\% = 100\%$$

$$\text{Nilai prediksi positif} = (a/n3) \times 100\% = 31/31 = 100\%$$

$$\text{Nilai prediksi negatif} = (d/n4) \times 100\% = 44/45 \times 100\% = 93.3\%$$

$$\text{Akurasi} = a+d / N \times 100\% = 31+44 / 76 \times 100\% = 96.05 \%$$

Tabel 4. Hasil pemeriksaan sampel serviks pada pasien chlamydia

Chlamydia			Chlamydia		
			Terdiagnosis	Bukan	Total
			chalmydia	chlamydia	
PCR	Sampel	Positif	32	0	32
	serviks	Negatif	3	41	44
Total			35	41	76

$$\text{Sensitivitas} = (a/n1) \times 100\% = 14/32 \times 100\% = 91.4\%$$

$$\text{Spesifisitas} = (d/n2) \times 100\% = 44/44 \times 100\% = 100\%$$

$$\text{Nilai prediksi positif} = (a/n3) \times 100\% = 14/14 = 100\%$$

$$\text{Nilai prediksi negatif} = (d/n4) \times 100\% = 93.1\%$$

$$\text{Akurasi} = a+d / N \times 100\% = 14 + 44 / 76 \times 100\% = 96.05\%$$



