

DAFTAR PUSTAKA

1. Lewis TV, Malviya S. A prospective cohort study of emergence agitation in the pediatric postanesthesia care unit. *Anesth Analg* 2003;96:1625-30.
2. Mason LJ. Emergence delirium. In: Pitfalls of pediatric anesthesia, New York: Loma Linda University Publisher;2010.p.150-5.
3. Braz GL, Muniz LD, Modolo SN. Emergence agitation in pediatric anesthesia: current features. *J pediatr (Rio J)* 2008;84(2):107-113.
4. Nasr VG, Hannallah RS. Emergence agitation in children: a review. *M.E. J. Anesth* 2011;21(2):175-9.
5. Mountain BW, Smithson L, Cramolini M, Wyatt TH, Newman M. Dexmedetomidine as a pediatric anesthetic premedication to reduce anxiety and to deter emergence delirium. *AANA J* 2011;79(3):219-23.
6. Meng QT, Xia ZY, Luo T, Wu Y, Tang LH, Zhao B, Chen JH, Chen X. Dexmedetomidine reduce emergence agitation after tonsillectomy in children by sevofluran anesthesia: a case-control study. *Int J Pediatr Otorhinolaryngol* 2012;28(3):1-6.
7. Syukry M, Clyde MC, Kalarickal PL, Ramadyani U. Does dexmedetomidine prevent emergence delirium in children after sevofluran-based general anesthesia?. *Pediatr Anest J* 2005;15(12):1098-104.
8. Syukry M, Cain JG. Dexmedetomidine prevents and treats agitation, delirium, and withdrawl. *Int J Trauma* 2007;17(1):24-6.
9. Guler G, Akin A, Tosun Z, Ors S, Esmaoglu A, Boyaci A. Single-dose dexmedetomidine reduces agitation and provides smooth extubation after pediatric adenotonsillectomy. *Pediatric Anesthesia* 2005;15:762-6
10. Ibacache ME, Munoz HR, Brandez V, Morales AR. Single dose dexmedetomidine reduce agitation after sevofluran anesthesia in children. *Anesth Analg* 2004;98(1):60-63.
11. Isik B, Arslan B, Tunga AD, Kurtipek O, Dexmedetomidine decrease emergence agitation in pediatric patients after sevofluran anesthesia without surgery. *Pediatr anesth* 2006;16(7):748-53.

12. Pieters BJ, Penn E, Nicklaus P, Bruegger D, Mehta B, Wheately R. Emergence delirium and postoperative pain in children undergoing adenotonsillectomy: a comparison of propofol vs sevofluran anesthesia. *Pediatr Anesth J* 2010;10(3):1-7.
13. Shari M, Burns, RN. Delirium during emergence from anesthesia: a case study. *J Crit Care Nurs* 2003;23(1):102-10.
14. Brown EN, Lydic R, Schiff ND. General anesthesia, sleep, and coma. *N Engl J Med* 2010;36(3):27-30.
15. Tobias JD. Dexmedetomidine: Applications in pediatric critical care and pediatric anesthesiology. *Pediatr Crit Care Med* 2007;8:115-131.
16. Maldonado JR. Delirium in the acute care setting characteristics, diagnosis, and treatment. *J Crit Care Clin* 2008;24:657-722.
17. Kim JH. Mechanism of emergence agitation induced by sevoflurane anesthesia. *Korean J Anesth* 2011;60(2):73-4.
18. Weldon BC, Watcha MF, White PF. Oral midazolam in children: effect of time and adjunctive therapy. *Anesth Analg* 1992;75(5):51-55.
19. Easley B, Tobias JD. Pro: Dexmedetomidine should be used for infants and children undergoing cardiac surgery. *J Cardiothoracic Anesth* 2008;22(1):147-51.
20. McClain BC. Newer modalities for pain managements. Presented at SPA Annual Meeting. 2006: 1-6.
21. Schieveld JNM. Paediatric delirium: where do we go from here?. an update on key issues and research questions. *Neth J Crit Care* 2010; 14(5):330-34.
22. Panzer O, Moitra V, Sladen R. Pharmacology of sedative-analgesic: dexmedetomidine, remifentanil, ketamin, volatile anesthetics, and the role of peripheral mu antagonists. *J Anesth Clinics* 2011;29(4):250-65.
23. Aarts A, Hagen VV, Russchen H. Does pharmacologic treatment prevent children from emergence agitation after sevofluran anesthesia? a systematic review. *Eras J Med* 2012;2(2):24-8.
24. Malviya S. Assessment of pain in children. Presennted at SPA annual meeting. 2006: 3-6.

25. Meyer RR, Munster P, Werner C, Barmbrink AM. Isoflurane is associated with a similar incidence of agitation/delirium as sevoflurane in young children: a randomised controlled study. *Pediatr Anesth* 2007;17(1):56-60.
26. Samira A, Bajwa FR, Fanzha DC, Allan M. A Comparison of emergence delirium scales following general anesthesia in children. *Pediatr Anaesth J* 2010;20(8):704-11
27. Dahlan MS. Menentukan besaran sampel dalam langkah-langkah membuat proposal penelitian bidang kedokteran dan kesehatan. Edisi 2. Jakarta: Sagung Seto;2012.p.79-85.