

## **DAFTAR PUSTAKA**

1. Salmah GS, Choy YC. Comparison of morphine with fentanyl added to intrathecal 0,5% hyperbaric bupivacaine for analgesia after caesarean section. Med J Malaysia, 2009; 64 : 71-4.
2. Morgan GE, Mikhail MS. Spinal, epidural and caudal blocks in clinical anesthesiology. 4<sup>th</sup> ed. New York: McGraw-Hill; 2006 : 289-323.
3. Edna Z, Yossi N, Uri R. Intrathecal anesthesia for elderly patients undergoing short transurethral procedures: A dose-finding study. JRMS. Dec 2002 : 43-6.
4. Patra P, Kapoor MC, Nair TGM. Spinal anesthesia with low dose bupivacaine and for endoscopic urological surgeries. J anesth clin pharmacol. 2005; 21(2) : 147-54.
5. Stoelting RK, Hillier SC (ed). Pharmacology & physiology in anesthetic practice. 4<sup>th</sup>ed. London: Lippincott Williams & Wilkins; 2006 : 190.
6. Palmer CM, Wallace M. Bupivacaine augments intrathecal fentanyl for labor analgesia. Anesthesiology, 1999; 91(1) : 84-9.
7. Khanna MS, Singh IK. Comparative evaluation of bupivacaine plain versus bupivacaine with in spinal anaesthesia in geriatric patients. Indian J anaesth. 2002; 46(3) : 199-203.
8. Grewal A. Dexmedetomidine. J anesteshiol clin pharmacol. 2011; 27(3) : 297-302.
9. Peter FD. Clinical anesthesia procedures of the Massachusetts general Hospital. 7<sup>th</sup>ed. 2007 : 247-72.
- 10.Paech MJ. Anesthesia for caesarean delivery. Obstetric anesthesia. New York: Oxford University Press; 2011 : 79-155.
- 11.Scott J, Flood P. Anesthesia for caesarean delivery. Obstetric anesthesia. 1<sup>st</sup>ed. Philadelphia: Mosby Elsevier; 2006.

- 12.Tsen LC. Anesthesia for caesarean delivery. Chesnut's obstetric anesthesia: principles and practice. 4<sup>th</sup>ed. Philadelphia: Mosby Elsevier; 2009.
- 13.Wong CA. Neuraxial anesthesia for adult genitourinary procedures in spinal and epidural anesthesia. New York: McGraw-Hill; 2007 : 237-46.
- 14.Hocking G, Wildsmith JA. Intrathecal drug spread. British J Anesth, 2004; 93(4) : 568-78.
- 15.Duke J. Anesthesia secrets . 3<sup>rd</sup>ed. 2006 : 500-6.
- 16.Yadav A. Short text of anesthesia. 2<sup>nd</sup>ed. 2004 : 116-27.
- 17.Cianni S, Rossi M, Casta A. Spinal anesthesia: an evergreen technique. Acta Biomed, 2008; 79 : 9-17.
- 18.Jeffrey H, Silverstein G, Alec R. Geriatric anesthesiology. 2<sup>nd</sup>ed. Berlin: Springer; 2007 : 3-37.
- 19.Kirsten M, Laura M. Bohn. Mu opioid receptor regulation and opiate responsiveness. The AAPS Journal. 2005 : 7(3).
- 20.Liu SS, McDonald SB. Current issues in spinal anesthesia. Anesthesiology. 2001 : 888–906.
- 21.Bakshi U. Adjuvant drugs in central neuraxial analgesia. J Anesth. 2010.
- 22.Miller RD. Bupivacaine. 6<sup>th</sup>ed. New York: Elsevier, Churchill Livingstone; 2006.
- 23.Beloeil H, Gentili M, Benhamou D. The effect of a peripheral block on inflammation-induced prostaglandin E2 and cyclooxygenase expression in rats. Anesth Analg, 2009; 109 : 943-50.
- 24.Chahl LA. Opioids - mechanisms of action. Aust Prescr, 1996; 19 : 63-5.

25. Eisenach MD. Alpha 2 adrenergic agonists for regional anesthesia. *Anesthesiology*. 1996.
26. Christiansson L. Update on adjuvant in regional anesthesia. *Period biol*, 2009; 111(2) : 161-70.
27. Yazbek VG, Aouad MM. Perioperative uses of dexmedetomidine. *M.E.J. Anesth*, 2006; 18(6) : 1043-58.
28. Gupta R, Bogra J, Verma R. Dexmedetomidine as an intrathecal adjuvant for postoperative analgesia. *Indian J Anaesth*, 2011; 55 : 347-51.
29. Guo TZ, Jiang JY, Buttermann AE. Dexmedetomidine injection into the locus ceruleus produces antinociception. *Anesthesiology*, 1996; 84(4) : 873-81.
30. Eid H, Shafie M, Youssef H. Dose-related prolongation of hyperbaric bupivacaine spinal anesthesia by dexmedetomidine. *Ain Shams J Anaesth*, 2011; 4(2) : 83-95.
31. Amin A, Fares K, Abd-Elbaky S. Efficacy of intrathecally administered dexmedetomidine versus fentanyl in patients undergoing major abdominal cancer surgery. *Pain physician*, 2012; 15 : 339-48.
32. Al-Mustafa M, Abu-Halaweh S. Effect of dexmedetomidine added to spinal bupivacaine of urological procedures. *Saudi Med J*, 2009; 30(3) : 365-70.
33. Kanazi GE, Aouad MT, Jabbour-Khoury SI. Effect of low-dose dexmedetomidine or clonidine on characteristics of bupivacaine spinal block. *Acta Anaesthesiol Scand*, 2006; 50(2) : 222-7.
34. Fyneface-Ogan S, Gogo O, Enyindah C. Comparative effects of single shot intrathecal bupivacaine with dexmedetomidine and bupivacaine with fentanyl on labor outcome. *Anesthesiology*, 2012 : 1-4.

