



## Rhodopsin promoter-EGFP fusion transgene expression in photoreceptor neurons of retina and pineal complex in mice

Andi Muhammad Ichsan<sup>a,b</sup>, Ichiro Kato<sup>a,c,\*</sup>, Toshiko Yoshida<sup>d</sup>, Kumi Takasawa<sup>a</sup>,  
Seiji Hayasaka<sup>b,c</sup>, Koichi Hiraga<sup>a</sup>

<sup>a</sup> Department of Biochemistry, Toyama Medical and Pharmaceutical University, School of Medicine, Sugitani 2630, Toyama 930-0194, Japan

<sup>b</sup> Department of Ophthalmology, Toyama Medical and Pharmaceutical University, School of Medicine, Toyama 930-0194, Japan

<sup>c</sup> COE Program in the 21st Century, Toyama Medical and Pharmaceutical University, School of Medicine, Toyama 930-0194, Japan

<sup>d</sup> Department of Anatomy, Toyama Medical and Pharmaceutical University, School of Medicine, Toyama 930-0194, Japan

Received 23 September 2004; received in revised form 3 December 2004; accepted 21 December 2004

### Abstract

Light detection in vertebrate eyes is mediated through retinal photoreceptor rod and cone cells that transduce light signals into electrical responses. The differentiation and synaptogenesis of photoreceptor cells are especially important since these cells are the main targets of degeneration in retinitis pigmentosa. We produced transgenic mice that express enhanced green fluorescent protein (EGFP) under the control of the mouse rhodopsin gene promoter. In Western blot analyses of transgenic retinal homogenates, expression of the endogenous rhodopsin gene was detected from post-natal day (P)8; however, EGFP expression in transgenic retinas was initially detected at P12, indicating delayed expression of the transgene. At P14, fluorescence microscopy showed a weak expression of EGFP in the transgenic retina. In the adult transgenic mice, the strongest EGFP expression was observed in the outer nuclear layer, and to a lesser extent in the outer plexiform layer as well as in the inner and outer segments. EGFP expression was also observed in the pineal stalk. The rhodopsin promoter-EGFP transgenic mice will be not only useful to assess rhodopsin gene promoter activity *in vivo*, but also for retinal transplant studies as a source of functional photoreceptor cells that are fluorescent green.

© 2005 Elsevier Ireland Ltd. All rights reserved.

**Keywords:** Transgenic mice; EGFP; Photoreceptor; Rhodopsin; Promoter; Retina


The retina has been the target of many developmental studies, because of interest in using the retina as a model for the central nervous system development. Histological studies of mammalian retinal neurogenesis have revealed the processes of neuroepithelial cell development into the highly-organized, multilayered retina [24]. Cell lineage tracing in mammals revealed that retinal progenitor cells are multipotent and retain their ability to generate various cell types such as ganglion cells, horizontal cells, cone-photoreceptors, amacrine cells, rod-photoreceptors, bipolar cells and Müller glia cells [17,28]. Many recent studies are focusing on the transcription factors required for retinal cell diversification

[1,3,7,10,18,19,23,25,26]. In many cases of human retinal degenerative disease, photoreceptor cells are the targets of neurodegeneration, while the neurons of the inner retina are well preserved [24].






Rhodopsin, which is present in mature rod-photoreceptor cells, is a 40-kDa transmembrane apoprotein that transduces light signals into a G-protein mediated signal cascade [16]. A point mutation in exon 1 of the human opsin gene has been shown to cause one form of autosomal dominant retinitis pigmentosa [8]. To date, more than 100 mutations in opsin gene which can lead to retinitis pigmentosa are reported (see for example tables in RetNet: <http://www.sph.uth.tmc.edu/Retnet/home.htm>). Interestingly, rhodopsin-null mice almost completely lose their photoreceptor cells within 3 months after birth [11]. These


\* Corresponding author. Tel.: +81 76 434 7227; fax: +81 76 434 5014.  
E-mail address: [ichikato@ms.toyama-mpu.ac.jp](mailto:ichikato@ms.toyama-mpu.ac.jp) (I. Kato).





# Neuroscience Letters





## Neuroscience Letters

[Guide for Authors](#)

[Submit Your Paper](#)

[Track Your Paper](#)

[Order Journal](#)

[View Articles](#)

[Open Access Options](#)

[Most Downloaded Articles](#)

[Recent Open Access Articles](#)

[Most Cited Articles](#)

[Special Issues](#)

[Recent Articles](#)

[News](#)

### Stay up-to-date

Register your interests and receive email alerts tailored to your needs

[Click here to sign up](#)

### Follow us

## Neuroscience Letters Editorial Board

### Editor-in-Chief

S.G. Waxman

Yale University School of Medicine, New Haven, Connecticut, USA

### Deputy Editors

Ausim Azzi

Temple University Hospital, Philadelphia, Pennsylvania, USA

Pamela Knapp

Virginia Commonwealth University, Richmond, Virginia, USA

### Associate Editors

B.A. Barres

Stanford University School of Medicine, Stanford, California, USA

J. Black

Yale University School of Medicine, New Haven, Connecticut, USA

T. Bonhoeffer

Max Planck Institut (MPI) für Neurobiologie, Martinsried, Germany

G. Bottini

Università degli Studi di Pavia, Pavia, Italy

T. Cummins

Indiana University School of Medicine, Indianapolis, Indiana, USA

R.D. Fields

National Institute of Child Health and Human Development (NICHD), Bethesda, Maryland, USA

K. Friston

A. Fuglevand

University of Arizona, Tucson, Arizona, USA

F.H. Gage

The Salk Institute for Biological Studies, La Jolla, California, USA

S. Grillner