

## DAFTAR PUSTAKA

- Abdulkader, M.D., T. (2008) 'PCR versus ELISA in diagnosis of human toxoplasmosis in Jeddah, Saudi Arabia', 38(3), pp. 707–714.
- Aramini, J. J. et al. (1999) 'Potential contamination of drinking water with Toxoplasma gondii oocysts', *Epidemiology and Infection*, 122(2), pp. 305–315. doi: 10.1017/S0950268899002113.
- Arias, M. L. et al. (1996) 'Seroepidemiology of Toxoplasmosis in humans: Possible transmission routes in Costa Rica', *Revista de Biología Tropical*, 44(2), pp. 377–381.
- Ayi, I. et al. (2010) 'Sero-epidemiology of toxoplasmosis amongst pregnant women in the greater Accra region of Ghana', *Ghana Medical Journal*, 43(3), pp. 107–114. doi: 10.4314/gmj.v43i3.55325.
- Bamba, S. et al. (2017) 'Seroprevalence and risk factors of Toxoplasma gondii infection in pregnant women from Bobo Dioulasso, Burkina Faso', *BMC Infectious Diseases*. BMC Infectious Diseases, 17(1), pp. 1–6. doi: 10.1186/s12879-017-2583-6.
- Begeman, I. J. et al. (2017) 'Point-of-care testing for Toxoplasma gondii IgG/IgM using Toxoplasma ICT IgG-IgM test with sera from the United States and implications for developing countries', *PLoS Neglected Tropical Diseases*, 11(6), pp. 1–15. doi: 10.1371/journal.pntd.0005670.

- Ben-David, H. (2006) 'Congenital Toxoplasmosis-Prenatal Aspects of Toxoplasma gondii infection', *Reproductive Toxicology*, 21(4), pp. 458–472.
- Berger, F. et al. (2009) 'Toxoplasmosis among pregnant women in France: Risk factors and change of prevalence between 1995 and 2003', *Revue d'Epidemiologie et de Sante Publique*, 57(4), pp. 241–248. doi: 10.1016/j.respe.2009.03.006.
- Bitnun, A. R. I. and Greg, H. A. (2019) *Fetal Implications of Maternal Infections in Pregnancy*. Fourth Edition, *Infectious Diseases*. Fourth Edition. Elsevier Ltd. doi: 10.1016/B978-0-7020-6285-8.00056-3.
- Centers for Disease Control and Prevention (2018) *Parasites-Toxoplasmosis (Toxoplasma Infection)*. Available at: <https://www.cdc.gov/parasites/toxoplasmosis/> (Accessed: 25 November 2020).
- Choi, W.-Y. et al. (1997) 'Foodborne Outbreaks of Human Toxoplasmosis', *The Journal of Infectious Diseases*, 5(175), pp. 1280–2.
- Cong, W. et al. (2015) 'Toxoplasma gondii Infection in Pregnant Women: A Seroprevalence and Case-Control Study in Eastern China', *BioMed Research International*. Hindawi Publishing Corporation, 2015. doi: 10.1155/2015/170278.
- Contini, C. (2014) 'Toxoplasmosis', in *The Infection Diseases VIII*, pp. 633–45. doi: 10.1016/B978-0-323-59648-0.00154-1.

- Darde, M. (2019) 'Seroprevalence and Risk Factors of *Toxoplasma gondii* Infection in Free-Range Chickens in Senegal, West Africa 1 ' , 1', XX(Xx), pp. 1–7. doi: 10.1089/vbz.2019.2481.
- Dubey, J. (2004) 'Toxoplasmosis-a waterborne zoonosis', *Veterinary Parasitology*, 126(1–2), pp. 57–72.
- Fajardo, H. V. et al. (2013) 'Seroprevalence and risk factors of toxoplasmosis in cattle from extensive and semi-intensive rearing systems at Zona da Mata, Minas Gerais state, Southern Brazil', *Parasites and Vectors*, 6(1), pp. 1–8. doi: 10.1186/1756-3305-6-191.
- Fiedler, K. et al. (1999) 'Toxoplasmosis-antibody seroprevalence in Mecklenburg-Western Pomerania', *Pubmed*, 121 95), pp. 239–43.
- Flegr, J. et al. (2014) 'Toxoplasmosis - A global threat. Correlation of latent toxoplasmosis with specific disease burden in a set of 88 countries', *PLoS ONE*, 9(3). doi: 10.1371/journal.pone.0090203.
- Fromont, E. G., Riche, B. and Rabilloud, M. (2009) 'Toxoplasma seroprevalence in a rural population in France: Detection of a household effect', *BMC Infectious Diseases*, 9, pp. 1–7. doi: 10.1186/1471-2334-9-76.
- Gandahusada, S. (1991) 'Study on the prevalence of toxoplasmosis in Indonesia: a review.', *The Southeast Asian journal of tropical medicine and public health*, 22 Suppl, pp. 93–98.
- Heukelbach, J. et al. (2007) 'Waterborne toxoplasmosis, northeastern

Brazil', *Emerging Infectious Diseases*, 13(2), pp. 287–289. doi: 10.3201/eid1302.060686.

Jiang, R. et al. (2018) 'Seroprevalence and associated risk factors of Toxoplasma gondii among Manchu pregnant women in northeastern China', *Microbial Pathogenesis*. Elsevier Ltd, 123(May), pp. 398–401. doi: 10.1016/j.micpath.2018.07.041.

Jones, J. and Dubey, J. (2010) 'Waterborne toxoplasmosis-recent developments', *Experimental Parasitology*, 124(1), pp. 10–25.

Jones, J. L. and Holland, G. N. (2010) 'Short report: Annual burden of ocular toxoplasmosis in the United States', *American Journal of Tropical Medicine and Hygiene*, 82(3), pp. 464–465. doi: 10.4269/ajtmh.2010.09-0664.

Karunajeewa, H. et al. (2001) 'Seroprevalence of varicella zoster virus, parvovirus B19 and Toxoplasma gondii in a Melbourne obstetric population: Implications for management', *Australian and New Zealand Journal of Obstetrics and Gynaecology*, 41(1), pp. 23–28. doi: 10.1111/j.1479-828X.2001.tb01289.x.

Kaye, A. (2011) 'Toxoplasmosis: Diagnosis, treatment, and prevention in congenitally exposed infants', *Journal of Pediatric Health Care*. Elsevier Ltd, 25(6), pp. 355–364. doi: 10.1016/j.pedhc.2010.04.008.

Konishi, E. et al. (2000) 'High Prevalence of Antibody to Toxoplasma gondii among Humans in Surabaya, Indonesia', 53, pp. 238–241.

Liu, X. *et al.* (2017) 'Detection of Toxoplasma gondii in chicken and soil of chicken farms in Nanjing region, China', pp. 1–8. doi: 10.1186/s40249-017-0277-3.

Lykins, J. *et al.* (2016) 'Understanding toxoplasmosis in the United States through "large data" analyses', *Clinical Infectious Diseases*, 63(4), pp. 468–475. doi: 10.1093/cid/ciw356.

Lykins, J. *et al.* (2018) 'Rapid, inexpensive, fingerstick, whole-blood, sensitive, specific, point-of-care test for anti-Toxoplasma antibodies', *PLoS Neglected Tropical Diseases*, 12(8), pp. 3–9. doi: 10.1371/journal.pntd.0006536.

Mahdy, M. A. K. *et al.* (2017) 'A community-based survey of Toxoplasma gondii infection among pregnant women in rural areas of Taiz governorate, Yemen: The risk of waterborne transmission', *Infectious Diseases of Poverty*. *Infectious Diseases of Poverty*, 6(1), pp. 4–9. doi: 10.1186/s40249-017-0243-0.

Many, A. and Koren, G. (2006) 'Toxoplasmosis during pregnancy.', *Canadian family physician Medecin de famille canadien*, 52, pp. 29–30, 32. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16477906%0Ahttp://www.ncbi.nlm.nih.gov/articlerender.fcgi?artid=PMC1479740>.

Mcleod, R., VanTubbergen, C. and Boyer, K. M. (2016) Chapter 290 – *Toxoplasmosis (Toxoplasma gondii)*. Twentieth, *Nelson Textbook of*

*Pediatrics*. Twentieth. Elsevier Inc. doi: 10.1016/B978-1-4557-7566-8.00290-8.

Minbaeva, G. et al. (2013) 'Toxoplasma gondii Infection in Kyrgyzstan: Seroprevalence, Risk Factor Analysis, and Estimate of Congenital and AIDS-Related Toxoplasmosis', *PLoS Neglected Tropical Diseases*, 7(2), pp. 1–7. doi: 10.1371/journal.pntd.0002043.

Montoya, Jose G., John C. Boothroyd, J. A. K. (2018) 'Toxoplasma gondii', in *Gerald M, Bennett JE, Dolin R, editors. Mandell, Douglas, and Bennett's Principles and practice of infectious diseases*. Elsevier, pp. 3122–3153. doi: 10.1016/B978-0-323-40161-6.00280-1.

Montoya, J. G. (2002) 'Laboratory Diagnosis of Toxoplasma gondii Infection and Toxoplasmosis', *The Journal of Infectious Diseases*, 185, pp. S73-82.

Montoya, J. G. (2019) 'Toxoplasmosis', in, pp. 2125–32. doi: 10.1016/B978-1-4557-5017-7.00349-4.

Montoya, J. G. and Liesenfeld, O. (2004) 'Toxoplasmosis', 363, pp. 1965–1976.

Morrison, D. A. and Höglund, J. (2005) 'Testing the hypothesis of recent population expansions in nematode parasites of human-associated hosts', *Heredity*, 94(4), pp. 426–434. doi: 10.1038/sj.hdy.6800623.

Mose, J. M. et al. (2016) 'Detection of Natural Toxoplasma gondii Infection in Chicken in Thika Region of Kenya Using Nested Polymerase Chain

Reaction', *BioMed Research International*, 2016. doi: 10.1155/2016/7589278.

Murebwayire, E. et al. (2017) 'Seroprevalence and risk factors of Toxoplasma gondii infection among pregnant women attending antenatal care in Kigali , Rwanda', 19(1), pp. 2–9.

Naot, Y. and Remington, J. S. (1980) 'An enzyme-linked immunosorbent assay for detection of IgM antibodies to Toxoplasma gondii: Use for diagnosis of acute acquired toxoplasmosis', *Journal of Infectious Diseases*, 142(5), pp. 757–766. doi: 10.1093/infdis/142.5.757.

Nissapatorn, V. et al. (2011) 'Toxoplasmosis-serological evidence and associated risk factors among pregnant women in southern Thailand', *American Journal of Tropical Medicine and Hygiene*, 85(2), pp. 243–247. doi: 10.4269/ajtmh.2011.10-0633.

Retmanasari, A. et al. (2017) 'Prevalence and Risk Factors for Toxoplasmosis in Middle Java, Indonesia', *EcoHealth*. Springer US, 14(1), pp. 162–170. doi: 10.1007/s10393-016-1198-5.

Robert-gangneux, F. and Dardé, M. (2012) 'Epidemiology of and Diagnostic Strategies for Toxoplasmosis Epidemiology of and Diagnostic Strategies for Toxoplasmosis', 25(2). doi: 10.1128/CMR.05013-11.

Rostami, A., Karanis, P. and Fallahi, S. (2018) 'Advances in serological , imaging techniques and molecular diagnosis of Toxoplasma gondii infection', *Infection*. Springer Berlin Heidelberg. doi: 10.1007/s15010-017-

1111-3.

Singh, S. et al. (2014) 'Serologic Prevalence of Toxoplasma gondii in Indian Women of Child Bearing Age and Effects of Social and Environmental Factors', *PLoS Neglected Tropical Diseases*, 8(3). doi: 10.1371/journal.pntd.0002737.

SMITH, J. L. (1991) 'Foodborne Toxoplasmosis', *Journal of Food Safety*, 12(1), pp. 17–57. doi: 10.1111/j.1745-4565.1991.tb00063.x.

Terazawa, A. et al. (2003) 'High Toxoplasma antibody prevalence among inhabitants in Jakarta, Indonesia', *Japanese Journal of Infectious Diseases*, 56(3), pp. 107–109.

Villard, O. et al. (2016) 'Serological diagnosis of Toxoplasma gondii infection: Recommendations from the French National Reference Center for Toxoplasmosis', 84, pp. 22–33. doi: 10.1016/j.diarmicrobio.2015.09.009.

Vueba, N. et al. (2020) 'Serological prevalence of toxoplasmosis in pregnant women in Luanda (Angola): Geospatial distribution and its association with socio-demographic and clinical-obstetric determinants', pp. 1–22. doi: 10.1371/journal.pone.0241908.

Wam, E. C. et al. (2016) 'Seroprevalence of Toxoplasma gondii IgG and IgM antibodies and associated risk factors in women of child-bearing age in Njinikom, NW Cameroon', *BMC Research Notes*. BioMed Central, 9(1), pp. 1–8. doi: 10.1186/s13104-016-2206-0.

Wilking, H. *et al.* (2016) 'Prevalence, incidence estimations, and risk factors of Toxoplasma gondii infection in Germany: A representative, cross-sectional, serological study', *Scientific Reports*. Nature Publishing Group, 6(February), pp. 1–9. doi: 10.1038/srep22551.

de Wit, L. A. *et al.* (2019) 'Potential public health benefits from cat eradication on islands', *PLoS Neglected Tropical Diseases*, 13(2), pp. 1–15. doi: 10.1371/journal.pntd.0007040.

Zemene, E. *et al.* (2012) 'Seroprevalence of Toxoplasma gondii and associated risk factors among pregnant women in Jimma town, Southwestern Ethiopia', *BMC Infectious Diseases*, 12, pp. 2–7. doi: 10.1186/1471-2334-12-337.

## DAFTAR LAMPIRAN

### Lampiran 1. Surat Rekomendasi Persetujuan Etik Penelitian



KEMENTERIAN PENDIDIKAN DAN KEBUDAYAAN  
 UNIVERSITAS HASANUDDIN FAKULTAS KEDOKTERAN  
 KOMITE ETIK PENELITIAN KESEHATAN  
 RSPTN UNIVERSITAS HASANUDDIN  
 RSUP Dr. WAHIDIN SUDIROHUSODO MAKASSAR  
 Sekretariat : Lantai 2 Gedung Laboratorium Terpadu  
 JL. PERINTIS KEMERDEKAAN KAMPUS TAMALANREA KM.10 MAKASSAR 90245.  
 Contact Person: dr. Agussalim Bukhari, MMed,PhD, Sp.GK. Telp. 081240850858, 0411 5780103. Fax : 0411-581431



#### **REKOMENDASI PERSETUJUAN ETIK**

Nomor : 444/UN4.6.4.5.31/ PP36/ 2020

Tanggal: 19 Agustus 2020

Dengan ini Menyatakan bahwa Protokol dan Dokumen yang Berhubungan Dengan Protokol berikut ini telah mendapatkan Persetujuan Etik :

No Protokol	UH20070283	No Sponsor Protokol	
Peneliti Utama	<b>dr. Nurul Fadilah</b>	Sponsor	
Judul Peneliti	Identifikasi Infeksi Toxoplasma gondii secara Serologis dan Biologi Molekuler pada Ibu Hamil di Kota Makassar serta Hubungannya dengan Faktor Resiko		
No Versi Protokol	2	Tanggal Versi	19 Agustus 2020
No Versi PSP	2	Tanggal Versi	19 Agustus 2020
Tempat Penelitian	<b>RS Universitas Hasanuddin dan Puskesmas di Kota Makassar</b>		
Jenis Review	<input type="checkbox"/> Exempted <input checked="" type="checkbox"/> Expedited <input type="checkbox"/> Fullboard Tanggal	Masa Berlaku 19 Agustus 2020 sampai 19 Agustus 2021	Frekuensi review lanjutan
Ketua Komisi Etik Penelitian Kesehatan FKUH	Nama <b>Prof.Dr.dr. Suryani As'ad, M.Sc.,Sp.GK (K)</b>	Tanda tangan 	
Sekretaris Komisi Etik Penelitian Kesehatan FKUH	Nama <b>dr. Agussalim Bukhari, M.Med.,Ph.D.,Sp.GK (K)</b>	Tanda tangan 	

Kewajiban Peneliti Utama:

- Menyerahkan Amandemen Protokol untuk persetujuan sebelum di implementasikan
- Menyerahkan Laporan SAE ke Komisi Etik dalam 24 Jam dan dilengkapi dalam 7 hari dan Lapor SUSAR dalam 72 Jam setelah Peneliti Utama menerima laporan
- Menyerahkan Laporan Kemajuan (progress report) setiap 6 bulan untuk penelitian resiko tinggi dan setiap setahun untuk penelitian resiko rendah
- Menyerahkan laporan akhir setelah Penelitian berakhir
- Melaporkan penyimpangan dari protokol yang disetujui (protocol deviation / violation)
- Mematuhi semua peraturan yang ditentukan

**Lampiran 2.** Dokumentasi Penelitian