

## DAFTAR PUSTAKA

- Aiuti, A., Webb, I.J., Bleul, C., Springer, T., and Gutierrez-Ramos, J.C. 1997. The chemokine SDF-1 is a chemoattractant for human CD34+ hematopoietic progenitor cells and provides a new mechanism to explain the mobilization of CD34+ progenitors to peripheral blood. *J. Exp. Med.* 185:111–120.
- Albertsson AM, Zhang X, Vontell R, Bi D, Bronson R, Supramaniam V, et al.  $\gamma\delta$  T Cells Contribute to Injury in The Developing Brain. *The American journal of pathology*. 2018;188(3):757-767.
- Antonelli M, Massimino M, Morra I, Garre ML, Gardiman MP, Buttarelli FR, Arcella A, Giangaspero F: Expression of pERK and pAKT in pediatric high grade astrocytomas: correlation with YKL40 and prognostic significance. *Neuropathology* 2012, 32(2):133-138.
- Badan Penelitian dan Pengembangan. 2014. Laporan Nasional Riset Kesehatan Dasar 2013. Jakarta, P. 136–139.
- Bading H, Ginty DD, Greenberg ME. Regulation of gene expression in hippocampal neurons by distinct calcium signaling pathways. *Science*. 1993 Apr 9;260(5105):181-6. doi: 10.1126/science.8097060. PMID: 8097060.
- Bae S., Kim H., Lee N., Won C., Kim H.R., Hwang Y.I., Song Y.W., Kang J.S., Lee W.J. Alpha-Enolase expressed on the surfaces of monocytes and macrophages induces robust synovial inflammation in rheumatoid arthritis. *J. Immunol.* 2012; 189:365–372. doi: 10.4049/jimmunol.1102073.
- Bajetto, A., Barbero, S., Bonavia, R., Piccioli, P., Pirani, P., Florio, T., et al. (2001). Stromal cell-derived factor-1alpha induces astrocyte proliferation through the activation of extracellular signal-regulated kinases 1/2 pathway. *J. Neurochem.* 77, 1226–1236. doi: 10.1046/j.1471-4159.2001.00350.
- Barbara Viviani, Stefano Bartesaghi, Emanuela Corsini, Pia Villa, Pietro Ghezzi, Angela Garau, Corrado L. Galli and Marina Marinovich: Erythropoietin protects primary hippocampal neurons increasing the expression of brain-derived neurotrophic factor. *Journal of Neurochemistry*, 2005, 93, 412–421. doi:10.1111/j.1471-4159.2005.03033
- Barnathan ES, Raghunath PN, Tomaszewski JE, Ganz T, Cines DB, Higazi A al-R. Immunohistochemical localization of defensin in human coronary vessels. *Am J Pathol* 1997; 150: 1009–20.
- Bathina S, Das UN. Brain-derived neurotrophic factor and its clinical implications. *Arch Med Sci*. 2015;11(6):1164-1178. doi:10.5114/aoms.2015.56342.
- Beauchamp K, et al, Pharmacology of Traumatic Brain Injury: Where is the “golden bullet”? 2008. *Molecular Medicine* 14(11-12):731-40, DOI:10.2119/2008-00050.Beauchamp.
- Begaz T, Kyriacou DN, Segal J, Bazarian JJ (2006) Serum biochemical markers for post-concussion syndrome in patients with mild traumatic brain injury. *J Neurotrauma* 23: 1201–1210.

- Berk, M.; Kapczinski, F.; Andreazza, A.; Dean, O.; Giorlando, F.; Maes, M.; Yücel, M.; Gama, C.; Dodd, S.; Dean, B. Pathways underlying neuroprogression in bipolar disorder: Focus on inflammation, oxidative stress and neurotrophic factors. *Neurosci. Biobehav. Rev.* 2011, 35, 804–817.
- Bernaudin M, Marti HH, Roussel S, Divoux D, Nouvelot A, MacKenzie ET, et al: A potential role for erythropoietin in focal permanent cerebral ischemia in mice. *J Cereb Blood Flow Metab* 19:643–651, 1999.
- Bleul, C.C., Fuhlbrigge, R.C., Casasnovas, J.M., Aiuti, A., and Springer, T.A. 1996. A highly efficacious lymphocyte chemoattractant, stromal cell-derived factor 1 (SDF-1). *J. Exp. Med.* 184:1101–1109.
- Boone, D.R.; Sell, S.L.; Micci, M.A.; Crookshanks, J.M.; Parsley, M.; Uchida, T.; Prough, D.S.; DeWitt, D.S.; Hellmich, H.L. Traumatic brain injury induced dysregulation of the circadian clock. *PLoS ONE* 2012, 7, e46204.
- Bradbury MW. The blood-brain barrier. *Exp Physiol.* 1993;78(4):453–72. 10. Cederberg D, Siesjo P. What has inflammation to do with traumatic brain injury? *Childs Nerv Syst.* 2010;26(2):221–6.
- Bramlett, H. M, Dietrich, W. D. Long-Term Consequences of Traumatic Brain Injury: Current Status of Potential Mechanisms of Injury and Neurological Outcomes', *Journal of Neurotrauma.* 2015; 32(23):1834– 1848.
- Brandes ME, Mai UE, Ohura K, Wahl SM. Type I transforming growth factor-beta receptors on neutrophils mediate chemotaxis to transforming growth factor-beta. *J Immunol.* 1991 Sep 1;147(5):1600-6. PMID: 1652608.
- Brines, M., and A. Cerami. 2006. Discovering erythropoietin's extra-hematopoietic functions: biology and clinical promise. *Kidney Int.* 70:246–250. <http://dx.doi.org/10.1038/sj.ki.5001546>
- Brines, ML, Ghezzi P, Keenan S, Agnello D, de Lanerolle N, Ceramic C, et al: Erythropoietin crosses the blood-brain barrier to protect against experimental brain injury. *Proc Natl Acad Sci U S A* 97:10526–10531, 2000.
- C. Li, C. Shi, J. Kim et al., “Erythropoietin promotes bone formation through EphrinB2/EphB4 signaling,” *Journal of Dental Research*, vol. 94, no. 3, pp. 455–463, 2014. 12 Stem Cells International.
- Cao F., Yang X.F., Liu W.G., Hu W.W., Li G., Zheng X.J., Shen F., Zhao X.Q., Lv S.T. Elevation of neuron-specific enolase and S-100 beta protein level in experimental acute spinal cord injury. *J. Clin. Neurosci.* 2008; 15: 541–544. doi: 10.1016/j.jocn. 2007. 014.
- Carmeliet P: Mechanisms of angiogenesis and arteriogenesis. *Nat Med* 6: 389-395, 2000.
- Carney N, Totten AM, O'Reilly C, Ullman JS, Hawryluk GW, Bell MJ, et al. Guidelines for the management of severe traumatic brain injury. Fourth edition. *Neurosurgery.* 2017;80:6–15.
- Chen S, Zhang L, Yan G, Cheng S, Fathy AH, Yan N, Zhao Y. Neutrophil-to-Lymphocyte Ratio Is a Potential Prognostic Biomarker in Patients with Ovarian Cancer: A Meta-Analysis. *Biomed Res Int.* 2017; 2017: 7943467. doi: 10.1155/2017/7943467. Epub 2017 Jul 26. PMID: 28815182; PMCID: PMC5549495.

- Chen SD, Wu CL, Hwang WC, Yang DI. More Insight into BDNF against Neurodegeneration: Anti-Apoptosis, Anti-Oxidation, and Suppression of Autophagy. *Int J Mol Sci.* 2017;18(3):545. Published 2017 Mar 3. doi:10.3390/ijms18030545.
- Cheng F, Yuan Q, Yang J, Wang W, Liu H. The prognostic value of serum neuron-specific enolase in traumatic brain injury: systematic review and meta-analysis. *PLoS One.* 2014;9(9):e106680. Published 2014 Sep 4. doi:10.1371/journal.pone.0106680.
- Cheng, Q. Yuan, J. Yang, W. Wang, H. Liu, The prognostic value of serum neuron-specific enolase in traumatic brain injury: systematic review and Meta-Analysis, *PLoS One* 9 (2014) e106680.
- Cheng, Xi, et Al. The Role of Sdf-1/Cxcr4/Cxcr7 in Neuronal Regeneration after Cerebral Ischemia. *Frontiers In Neuroscience*, 2017, 11: 590.
- Cherian, J.C. Goodman, C. Robertson, Neuroprotection with erythropoietin administration following controlled cortical impact injury in rats, *J. Pharmacol. Exp. Ther.* 322 (2007) 789–794.
- Chong Yang Ma, et al, Erythropoietin Pathway: A Potential Target for the Treatment of Depression, *Int. J. Mol. Sci.* 2016, 17(5), 677; <https://doi.org/10.3390/ijms17050677>.
- Clark RK, Lee EV, Fish CJ, White RF, Price WJ, Jonak ZL, Feuerstein GZ, Barone FC: Development of tissue damage, inflammation and resolution following stroke: an immunohistochemical and quantitative planimetric study. *Brain Res Bull* 1993, 31:565-572.
- Clausen F, Lorant T, Lewen A, Hillered L: T lymphocyte trafficking: a novel target for neuroprotection in traumatic brain injury. *J Neurotrauma* 2007, 24:1295-1307.
- Criscuolo, C.; Fabiani, C.; Bonadonna, C.; Origlia, N.; Domenici, L. BDNF prevents amyloid-dependent impairment of LTP in the entorhinal cortex by attenuating p38 MAPK phosphorylation. *Neurobiol. Aging* 2015, 36, 1303–1309.
- Dash, Pramod K., et Al. Biomarkers for The Diagnosis, Prognosis, and Evaluation of Treatment Efficacy for Traumatic Brain Injury. *Neurotherapeutics*, 2010, 7.1: 100-114.
- De Silva, Mary J., et Al. Patient Outcome after Traumatic Brain Injury in High-, Middle-And Low-Income Countries: Analysis of Data On 8927 Patients In 46 Countries. *International Journal of Epidemiology*, 2009, 38.2: 452-458.
- Dechant G, Barde YA. The neurotrophin receptor p75(NTR): novel functions and implications for diseases of the nervous system. *Nat Neurosci.* 2002;5:1131–1136.
- Deng, Quan-Jun; Xu, Xiao-Feng; Ren, Jing. Effects Of Sdf-1/Cxcr4 on The Repair of Traumatic Brain Injury in Rats by Mediating Bone Marrow Derived Mesenchymal Stem Cells. *Cellular and Molecular Neurobiology*, 2018, 38.2: 467-477.
- Dewan, M.C., Rattani, A., Gupta, S., Baticulon, R.E., Hung, Y.-C., Punchak, M., Agrawal, A., Adeleye, A.O., Shrime, M.G., Rubiano, A.M., Rosenfeld, J.V., Park, K.B., 2018. Estimating the global incidence of traumatic brain injury. *J. Neurosurg.* 1–18. <https://doi.org/10.3171/2017.10.JNS17352>.
- Dharmajaya, Ridha. Brain Derived Neurotrophic Factor (BDNF) Pasca cedera Kepala Berat sebagai Faktor Peramalan Perjalanan Penyakit {(Brain Derived Neurotrophic Factor (BDNF)} As A Prognostic Factor in Severe Head Injury)}. *Indonesian Journal of Clinical Pathology and Medical Laboratory*, 2018, 21.1: 61-66.

- Dzietko M., Felderhoff-Mueser U., Siffringer M., Krutz B., Bittigau P., Thor F., Heumann R., Buhrer C., Ikonomidou C. and Hansen H. H. (2004) Erythropoietin protects the developing brain against N-methyl-D-aspartate receptor antagonist neurotoxicity. *Neurobiol. Dis.* 15, 177–187.
- Egea-Guerrero J.J., Murillo-Cabezas F., Rodriguez-Rodriguez A., Gordillo-Escobar E., Revuelto-Rey J., Munoz-Sanchez M.A., Leon-Justel A., Vilches-Arenas A. (An experimental model of mass-type brain damage in the rat: Expression of brain damage based on neurospecific enolase and protein S100B) *Med. Intensiv.* 2014; 38:218–225. doi: 10.1016/j.medint.2013.03.015.
- Ehrenreich, H., Aust, C., Krampe, H. et al. Erythropoietin: Novel Approaches to Neuroprotection in Human Brain Disease. *Metab Brain Dis* 19, 195–206 (2004). <https://doi.org/10.1023/B:MEBR.0000043969.96895.3c>
- Eldadah BA, Faden AI. Caspase pathways, neuronal apoptosis, and CNS injury. *J Neurotrauma.* 2000; 17(10): 811–29.
- EllinorKenne. et al. Neutrophil depletion reduces edema formation and tissue loss following traumatic brain injury in mice. *Journal of Neuroinflammation* 2012, 9:17. <http://www.jneuroinflammation.com/content/9/1/17>.
- Elliott, E.; Atlas, R.; Lange, A.; Ginzburg, I. Brain-derived neurotrophic factor induces a rapid dephosphorylation of tau protein through a PI-3 Kinase signalling mechanism. *Eur. J. Neurosci.* 2005, 22, 1081–1089.
- Elson, D.A. et al. Induction of hypervascularity without leakage or inflammation in transgenic mice overexpressing hypoxia-inducible factor-1 $\alpha$ . *Genes Dev.* 15, 2520–2532 (2001).
- Ergun R., Bostancı U., Akdemir G., Beskonaklı E., Kaptanoglu E., Gursoy F., Taskin Y. Prognostic value of serum neuron-specific enolase levels after head injury. *Neurol. Res.* 1998;20: 418–420. doi: 10.1080/01616412.1998.11740541.
- Faila MD, Conley,YP, Wagner AK. BDNF in Traumatic Brain Injury-Related Mortality: Interrelationships Between Genetics and Acute Systemic Central Nervous System BDNF Profiles. Sage Journals, 2015, <https://doi.org/10.1177/1545968315586465>.
- Fisher Jw. Erythropoietin: Physiology and Pharmacology Update. Society For Experimental Biology and Medicine, 2003; 216: 358–69.
- Fleitas, C.; Piñol-Ripoll, G.; Marfull, P.; Rocandio, D.; Ferrer, I.; Rampon, C.; Egea, J.; Espinet, C. proBDNF is modified by advanced glycation end products in Alzheimer's disease and causes neuronal apoptosis by inducing p75 neurotrophin receptor processing. *Mol. Brain* 2018, 11, 68.
- Franklin J. L. and Johnson E. M. Jr (1992) Suppression of programmed neuronal death by sustained elevation of cytoplasmic calcium. *Trends Neurosci.* 15, 501–508.
- Frisen J, et al. Characterization of glial trkB receptors: differential response to injury in the central and peripheral nervous systems. *Proc Natl Acad Sci USA.* 1993;90:4971–4975.
- Fuadi, Iwan; Bisri, Tatang. Efek Proteksi Otak Erythropoietin. *Jurnal Neuroanestesi Indonesia,* 2015, 4.2: 149-56.
- Galgano M, Toshkezi G, Qiu X, Russell T, Chin L, Zhao LR. Traumatic Brain Injury: Current Treatment Strategies and Future Endeavors. *Cell Transplant.* 2017;26(7):1118-1130. doi:10.1177/0963689717714102.

- Ganju RK, Brubaker SA, Meyer J, Dutt P, Yang Y, Qin S, Newman W and Groopman JE: The  $\alpha$ -chemokine, stromal cell-derived factor-1alpha, binds to the transmembrane G-protein-coupled CXCR-4 receptor and activates multiple signal transduction pathways. *J Biol Chem* 273: 23169-23175, 1998.
- Ghajar, Jamshid. Traumatic Brain Injury. *The Lancet*, 2000, 356.9233: 923-929.
- Giulian D, Chen J, Ingeman JE, George JK, Noponen M. The role of mononuclear phagocytes in wound healing after traumatic injury to adult mammalian brain. *J Neurosci*. 1989;9(12): 4416–29. Dong et al. *Journal of Neuroinflammation* (2016) 13:197 Page 12 of 13.
- Gonzalez FF, McQuillen P, Mu D, Chang Y, Wendland M, Vexler Z, Ferriero DM. Erythropoietin enhances long-term neuroprotection and neurogenesis in neonatal stroke. *Dev Neurosci* 2007;29(4–5):321–330.
- Gradisek P, Osredkar J, Korsic M, Kremzar B. Multiple indicators model of long-term mortality in traumatic brain injury. *Brain Inj.* 2012;26(12):1472-81. doi: 10.3109/02699052.2012.694567. Epub 2012 Jun 21. PMID: 22721420.
- Grasso G, Sfacteria A, Cerami A, Brines M. Erythropoietin as a tissue- protective cytokine in brain injury: what do we know and where do we go? *Neuroscientist* 2004;10(2):93–98.
- Greve, Mark W.; Zink, Brian J. Pathophysiology of Traumatic Brain Injury. *Mount Sinai Journal of Medicine: A Journal of Translational and Personalized Medicine: A Journal of Translational and Personalized Medicine*, 2009, 76.2: 97-104.
- Guerriero Rm, Giza Cc, Rotenberg A. Glutamate and Gaba Imbalance Following Traumatic Brain Injury. *CurrNeurolNeurosci*. 2015;15(5):1-20.
- Guo R, Chai LL, Chen L, Chen WG, Ge LP, Li XG, Li HL, Li SR, Cao C. Stromal cell-derived factor 1 (SDF-1) accelerated skin wound healing by promoting the migration and proliferation of epidermal stem cells. *In Vitro Cell Dev Biol Anim*. 2015;51(6):578–585.
- Gururaj G., An epidemiological approach to prevention – prehospital care and rehabilitation in neurotrauma, *Neurology India* 43(3) (1995), 95–105.
- Gustafsson, D.; Klang, A.; Thams, S.; Rostami, E. The Role of BDNF in Experimental and Clinical Traumatic Brain Injury. *Int. J. Mol. Sci.* 2021, 22, 3582. <https://doi.org/10.3390/ijms22073582>.
- Haase VH: Hypoxic regulation of erythropoiesis and iron metabolism. *Am J Physiol Renal Physiol* 2010;299: F1–F13.
- Hafner A., Glavan G., Obermajer N., Zivin M., Schliebs R., Kos J. Neuroprotective role of gamma-enolase in microglia in a mouse model of Alzheimer's disease is regulated by cathepsin X. *Aging Cell*. 2013; 12:604–614. doi: 10.1111/acel.12093.
- Haiga, Yuri, Amir, Darwin, Syafrita, Yuliarni. Association Between Spesific Enolase Serum Levels and Outcome Acute Ischemic Stroke Onset 1-Month. *Indonesian Journal of Clinical Pathology and Medical Laboratory*, 2019, 25.2: 135-139.
- Haque A, Ray SK, Cox A, Banik NL. Neuron specific enolase: a promising therapeutic target in acute spinal cord injury. *Metab Brain Dis*. 2016 Jun;31(3): 487-95. doi: 10.1007/s11011-016-9801-6. Epub 2016 Feb 5. PMID: 26847611; PMCID: PMC4864119.

- Hazeldine J, Lord JM, Belli A. Traumatic Brain Injury and Peripheral Immune Suppression: Primer and Prospectus. *Front Neurol.* 2015 Nov 5;6:235. doi: 10.3389/fneur.2015.00235. PMID: 26594196; PMCID: PMC4633482.
- Hellewell SC, Yan EB, Alwis DS, et al. Erythropoietin improves motor and cognitive deficit, axonal pathology, and neuroinflammation in a combined model of diffuse traumatic brain injury and hypoxia, in association with upregulation of the erythropoietin receptor. *J Neuroinflammation.* 2013; 10:156.
- Hernández, Carolina Castillo, et al. Neuroprotective Effects of Erythropoietin on Neurodegenerative and Ischemic Brain Diseases: The Role of Erythropoietin Receptor. *Neural Regeneration Research,* 2017, 12.9: 1381.
- Hikmi, Syauq, et al. Hubungan antara Hasil Pemeriksaan Awal Laboratorium Klinik dengan Glasgow Outcome Scale Extended (Gose) pada Penderita Cedera Otak Berat. 2018. Phd Thesis. Universitas Airlangga.
- Hirota, Kiichi & Semenza, Gregg. (2006). Hirota K, Semenza GL.. Regulation of angiogenesis by hypoxia-inducible factor 1. *Crit Rev Oncol Hematol* 59: 15-26. Critical reviews in oncology/hematology. 59. 15-26. 10.1016/j.critrevonc.2005.12.003.
- Hofman K, Primack A, Keusch, G, Hrynkow S. Addressing the Growing Burden of Trauma and Injury in Low-And Middle-Income Country. 2005. *American Journal of Public Health,* Vol. 95, No. 1, Pp. 13–17.
- Holmin S, Mathiesen T, Shetye J, Biberfeld P. Intracerebral inflammatory response to experimental brain contusion. *Acta Neurochir (Wien).* 1995;132(1-3):110–9.
- Huang EJ, Reichardt LF. Neurotrophins: roles in neuronal development and function. *Annu Rev Neurosci.* 2001;24:677-736. doi: 10.1146/annurev.neuro.24.1.677. PMID: 11520916; PMCID: PMC2758233.
- Huang, Chun-Ta, et al. Erythropoietin Reduces Nerve Demyelination, Neuropathic Pain Behavior and Microglial MAPK S Activation Through Erythropoietin Receptors on Schwann Cells in A Rat Model of Peripheral Neuropathy. *Glia,* 2018, 66.11: 2299-2315.
- Hyder AA, Wunderlich CA, Puvanachandra P, Gururaj G, Kobusingye OC. The impact of traumatic brain injuries: a global perspective. *NeuroRehabilitation.* 2007;22(5):341-53. PMID: 18162698.
- Itoh, Tatsuki, et al. The Relationship Between Sdf-1 A/Cxcr4 and Neural Stem Cells Appearing in Damaged Area After Traumatic Brain Injury in Rats. *Neurological Research,* 2009, 31.1: 90-102.
- J. M. Hallenbeck, A. J. Dutka, T. Tanishima, P. M. Kochanek, K. K. Kumaroo, C. B. Thompson, T. P. Obrenovitch, T. J. Contreras, Polymorphonuclear leukocyte accumulation in brain regions with low blood flow during the early post ischemic period. *Stroke* 17, 246–253 (1986).
- Jacobs, K., Shoemaker, C., Rudersdorf, R., Neill, S.D., Kaufmann, R.J., Mufson, A., Seehra, J., Jones, S.S., Hewick, R., And Fritsch, E.F. (1985). Isolation and Characterization of Genomic and Cdna Clones of Human Erythropoietin. *Nature.* 313(6005): 806-810. Pmid:3838366.
- Jaquet K, Krause K, Tawakol-Khodai M, Geidel S, Kuck KH. Erythropoietin and VEGF exhibit equal angiogenic potential. *Microvasc Res.* 2002 Sep;64(2):326-33. doi: 10.1006/mvre.2002.2426. PMID: 12204656.

- Jelkmann, W. (2013). Physiology And Pharmacology of Erythropoietin. *Transfus Med Hemother.* 40(5):302–309 Doi: 10.1159/000356193.
- Jennifer C. et al. The cellular inflammatory response in human spinal cords after injury, *Brain* (2006), 129, 3249–3269. doi:10.1093/brain/awl296.
- Jin W, Kong J, Lu T, et al. Erythropoietin prevents secondary brain injury induced by cortical lesion in mice: possible involvement of Nrf2 signaling pathway. *Ann Clin Lab Sci.* 2011; 41:25-32.
- Johnsson P., Blomquist S., Luhrs C., Malmkvist G., Alling C., Solem J.O., Stahl E. Neuron-specific enolase increases in plasma during and immediately after extracorporeal circulation. *Ann. Thorac. Surg.* 2000;6 9:750–754. doi: 10.1016/S0003-4975(99)01393-4.
- Juul, Sandra E., et al. A Randomized Trial of Erythropoietin for Neuroprotection in Preterm Infants. *New England Journal of Medicine*, 2020, 382.3: 233-243.
- Kaczorowski SL, Schiding JK, Toth CA, Kochanek PM (1995): Effect of soluble complement receptor-1 on neutrophil accumulation after traumatic brain injury in rats. *J Cerebral Blood Flow Metabolism* 15: 860–864.
- Kastilong M, Subrata I, Tangkudung G, Herlyani Khosama. Neutrophyl lymphocyte ratio and the outcome of traumatic Brain injury. *Jurnal Sinaps*, Vol. 1 No. 2(2018), Im. 20-28.
- Kellert, B., McPherson RJ., Juul, SE. A Comparison of High-Dose Recombinant Erythropoietin Treatment Regimens in Brain-Injured Neonatal Rats, *Pediatric Research* 61, 2007, 451-455.
- Khan M, Im YB, Shunmugavel A, Gilg AG, Dhindsa RK, Singh AK, Singh I: Administration of S-nitrosoglutathione after traumatic brain injury protects the neurovascular unit and reduces secondary injury in a rat model of controlled cortical impact. *J Neuroinflammation* 2009, 6:32.
- Klein R, Nanduri V, Jing SA, et al. The trkB tyrosine protein kinase is a receptor for brain-derived neurotrophic factor and neurotrophin-3. *Cell.* 1991; 66(2): 395-403. doi: 10.1016/0092-8674(91)90628-c
- Kleine T.O., Benes L., Zofel P. Studies of the brain specificity of S100B and neuron-specific enolase (NSE) in blood serum of acute care patients. *Brain Res. Bull.* 2003; 61:265–279. doi: 10.1016/S0361-9230(03)00090-X.
- Kochanek, Pm, Berger, Rp, Fink, El, Au, Ak, Bayır, H, Bell, Mj, Dixon, Ce, & Clark, Rsb 2013, The Potential for Bio-Mediators and Biomarkers in Pediatric Traumatic Brain Injury and Neurocritical Care, *Front Neurol*, Vol.4, P1-9.
- Konrad FM, Braun S, Ngamsri KC, Vollmer I, Reutershan J. Heme oxygenase-1 attenuates acute pulmonary inflammation by decreasing the release of segmented neutrophils from the bone marrow. *Am J Physiol Lung Cell Mol Physiol* 2014; 307: L707–L717.
- Korhonen L, Riikonen R, Nawa H, Lindholm D. Brain Derived Neurotrophic Factor is Increased in Cerebrospinal Fluid of Children Suffering from Asphyxia. *Neuroscience Letters* 1998; 240: 151–154.
- Korley, F.K.; Diaz-Arrastia, R.; Wu, A.H.; Yue, J.K.; Manley, G.T.; Sair, H.I.; Van Eyk, J.; Everett, A.D; Track-Tbi Investigators; Okonkwo, D.O.; et al. Circulating Brain-Derived Neurotrophic Factor has Diagnostic and Prognostic Value in Traumatic Brain Injury. *J. Neurotrauma* 2016, 33, 215–225.

- Korte M, Carroll P, Wolf E, Brem G, Thoenen H, Bonhoeffer T. Hippocampal long-term potentiation is impaired in mice lacking brain-derived neurotrophic factor. *Proc Natl Acad Sci U S A*. 1995 Sep 12;92(19):8856-60. doi: 10.1073/pnas.92.19.8856. PMID: 7568031; PMCID: PMC41066.
- Kremer KN, Peterson KL, Schneider PA, Meng XW, Dai H, Hess AD, et al. CXCR4 chemokine receptor signaling induces apoptosis in acute myeloid leukemia cells via regulation of the Bcl-2 family members Bcl-XL, Noxa, and Bak. *J Biol Chem*. 2013; 288(32):22899–914.
- Krueger, F., Pardini, M., Huey, E. D., Raymont, V., Solomon, J., Lipsky, R. H., et al. (2011). The role of the Met66 brain-derived neurotrophic factor allele in the recovery of executive functioning after combat-related traumatic brain injury. *J. Neurosci.* 31, 598–606. doi: 10.1523/JNEUROSCI.1399-10.2011.
- La Ferla K, Reimann C, Jelkmann W, HellwigBürgel T: Inhibition of erythropoietin gene expression signaling involves the transcription factors GATA-2 and NF- $\kappa$ B. *FASEB J* 2002;16: 1811–1813.
- Lambeth JD. Nox enzymes. ROS, and chronic disease: an example of antagonistic pleiotropy. *Free Radic Biol Med* 2007; 43:332–47.
- Lappin, TR, The Cellular Biology of Erythropoietin Receptors. 2013, *The oncologist* 8 Suppl 1 (90001):15-8, DOI: 10.1634/theoncologist.8-suppl\_1-15.
- Lee KS. Estimation of the incidence of head injury in Korea: an approximation based on national traffic accident statistics. *J Korean Med Sci*. 2001;16:342–346.
- Leibrock J, Lottspeich F, Hohn A, Hofer M, Hengerer B, Masiakowski P, Thoenen H, Barde YA. Molecular cloning and expression of brain-derived neurotrophic factor. *Nature*. 1989 Sep 14;341(6238):149-52. doi: 10.1038/341149a0. PMID: 2779653.
- Levin HS, Shum D, Chan RC. Understanding traumatic brain injury: current research and future directions. New York (NY): Oxford University Press; 2014.
- Li, J., Guo, W., Xiong, M., Han, H., Chen, J., Mao, D., Zeng, Y. (2015). Effect of SDF-1/CXCR4 axis on the migration of transplanted bone mesenchymal stem cells mobilized by erythropoietin toward lesion sites following spinal cord injury. *International Journal of Molecular Medicine*, 36, 1205–1214.
- Liao ZB, Jiang GY, Tang ZH, et al. Erythropoietin can promote survival of cerebral cells by downregulating Bax gene after traumatic brain injury in rats. *Neurol India*. 2009; 57:722-728.
- Lima, Je, Takayanagui, Om, Garcia, Lv, & Danleite, Jp 2004, ‘Use of Neuron-Specific Enolase for Assessing the Severity and Outcome in Patients with Neurological Disorders’, *Brazilian Journal of Medical and Biological Research*, Vol.37, P19-26.
- Liu SC, Alomran R, Chernikova SB, Lartey F, Stafford J, Jang T, Merchant M, Zboralski D, Zollner S, Kruschinski et al: Blockade of SDF-1 after irradiation inhibits tumor recurrences of autochthonous brain tumors in rats. *Neuro Oncol* 2014, 16(1):21-28.
- Liu XB, Wang JA, Yu SP, Keogh CL, Wei L. Therapeutic strategy of erythropoietin in neurological disorders. *CNS NeurolDisord Drug Targets* 2008;7(3):227–234. [PubMed: 18673207]
- Liu, Motao, Et Al. Efficacy and safety of erythropoietin for traumatic brain injury. *BMC Neurology* 20, Article number: 399, 2020.

- Liu, Wenchao, Et Al. Neuroprotective Effect of Erythropoietin in Treatment of Traumatic Brain Injury in Rats. *Chinese Journal of Trauma*, 2017, 33.11: 1041-1048.
- Liu, X.; Duan, B.; Cheng, Z.; Jia, X.; Mao, L.; Fu, H.; Kong, D. SDF-1/CXCR4 axis modulates bone marrow mesenchymal stem cell apoptosis, migration and cytokine secretion. *Protein Cell* 2011, 2, 845–854.
- Loetscher, M., T. Geiser, T. O'Reilly, R. Zwahlen, M. Baggio, and B. Moser. 1994. Cloning of a human seven-transmembrane domain receptor, LESTR, that is highly expressed in leukocytes. *J. Biol. Chem.* 269:232–237.
- Luo, L.; Liu, X.-L.; Li, J.; Mu, R.-H.; Liu, Q.; Yi, L.-T.; Geng, D. Macranthol promotes hippocampal neuronal proliferation in mice via BDNF-TrkB-PI3K/Akt signaling pathway. *Eur. J. Pharmacol.* 2015, 762, 357–363.
- Ma C, Cheng F, Wang X, et al. Erythropoietin Pathway: A Potential Target for the Treatment of Depression. *Int J Mol Sci.* 2016;17(5):677. Published 2016 May 6. doi:10.3390/ijms17050677.
- Maan, Zeshaan N., Et Al. Stromal Cell-Derived Factor 1 (Sdf-1) Signaling and Tissue Homeostasis. In: *Regenerative Medicine and Plastic Surgery*. Springer, Cham, 2019. P. 47-59.
- McDonald, SJ, et al, The effect of concomitant peripheral injury on traumatic brain injury pathobiology and outcome. *Journal of Neuroinflammation* 13, Article number: 90, 2016.
- Maekawa T, Ishii T. Chemokine/receptor dynamics in the regulation of hematopoiesis. *Intern Med.* 2000 Feb;39(2):90-100. doi: 10.2169/internalmedicine.39.90. PMID: 10732823.
- Mahmood A, Lu D, Qu C, Goussev A, Zhang ZG, Lu C, Chopp M. Treatment of traumatic brain injury in rats with erythropoietin and carbamylated erythropoietin. *J Neurosurg* 2007; 107 (2): 392–397. doi: 10.3171/JNS-07/08/0392.
- Mallet, Robert T.; Ryou, Myoung-Gwi. Erythropoietin: Endogenous Protection of Ischemic Brain. In: *Vitamins and Hormones*. Academic Press, 2017. P. 197-232.
- Mao, X.-Y.; Cao, Y.-G.; Ji, Z.; Zhou, H.-H.; Liu, Z.-Q.; Sun, H.-L. Topiramate protects against glutamate excitotoxicity via activating BDNF/TrkB-dependent ERK pathway in rodent hippocampal neurons. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* 2015, 60, 11–17.
- Marangos, P.J.; Schmeichel, D.E.; Parma, A.M.; Goodwin, F.K. Developmental profile of neuron-specific (NSE) and non-neuronal (NNE) enolase. *Brain Res.* 1980, 190, 185–193.
- Marti HH. Erythropoietin and the hypoxic brain. *J Exp Biol* 2004;207(Pt 18):3233–3242. [PubMed: 15299044]
- McGee S., Havens A., Shiozawa Y., Jung Y., Taichman R. Effects of erythropoietin on the bone microenvironment. *Growth Factors*. 2012; 30:22–28.
- Meng, Y. Xiong, A. Mahmood, Y. Zhang, C. Qu, M. Chopp, Dose-dependent neurorestorative effects of delayed treatment of traumatic brain injury with recombinant human erythropoietin in rats, *J. Neurosurg.* 115 (2011) 550–560.
- Mercier, Eric, Et Al. Prognostic Value of Neuron-Specific Enolase (NSE) For Prediction of Post-Concussion Symptoms Following a Mild Traumatic Brain Injury: A Systematic Review. *Brain Injury*, 2018, 32.1: 29-40.
- Meric, Emine, et Al. The Prognostic Value of Neuron-Specific Enolase in Head Trauma Patients. *The Journal of Emergency Medicine*, 2010, 38.3: 297-301.

- Merrick, C. Advanced Trauma Life Support. 10th Edn, Anaesthesia. 10th Edn. Chicago: American College of Surgeons. 2018.
- Morganti-Kossman MC, Lenzlinger PM, Hans V, Stahel P, Csuka E, Ammann E, et al. Production of cytokines following brain injury: beneficial and deleterious for the damaged tissue. *Mol Psychiatry*. 1997;2(2):133–6.
- Morishita E., Masuda S., Nagao M., Yasuda Y. and Sasaki R. (1997) Erythropoietin receptor is expressed in rat hippocampal and cerebral cortical neurons, and erythropoietin prevents in vitro glutamate-induced neuronal death. *Neuroscience* 76, 105–116.
- Moxon-Emre I, Schlichter LC: Neutrophil depletion reduces blood-brain barrier breakdown, axon injury, and inflammation after intracerebral hemorrhage. *J NeuropatholExpNeurol* 2011, 70:218-235.
- Murthy, T. V. S. P., Et Al. Secondary Brain Injury: Prevention and Intensive Care Management. *Indian Journal of Neurotrauma*, 2005, 2.01: 7-12.
- Nagasawa, T., et al. 1996. Defects of B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the CXC chemokine PBSF/SDF-1. *Nature*.382:635–638.
- Naranda T, Kaufman RI, Li J, Wong K, Boge A, Hallen D, Fung KY, Duncan MW, Andersen N, Goldstein A, Olsson L. Activation of erythropoietin receptor through a novel extracellular binding site. *Endocrinology* 2002;143(6): 2293–2302. [PubMed: 12021194]
- Nasution, et al. Modification of the Marmarou model in developing countries. *Annals of Medicine and Surgery*, 57, 109-113, 2020.
- Nekoui, Alireza; Blaise, Gilbert. Erythropoietin and Non-Hematopoietic Effects. *The American Journal of The Medical Sciences*, 2017, 353.1: 76-81.
- Ng, S. Y. And Lee, A. Y. W. Traumatic Brain Injuries: Pathophysiology and Potential Therapeutic Targets', *Frontiers in Cellular Neuroscience*. Frontiers Media S.A. 2019.
- Notopoero, P. B. Eritropoetin Fisiologi, Aspek Klinik, dan Laboratorik. *Indonesian Journal of Clinical Pathology and Medical Laboratory*, 2018, 14.1: 28-36.
- Ogilvie M., Yu X., Nicolas-Metral V., Pulido S.M., Liu C., Ruegg U.T., Noguchi C.T. Erythropoietin stimulates proliferation and interferes with differentiation of myoblasts. *J. Biol. Chem.* 2000;275: 39754–39761.
- Orlando, Ilaria Mc, Et Al. Distal and Proximal Hypoxia Response Elements Cooperate to Regulate Organ-Specific Erythropoietin Gene Expression. *Haematologica*, 2019.
- Ostrowski, Daniela; Heinrich, Ralf. Alternative Erythropoietin Receptors in The Nervous System. *Journal Of Clinical Medicine*, 2018, 7.2: 24.
- Pillai A, Terry AV Jr, Mahadik SP. Differential effects of long-term treatment with typical and atypical antipsychotics on NGF and BDNF levels in rat striatum and hippocampus. *Schizophr Res.* 2006 Feb 15;82(1):95-106. doi: 10.1016/j.schres.2005.11.021. Epub 2006 Jan 25. PMID: 16442781.
- Pillarisetti, K.; Gupta, S.K. Cloning and relative expression analysis of rat stromal cell derived factor-1 (SDF-1)1: SDF-1 alpha mRNA is selectively induced in rat model of myocardial infarction. *Inflammation* 2001, 25, 293–300.

- Pislar A., Bozic B., Zidar N., Kos J. Inhibition of cathepsin X reduces the strength of microglial-mediated neuroinflammation. *Neuropharmacology*. 2017; 114:88–100. doi: 10.1016/j.neuropharm.2016.11.019.
- Polcyn, R.; Capone, M.; Hossain, A.; Matzelle, D.; Banik, N.L.; Haque, A. Neuron specific enolase is a potential target for regulating neuronal cell survival and death: Implications in neurodegeneration and regeneration. *Neuroimmunol. Neuroinflamm.* 2017, 4, 254–257.
- Ponomaryov T, Peled A, Petit I, et al. Induction of the chemokine stromal-derived factor-1 following DNA damage improves human stem cell function. *J Clin Invest.* 2000;106(11):1331-1339. doi:10.1172/JCI10329.
- Povlishock JT, Katz DI. Update of neuropathology and neurological recovery after traumatic brain injury. *J Head Trauma Rehabil.* 2005;20(1):76–94.
- Prins, Mayumi, et Al. The Pathophysiology of Traumatic Brain Injury at A Glance. *Disease Models & Mechanisms*, 2013, 6.6: 1307-1315.
- Qin C, Zhou S, Xiao Y, Chen L. Erythropoietin enhances mitochondrial biogenesis in cardiomyocytes exposed to chronic hypoxia through Akt/eNOS signalling pathway. *Cell Biol Int.* 2014; 38:335–342.
- R. F. Wynn, C. A. Hart, C. Corradi-Perini et al., “A small proportion of mesenchymal stem cells strongly expresses functionally active CXCR4 receptor capable of promoting migration to bone marrow, *Blood*, vol. 104, no. 9, pp. 2643 – 2645, 2004.
- R. S. Clark, J. K. Schiding, S. L. Kaczorowski, D. W. Marion, P. M. Kochanek, Neutrophil accumulation after traumatic brain injury in rats: Comparison of weight drop and controlled cortical impact models. *J. Neurotrauma* 11, 499–506 (1994).
- Ramont L, Thoannes H, Volondat A, Chastang F, Millet MC, Maquart FX (2005) Effects of hemolysis and storage condition on neuron-specific enolase (NSE) in cerebrospinal fluid and serum: implications in clinical practice. *Clin Chem Lab Med* 43:1215– 1217
- Rivard, A. et al. Age-dependent defect in vascular endothelial growth factor expression is associated with reduced hypoxia-inducible factor 1 activity. *J. Biol. Chem.* 275, 29643–29647 (2000).
- Rohe, M.; Synowitz, M.; Glass, R.; Paul, S.M.; Nykjaer, A.; Willnow, T.E. Brain-derived neurotrophic factor reduces amyloidogenic processing through control of SORLA gene expression. *J. Neurosci.* 2009, 29, 15472–15478.
- Rutland-Brown W, Langlois JA, Thomas KE, Xi YL. Incidence of traumatic brain injury in the United States, 2003. *J Head Trauma Rehabil.* 2006;21(6):544.
- Salehi Z, Mashayekhi F. Brain-Derived Neurotrophic Factor Concentrations in The Cerebrospinal Fluid of Patients with Parkinson’s Disease. *J Of Clin Neuroscience* 2009; 16 (1): 90–93.
- Sawhney S., Hood K., Shaw A., Braithwaite A.W., Stubbs R., Hung N.A., Royds J.A., Slatter T.L. Alpha-enolase is upregulated on the cell surface and responds to plasminogen activation in mice expressing a 133p53 alpha mimic. *PLoS ONE.* 2015;10: e0116270. doi: 10.1371/journal.pone.0116270.
- Schaefer A., Magocsi M. and Marquardt H. (1997). Signalling mechanisms in erythropoiesis: the enigmatic role of calcium. *Cell. Signal.* 9, 483–495.
- Schluesener H, Meyermann R. Neutrophilic defensins penetrate the blood-brain barrier. *J Neurosci Res* 1995; 42: 718–23.

- Schmitt AB, Buss A, Breuer S, Brook GA, Pech K, Martin D, et al. Major histocompatibility complex class II expression by activated microglia caudal to lesions of descending tracts in the human spinal cord is not associated with a T cell response. *Acta Neuropathol (Berl)* 2000; 100: 528–36.
- Schoettle RJ, Kochanek PM, Magargee MJ, Uhl MW, Nemoto EM (1990): Early polymorphonuclear leukocyte accumulation correlates with the development of posttraumatic cerebral edema in rats. *J Neurotrauma* 7: 207–217.
- Schober ME, et al. Developmental traumatic brain injury decreased brain derived neurotrophic factor expression late after injury. *Metab Brain Dis.* 2012.
- Semple BD, Bye N, Ziebell JM, Morganti-Kossmann MC: Deficiency of the chemokine receptor CXCR2 attenuates neutrophil infiltration and cortical damage following closed head injury. *Neurobiol Dis.* 2010, 40:394-403.
- Shih, Hong-Mou; Wu, Chih-Jen; Lin, Shuei-Liong. Physiology And Pathophysiology of Renal Erythropoietin-Producing Cells. *Journal of The Formosan Medical Association*, 2018, 117.11: 955-963.
- Shimizu, E., Hashimoto, K., and Iyo, M. (2004). Ethnic difference of the BDNF 196G/A (val66met) polymorphism frequencies: the possibility to explain ethnic mental traits. *Am. J. Med. Gen. Part B* 126, 122–123. doi: 10.1002/ajmg.b.20118.
- Simon, D.; Nascimento, R.I.; Filho, E.M.; Bencke, J.; Regner, A. Plasma Brain-Derived Neurotrophic Factor Levels after Severe Traumatic Brain Injury. *Brain Inj.* 2016, 30, 23–28.
- Siren A. L., Fratelli M., Brines M. et al. (2001) Erythropoietin prevents neuronal apoptosis after cerebral ischemia and metabolic stress. *Proc. Natl Acad. Sci. USA* 98, 4044–4049
- Smith M. Monitoring intracranial pressure in traumatic brain injury. *Anesth Analg.* 2008;106(1):240–248.
- Soares HD, Hicks RR, Smith D, McIntosh TK: Inflammatory leukocytic recruitment and diffuse neuronal degeneration are separate pathological processes resulting from traumatic brain injury. *J Neurosci* 1995, 15:8223-8233.
- Soehnlein O, Zernecke A, Eriksson EE, Rothfuchs AG, Pham CT, Herwald H, Bidzhekov K, Rottenberg ME, Weber C, Lindbom L: Neutrophil secretion products pave the way for inflammatory monocytes. *Blood* 2008, 112:1461-1471.
- Staller, P. et al. Chemokine receptor CXCR4 downregulated by von Hippel-Lindau tumour suppressor pVHL. *Nature* 425, 307–311 (2003).
- Sun, W., Liu, J., Huan, Y., & Zhang, C. (2013). Intracranial injection of recombinant stromal-derived factor-1 alpha (SDF-1 $\alpha$ ) attenuates traumatic brain injury in rats. *Inflammation Research*, 63(4), 287–297. doi:10.1007/s00011-013-0699-8.
- Suzuki, Norio, et Al. Regulation of Hypoxia-Inducible Gene Expression After HIF Activation. *Experimental Cell Research*, 2017, 356.2: 182-186.
- Tabish A, N.A. Lone, W.M. Afzal and A. Salam, The incidence and severity of injury in children hospitalized for traumatic brain injury in Kashmir, *Injury* 37(5) (2006), 410–415.
- Takeda K, et Al. Brain-Derived Neurotrophic Factor Inhibits Intercellular Adhesion Molecule-1
- Tamura, Tetsuya, et Al. Neuroprotective Erythropoietin Attenuates Microglial Activation, Including Morphological Changes, Phagocytosis, And Cytokine Production. *Brain Research*, 2017, 1662: 65-74.

- Tang, Mary E.; Lobel, Darlene A. Severe Traumatic Brain Injury: Maximizing Outcomes. Mount Sinai Journal of Medicine: A Journal of Translational and Personalized Medicine: A Journal of Translational and Personalized Medicine, 2009, 76.2: 119-128.
- Tao X., Finkbeiner S., Arnold D. B., Shaywitz A. J. and Greenberg M. E. (1998)  $\text{Ca}^{2+}$  influx regulates BDNF transcription by a CREB family transcription factor-dependent mechanism. *Neuron* 20,709–726.
- Teasdale, G. and B. Jennett (1974). "Assessment of coma and impaired consciousness. A practical scale." *Lancet* 2(7872): 81-4.
- Thelin EP, Jeppsson E, Frostell A, Svensson M, Mondello S, Bellander BM, Nelson DW (2016) Utility of neuron-specific enolase in traumatic brain injury; relations to S100B levels, outcome, and extracranial injury severity. *Crit Care* 20:285.
- Tugyan, Kazım, Et Al. Neuroprotective Effect of Erythropoietin on Nandrolone Decanoate-Induced Brain Injury in Rats. *Neuroscience Letters*, 2013, 533: 28-33.
- Tyler WJ, Perrett SP, Pozzo-Miller LD. The role of neurotrophins in neurotransmitter release. *Neuroscientist*. 2002;8:524–531.
- Villa P, Bigini P, Mennini T, Agnello D, Laragione T, Cagnotto A, Viviani B, Marinovich M, Cerami A, Coleman TR, Brines M, Ghezzi P. Erythropoietin selectively attenuates cytokine production and inflammation in cerebral ischemia by targeting neuronal apoptosis. *J Exp Med*. 2003 Sep 15;198(6):971-5. doi: 10.1084/jem.20021067.
- Wagner AK, McCullough EH, Niyonkuru C, Ozawa H, Loucks TL, et al. Acute serum hormone levels: Characterization and prognosis after severe traumatic brain injury. *J Neurotrauma*. 2011;28(6):871–88.
- Wang, Kevin K., Et Al. An Update on Diagnostic and Prognostic Biomarkers for Traumatic Brain Injury. *Expert Review of Molecular Diagnostics*, 2018, 18.2: 165-180.
- Wang, Y.; Hameed, M.Q.; Rakhade, S.N.; Iglesias, A.H.; Muller, P.A.; Mou, D.L.; Rotenberg, A. Hippocampal immediate early gene transcription in the rat fluid percussion traumatic brain injury model. *Neuroreport* 2014, 25, 954–959.
- Werner C, Engelhard K. Pathophysiology of traumatic brain injury. *Br J Anaesth*. 2007 Jul;99(1):4-9. doi: 10.1093/bja/aem131. PMID: 17573392.
- Whalen MJ, Carlos TM, Kochanek PM, Clark RS, Heineman S, Schiding JK, Franicola D, Memarzadeh F, Lo W, Marion DW, Dekosky ST: Neutrophils do not mediate blood-brain barrier permeability early after controlled cortical impact in rats. *J Neurotrauma* 1999, 16:583-594.
- Wilson EH, Weninger W, Hunter CA. Trafficking of immune cells in the central nervous system. *J Clin Invest*. 2010 May;120(5):1368-79. doi: 10.1172/JCI41911. Epub 2010 May 3. PMID: 20440079; PMCID: PMC2860945.
- Wojchowski DM, Gregory RC, Miller CP, Pandit AK, Pircher TJ. Signal transduction in the erythropoietin receptor system. *Exp Cell Res*. 1999; 253:143–156. [PubMed: 10579919]
- Wu HM, Huang SC, Hattori N, Glenn TC, Vespa PM, et al. (2004) Selective metabolic reduction in gray matter acutely following human traumatic brain injury. *J Neurotrauma* 21: 149–161.

- Wurzelmann, M.; Romeika, J.; Sun, D. Therapeutic potential of brain-derived neurotrophic factor (BDNF) and a small molecular mimic of BDNF for traumatic brain injury. *Neural Regen. Res.* 2017, 12, 7–12.
- Xiong Y, Mahmood A, Meng Y, Zhang Y, Qu C, Schallert T, Chopp M. Delayed administration of erythropoietin reducing hippocampal cell loss, enhancing angiogenesis and neurogenesis, and improving functional outcome following traumatic brain injury in rats: Comparison of treatment with single and triple dose. *J Neurosurg.* 2009.
- Xiong Y, Mahmood A, Qu C, et al. Erythropoietin improves histological and functional outcomes after traumatic brain injury in mice in the absence of the neural erythropoietin receptor. *J Neurotrauma.* 2010; 27:205-215.
- Yang, K.; Perez-Polo, J.R.; Mu, X.S.; Yan, H.Q.; Xue, J.J.; Iwamoto, Y.; Liu, S.J.; Dixon, C.E.; Hayes, R.L. Increased expression of brain-derived neurotrophic factor but not neurotrophin-3 mRNA in rat brain after cortical impact injury. *J. Neurosci. Res.* 1996, 44, 157–164.
- Yanuartono, Yanuartono, et Al. Recombinant Human Erythropoietin: Manfaat dalam Bidang Kedokteran. *Jurnal Sain Veteriner*, 2019, 37.1: 49-60.
- Young Jae Kim, Yong-Wook Jung. Systemic injection of recombinant human erythropoietin after focal cerebral ischemia enhances oligodendroglial and endothelial progenitor cells in rat brain. 2010. *Anatomy and cell biology ACB*, doi:10.5115/acb.2010.43.2.140.
- Yu, Xiaobing, et Al. Erythropoietin Receptor Signalling Is Required for Normal Brain Development. *Development*, 2002, 129.2: 505-516.
- Zafra F., Hengerer B., Leibrock J., Thoenen H. and Lindholm D. (1990). Activity dependent regulation of BDNF and NGF mRNAs in the rat hippocampus is mediated by non-NMDA glutamate receptors. *EMBO J.* 9, 3545–3550.
- Zhang Y., Wang L., Dey S., Alnaeeli M., Suresh S., Rogers H., Teng R., Noguchi C.T. Erythropoietin action in stress response, tissue maintenance and metabolism. *Int. J. Mol. Sci.* 2014;15: 10296–10333.
- Zhang, Xiao Qian, et Al. Increased Protein Expression Levels of P CREB, BDNF and SDF-1/Cxcr4 in The Hippocampus May Be Associated with Enhanced Neurogenesis Induced by Environmental Enrichment. *Molecular Medicine Reports*, 2016, 14.3: 2231-2237.
- Zheng, J.; Liang, J.; Deng, X.; Chen, X.; Wu, F.; Zhao, X.; Luo, Y.; Fu, L.; Jiang, Z. Mitogen activated protein kinase signaling pathways participate in the active principle region of Buyang Huan wu decoction-induced differentiation of bone marrow mesenchymal stem cells. *Neural Regen. Res.* 2012, 7, 1370–1377. [PubMed]
- Zhu ZF, Wang QG, Han BJ, William CP: Neuroprotective effect and cognitive outcome of chronic lithium on traumatic brain injury in mice. *Brain Res Bull* 2010, 83:272-277.

## LAMPIRAN

### 1. Rekomendasi Persetujuan Etik

<p>KEMENTERIAN PENDIDIKAN DAN KEBUDAYAAN UNIVERSITAS HASANUDDIN FAKULTAS KEDOKTERAN KOMITE ETIK PENELITIAN KESEHATAN RSPTN UNIVERSITAS HASANUDDIN RSUP Dr. WAHIDIN SUDIROHUSODO MAKASSAR Sekretariat : Lantai 2 Gedung Laboratorium Terpadu JL.PERINTIS KEMERDEKAAN KAMPUS TAMALANREA KM.10 MAKASSAR 90245. Contact Person: dr. Agussalim Bukhari.,M.Med.,Ph.D.,Sp.GK. TELP. 081241850858, 0411 5780103, Fax : 0411-581431</p> 			
<p><b>REKOMENDASI PERSETUJUAN ETIK</b> Nomor : 408/UN4.6.4.5.31/ PP36/ 2021</p>			
Dengan ini Menyatakan bahwa Protokol dan Dokumen yang Berhubungan Dengan Protokol berikut ini telah mendapatkan Persetujuan Etik :			
No Protokol	UH21050289	No Sponsor Protokol	
Peneliti Utama	dr. Muhammad Fadli Said, SpBS	Sponsor	
Judul Peneliti	Pengaruh Pemberian Eritropoietin terhadap Ekspresi Brain-Derived Neutrophic Factor mRNA (BDNF mRNA), kadar Stromal Cell-Derived Factor-1(SDF-1), dan Neuron Specific Enolase (NSE) pada Tikus Sprague Dawley Model Cedera Otak.		
No Versi Protokol	2	Tanggal Versi	17 Juni 2021
No Versi PSP		Tanggal Versi	
Tempat Penelitian	<b>Laboratorium Animal FKUH dan Laboratorium Mikrobiologi FK Universitas Hasanuddin Makassar</b>		
Jenis Review	<input type="checkbox"/> Exempted <input checked="" type="checkbox"/> Expedited <input type="checkbox"/> Fullboard Tanggal	Masa Berlaku 18 Juni 2021 sampai 18 Juni 2022	Frekuensi review lanjutan
Ketua Komisi Etik Penelitian Kesehatan FKUH	Nama Prof.Dr.dr. Suryani As'ad, M.Sc.,Sp.GK (K)		
Sekretaris Komisi Etik Penelitian Kesehatan FKUH	Nama dr. Agussalim Bukhari, M.Med.,Ph.D.,Sp.GK (K)		
Kewajiban Peneliti Utama:			
<ul style="list-style-type: none"><li>Menyerahkan Amandemen Protokol untuk persetujuan sebelum di implementasikan</li><li>Menyerahkan Laporan SAE ke Komisi Etik dalam 24 Jam dan dilengkapi dalam 7 hari dan Lapor SUSAR dalam 72 Jam setelah Peneliti Utama menerima laporan</li><li>Menyerahkan Laporan Kemajuan (progress report) setiap 6 bulan untuk penelitian resiko tinggi dan setiap setahun untuk penelitian resiko rendah</li><li>Menyerahkan laporan akhir setelah Penelitian berakhir</li><li>Melaporkan penyimpangan dari prokol yang disetujui (protocol deviation / violation)</li><li>Mematuhi semua peraturan yang ditentukan</li></ul>			

## 2. Perlakuan Kelompok Hewan Coba



Penimbangan hewan coba



Eritropoietin yang digunakan



Alat Marmarou



Insisi dan craniotomi tikus



Perlakuan cedera otak



Pengambilan sampel serum



Injeksi eritropoietin



Pengambilan sampel jaringan.

### 3. Pemeriksaan PCR untuk menilai Ekspresi mRNA BDNF



GreenPrimer RTPCR Gen BDNF / Housekeeping Gen GAPDH



PCR Master Mix SYBR



Pembuatan PCR Mix



Nanodrop



Memasukkan template ke dalam PCR Mix



Mesin PCR



Analisa hasil PCR

#### 4. Pemeriksaan ELISA untuk Menilai Kadar SDF-1 dan NSE



Kit Elisa Rat ENO2 dan SDF1



Kit Rat SDF1 Elisa Kit Cat. No. LS-F21608



Menyiapkan semua reagen, standar, dan sampel.

Mengisi sampel dan standar kedalam plate





Inkubasi Plate ke waterbath 37°C selama 90 menit  
lalu kosongkan plate



Isi Biotin ke dalam plate lalu  
inkubasi 1 jam



Buang Biotin lalu Cuci Plate 3X  
kedalam Plate



Mengisi plate dengan HRP Conjugate 1X



Buang Conjugate lalu Cuci Plate 5X



Isi TMB Substrate ke dalam plate lalu inkubasi, standar dan sampel akan berwarna biru





Plate setelah pengisian Substrate.



Stop reaksi dengan penambahan reagen Stop Reaction,  
Sampel dan standar akan berubah dari biru ke kuning.



Plate setelah stop reaksi.



Baca plate ke dalam mesin elisa dengan filter 450 nm



Analisa hasil tes elisa



Mochammad Hatta  
BTN ANTARA B6/6,  
Perintis Kemerdekaan, Km.9  
90245

**OG250521**

Order date : 2021/05/25  
Packing date : 2021/05/29  
Page : 1 / 1

Oligo	BDNF Forward					
SEQ	5'-GAGAAGAGTGATGACCACCT-3' (21mer)					
GC%	MW		Yield		scale (umoles)	Tm°C
	calculated	measured	OD	nmol		
41.7	6,471.8	6,436.8	18.3	76.7	0.2	56.8
vol. for 100pmol/ul	Purification		Modification			
325	MOPC					

Oligo	BDNF Reverse					
SEQ	5'-TCACGTGCTAAAGTGTCAAG-3' (21mer)					
GC%	MW		Yield		scale (umoles)	Tm°C
	calculated	measured	OD	nmol		
41.7	6,511.4	6,388.2	18.4	74.1	0.2	56.8
vol. for 100pmol/ul	Purification		Modification			
331	MOPC					


Oligo	GAPDH Forward					
SEQ	5'-AGGTCGGTGTGAACGGATTG-3' (21mer)					
GC%	MW		Yield		scale (umoles)	Tm°C
	calculated	measured	OD	nmol		
43.4	6,569.2	6,538.2	17.7	72.7	0.2	58.8
vol. for 100pmol/ul	Purification		Modification			
305	MOPC					

Oligo	GAPDH Reverse					
SEQ	5'-GGGGTCGTTGATGGCAACA-3' (19mer)					
GC%	MW		Yield		scale (umoles)	Tm°C
	calculated	measured	OD	nmol		
43.4	6,793.4	6,471.6	17.4	71.9	0.2	58.8
vol. for 100pmol/ul	Purification		Modification			
302	MOPC					

World Meridian 10F, Gasan-dong, Geumcheon-gu, Seoul, 153-781 KOREA TEL +82-2-2113-7363 | FAX +82-2-2113-7120

Sheet Primer Rat BDNF



## Rat SDF1 / CXCL12 ELISA Kit (Sandwich ELISA)

# User Manual

Catalog No. LS-F21608

It is important that you read this entire manual carefully before starting your experiment.

**This kit is for Research Use Only. Not for Diagnostic Use.**  
This kit is not approved for use in humans or for clinical diagnosis.

SDF-1 ELISA KIT



## Rat ENO2 / NSE ELISA Kit (Sandwich ELISA)

# User Manual

Catalog No. LS-F5577

It is important that you read this entire manual carefully before starting your experiment.

**This kit is for Research Use Only. Not for Diagnostic Use.**  
This kit is not approved for use in humans or for clinical diagnosis.

NSE ELISA KIT

**5. Hasil pemeriksaan RTPCR Gen Rat BDNF dengan Housekeeping Gen Rat GAPDH**

**Plate 1**

NO	Sampel	Ekspresi (Fold change)
1	LA01	6,278
2	LA02	6,844
3	LA03	6,040
4	LA04	5,468
5	LA05	5,781
6	LB01	7,357
7	LB02	8,647
8	LB03	7,878
9	LB04	8,140
10	LB05	7,560
11	LC01	8,412
12	LC02	9,821
13	LC03	9,620
14	LC04	9,248
15	LC05	8,907

**Plate 2**

NO	Sampel	Ekspresi (Fold change)
1	LA06	6,326
2	LA07	5,117
3	LA08	5,438
4	LA09	5,812
5	LA10	6,013
6	LA11	10,424
7	LA12	10,376
8	LA13	10,075
9	LA14	9,792
10	LA15	10,197
11	LB06	6,485
12	LB07	6,236
13	LB08	6,216
14	LB09	6,709
15	LB10	6,588
16	LC06	7,220
17	LC07	6,983
18	LC08	7,344
19	LC09	7,095
20	LC10	7,474

**6. Hasil Pemeriksaan Rat SDF-1/CXCL12 ELISA Kit KATALOG LS-F21608**

No	No Sampel	Konsentrasi (ng/ml)
1	SA01	7,984
2	SA02	7,698
3	SA03	8,127
4	SA04	6,860
5	SA05	7,146
6	SA06	6,411
7	SA07	8,290
8	SA08	6,983
9	SA09	7,514
10	SA10	6,656
11	SA11	1,978
12	SA12	2,979
13	SA13	1,712
14	SA14	3,387
15	SA15	2,693
16	SB01	5,798
17	SB02	6,676
18	SB03	6,472
19	SB04	6,309
20	SB05	6,329
21	SB06	9,373
22	SB07	9,005
23	SB08	9,148
24	SB09	9,598
25	SB10	10,456
26	SC01	3,959
27	SC02	4,245
28	SC03	5,389
29	SC04	4,613
30	SC05	5,063
31	SC06	11,763
32	SC07	11,354
33	SC08	13,070
34	SC09	12,662
35	SC10	13,275

**7. Hasil Pemeriksaan Rat ENO2/NSE ELISA KIT Katalog LS-F5577**

No	No Sampel	Konsentrasi (ng/ml)
1	SA01	19,059
2	SA02	17,548
3	SA03	23,759
4	SA04	22,081
5	SA05	20,234
6	SA06	18,471
7	SA07	17,212
8	SA08	24,599
9	SA09	20,990
10	SA10	18,136
11	SA11	3,195
12	SA12	1,432
13	SA13	7,056
14	SA14	4,034
15	SA15	5,629
16	SB01	14,359
17	SB02	16,877
18	SB03	19,563
19	SB04	18,723
20	SB05	16,709
21	SB06	26,445
22	SB07	32,153
23	SB08	30,307
24	SB09	31,230
25	SB10	28,124
26	SC01	10,330
27	SC02	13,519
28	SC03	15,869
29	SC04	11,673
30	SC05	9,322
31	SC06	34,084
32	SC07	36,938
33	SC08	38,952
34	SC09	35,511
35	SC10	41,050

**Tabel x. Instrumen pencatatan berat tikus dan dosis obat yang diberikan.**

Nomor	Kelompok	Berat Badan (gram)	Dosis EPO 30.000 U/kg	Dosis Ketamin 50 mg/kg
1.	A	203		
2.	A	205		
3.	A	209		
4.	A	211		
5.	A	196		
6.	B	210		10.5 mg
7.	B	213		10.65 mg
8.	B	200		10 mg
9.	B	197		9.85 mg
10.	B	199		9.95 mg
11.	C	205	6.150 U	10.25 mg
12.	C	213	6.390 U	10.65 mg
13.	C	198	5.940 U	9.9 mg
14.	C	207	6.210 U	10.35 mg
15.	C	205	6.150 U	10.25 mg

**Kelompok A (1-5)** : Trauma (-), EPO (-)

**Kelompok B (6-10)** : Trauma (+), EPO (-)

**Kelompok C (11-15)** : Trauma (+), EPO (+)