

DAFTAR PUSTAKA

1. NKF-KDIGO. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. ISN. 2013;3(1):1–163.
2. Depkes. InfoDATIN Pusat Data dan Informasi Kementerian Kesehatan RI: Situasi Penyakit Ginjal Kronis. 2017;1–10. Available from: www.depkes.go.id/resources/download/pusdatin/infodatin/
3. Hill NR, Fatoba ST, Oke JL, Hirst JA, O’Callaghan CA, Lasserson DS, et al. Global prevalence of chronic kidney disease - A systematic review and meta-analysis. PLoS One. 2016;11(7):1–18.
4. Bill F, Foundation MG. Global , regional , and national burden of chronic kidney disease , 1990 – 2017 : a systematic analysis for the Global Burden of Disease Study 2017. 2017;
5. Riskesdas. Laporan Riskesdas Tahun 2018 Nasional. 2018;
6. Mardhatillah M, Arsin A, Syafar M, Hardianti A. Ketahanan Hidup Pasien Penyakit Ginjal Kronik Yang Menjalani Hemodialisis Di Rsup Dr. Wahidin Sudirohusodo Makassar. J Kesehat Masy Marit. 2020;3(1):21–33.
7. 11 th Report Of Indonesian Renal Registry. 11th Report Of Indonesian Renal Registry 2018. Irr [Internet]. 2018;1–46. Available from:

[https://www.indonesianrenalregistry.org/data/IRR 2018.pdf](https://www.indonesianrenalregistry.org/data/IRR_2018.pdf)

8. Diaz-Martinez J, Campa A, Delgado-Enciso I, Hain D, George F, Huffman F, et al. The relationship of blood neutrophil-to-lymphocyte ratio with nutrition markers and health outcomes in hemodialysis patients. *Int Urol Nephrol*. 2019;51(7):1239–47.
9. Kwon YE, Kee YK, Yoon CY, Han IM, Han SG, Park KS, et al. Change of nutritional status assessed using subjective global assessment is associated with all-cause mortality in incident dialysis patients. *Med (United States)*. 2016;95(7):e2714.
10. Takahashi H, Ito Y, Ishii H, Aoyama T, Kamoi D, Kasuga H, et al. Geriatric nutritional risk index accurately predicts cardiovascular mortality in incident hemodialysis patients. *J Cardiol [Internet]*. 2014;64(1):32–6. Available from: <http://dx.doi.org/10.1016/j.jjcc.2013.10.018>
11. Lin TY, Hung SC. Geriatric nutritional risk index is associated with unique health conditions and clinical outcomes in chronic kidney disease patients. *Nutrients*. 2019;11(11).
12. Yamada K, Furuya R, Takita T, Maruyama Y, Yamaguchi Y, Ohkawa S. Simplified nutritional screening tools for patients on maintenance hemodialysis 1–3. 2018;(April):106–13.
13. Xiong J, Wang M, Zhang Y, Nie L, He T, Wang Y, et al. Association of geriatric

nutritional risk index with mortality in hemodialysis patients: A meta-analysis of cohort studies. *Kidney Blood Press Res.* 2018;43(6):1878–89.

14. Catabay C, Obi Y, Streja E, Soohoo M, Park C, Rhee CM, et al. Lymphocyte Cell Ratios and Mortality among Incident Hemodialysis Patients. *Am J Nephrol.* 2017;46(5):408–16.
15. Kato S, Abe T, Lindholm B, Maruyama S. Neutrophil/lymphocyte ratio: A promising prognostic marker in patients with chronic kidney disease. *Inflamm Cell Signal.* 2015;1–5.
16. Arora P. Chronic kidney disease. *MEDSCAPE.* 2019;
17. López-Novoa JM, Martínez-Salgado C, Rodríguez-Peña AB, Hernández FJL. Common pathophysiological mechanisms of chronic kidney disease: Therapeutic perspectives. *Pharmacol Ther* [Internet]. 2010;128(1):61–81. Available from: <http://dx.doi.org/10.1016/j.pharmthera.2010.05.006>
18. Yao Q, Axelsson J, Stenvinkel P, Lindholm B. Chronic systemic inflammation in dialysis patients: An update on causes and consequences. *ASAIO J.* 2004;50(6).
19. Obi Y, Qader H, Kovesdy CP, Kalantar-Zadeh K. Latest consensus and update on protein-energy wasting in chronic kidney disease. *Curr Opin Clin Nutr Metab Care.* 2015;18(3):254–62.

20. Carrero JJ, Stenvinkel P, Cuppari L, Ikizler TA, Kalantar-Zadeh K, Kaysen G, et al. Etiology of the Protein-Energy Wasting Syndrome in Chronic Kidney Disease: A Consensus Statement From the International Society of Renal Nutrition and Metabolism (ISRNM). *J Ren Nutr.* 2013;23(2):77–90.
21. Nitta K, Tsuchiya K. Recent advances in the pathophysiology and management of protein-energy wasting in chronic kidney disease. *Ren Replace Ther* [Internet]. 2016;2(1):1–12. Available from: <http://dx.doi.org/10.1186/s41100-016-0015-5>
22. Rani VN, Kavimani S, Soundararajan P, Chamundeeswari D, Kannan G. Correlation between anthropometry, biochemical markers and subjective global assessment-dialysis malnutrition score as predictors of nutritional status of the maintenance hemodialysis patients. *Int J Med Res Heal Sci.* 2015;4(4):852.
23. Spatola L, Finazzi S, Calvetta A, Reggiani F, Morenghi E, Santostasi S, et al. Subjective Global Assessment-Dialysis Malnutrition Score and cardiovascular risk in hemodialysis patients: an observational cohort study. *J Nephrol.* 2018;31(5):757–65.
24. Xie Y, Zhang H, Ye T, Ge S, Zhuo R, Zhu H. The Geriatric Nutritional Risk Index Independently Predicts Mortality in Diabetic Foot Ulcers Patients Undergoing Amputations. *J Diabetes Res.* 2017;2017.
25. Valga F, Monzón T, Henriquez F, Antón-Pérez G. Neutrophil-to-lymphocyte

- and platelet-to-lymphocyte ratios as biological markers of interest in kidney disease. *Nefrologia*. 2019;39(3):243–9.
26. Bronze-Da-Rocha E, Santos-Silva A. Neutrophil elastase inhibitors and chronic kidney disease. *Int J Biol Sci*. 2018;14(10):1343–60.
 27. Amarya S, Singh K, Sabharwal M. Changes during aging and their association with malnutrition. *J Clin Gerontol Geriatr*. 2015;6(3):78–84.
 28. MV Pahl NV. Immune Function in Chronic Kidney Disease in Chronic Renal Disease. In 2015.
 29. Shaarawy A, Baki AH, Teama NM, Abdel Halim RM, Eldin Fahim NA, Sultan RA. Neutrophils to Lymphocytes Ratio is an Easy Non Expensive Marker of Inflammation in Hemodialysis Patients. *J Clin Exp Nephrol*. 2018;03(04):1–6.
 30. Azab B, Camacho-Rivera M, Taioli E. Average values and racial differences of neutrophil lymphocyte ratio among a nationally representative sample of United States subjects. *PLoS One*. 2014;9(11).
 31. Turkmen K, Erdur FM, Ozcicek F, Ozcicek A, Akbas EM, Ozbicer A, et al. Platelet-to-lymphocyte ratio better predicts inflammation than neutrophil-to-lymphocyte ratio in end-stage renal disease patients. *Hemodial Int*. 2013;17(3):391–6.
 32. Stenvinkel P, Chung SH, Heimbürger O, Lindholm B. Malnutrition,

inflammation, and atherosclerosis in peritoneal dialysis patients. *Perit Dial Int.* 2001;21(SUPPL. 3).

33. Turkmen K, Kayikcioglu H, Ozbek O, Solak Y, Kayrak M, Samur C, et al. The relationship between epicardial adipose tissue and malnutrition, inflammation, atherosclerosis/calcification syndrome in ESRD patients. *Clin J Am Soc Nephrol.* 2011;6(8):1920–5.
34. Behairy M. Neutrophil to Lymphocyte Ratio and Platelet to Lymphocyte Ratio in the Evaluation of Inflammation and Nutritional Status Among Predialysis Chronic Kidney Disease Patients. *Nephrol - Dial - Transplant.* 2019;
35. Ahab E. Neutrophil to lymphocyte and platelet to lymphocyte ratio. A novel markers of inflammation in chronic hemodialysis patients. *Nephrol Dial Transplant.* 2014;29(suppl 3).
36. Nelly N, Widaningsih Y, Mangarengi F. Nilai rasio netrofil limfosit dan rasio trombosit limfosit pada pasien chronic kidney disease sebelum dan setelah hemodialisa. *Intisari Sains Medis.* 2019;10(2):230–4.
37. Kanasaki K, Kitada M, Koya D. Pathophysiology of the aging kidney and therapeutic interventions. Vol. 35, *Hypertension Research.* 2012. p. 1121–8.
38. Goldberg I, Krause I. the Role of Gender in Chronic Kidney Disease. *Cit EMJ.* 2016;1(2):58–64.

39. Adjei DN, Stronks K, Adu D, Snijder MB, Modesti PA, Peters RJG, et al. Relationship between educational and occupational levels, and Chronic Kidney Disease in a multi-ethnic sample- The HELIUS study. *PLoS One*. 2017;12(11):1–14.
40. Min R, Wang H, Zhang X, Li X, Fang P, Bai X. Facing the urban–rural gap in patients with chronic kidney disease: Evidence from inpatients with urban or rural medical insurance in central China. *PLoS One*. 2018;13(12):1–14.
41. Hassanin IA, Hassanein H, Elmenshawy P, El-Gameel D, Elsheikh AA, El-Kobrosly A, et al. Malnutrition score and Body Mass Index as nutritional screening tools for hemodialysis patients. *Clin Nutr ESPEN*. 2021;42:403–6.
42. Alba MM, Citarelli AN, Menni F, Agricola M, Braicovich A, De Horta E, et al. Tobacco and end stage renal disease: A multicenter, cross-sectional study in Argentinian Northern Patagonia. *Tob Induc Dis*. 2015;13(1):1–8.
43. Kaya T, Açıkgöz SB, Yıldırım M, Nalbant A, Altaş AE, Cinemre H. Association between neutrophil-to-lymphocyte ratio and nutritional status in geriatric patients. *J Clin Lab Anal*. 2019;33(1).
44. Foley RN, Collins AJ. End-stage renal disease in the United States: An update from the United States Renal Data System. *J Am Soc Nephrol*. 2007;18(10):2644–8.
45. Collins AJ, Foley RN, Gilbertson DT, Chen SC. United States Renal Data

System public health surveillance of chronic kidney disease and end-stage renal disease. *Kidney Int Suppl.* 2015;5(1):2–7.

46. Antunes SA, Canziani MEF, Campos AF, Vilela RQB. Hypoalbuminemia seems to be associated with a higher rate of hospitalization in hemodialysis patients. *J Bras Nefrol.* 2016;38(1):70–5.
47. Qian X, Guo X, Wang H, Ye M, Zhao Y, Xue G, et al. Neutrophil-lymphocyte Ratio (NLR) and Platelet-Lymphocyte Ratio (PLR) Predict Severity and Prognosis of Lower Limb Arteriosclerosis Obliterans. *Ann Vasc Surg.* 2019;
48. Zhao WM, Tao SM, Liu GL. Neutrophil-to-lymphocyte ratio in relation to the risk of all-cause mortality and cardiovascular events in patients with chronic kidney disease: a systematic review and meta-analysis. *Ren Fail.* 2020;42(1):1059–66.
49. Zhang J, Lu X, Wang S, Li H. High Neutrophil-to-Lymphocyte Ratio and Platelet-to-Lymphocyte Ratio Are Associated with Poor Survival in Patients with Hemodialysis. *Biomed Res Int.* 2021;2021.
50. Baumeister SE, Fischer B, Döring A, Koenig W, Zierer A, John J, et al. The Geriatric Nutritional Risk Index predicts increased healthcare costs and hospitalization in a cohort of community-dwelling older adults: Results from the MONICA/KORA Augsburg cohort study, 1994-2005. *Nutrition.* 2011;27(5):534–42.

51. Kitamura M, Izawa KP, Yaekura M, Mimura Y, Ikeda Y, Nagashima H, et al. Relationship among activities of daily living, nutritional status, and 90 day readmission in elderly patients with heart failure. *Int J Environ Res Public Health*. 2019;16(24).
52. Kobayashi I, Ishimura E, Kato Y, Okuno S, Yamamoto T, Yamakawa T, et al. Geriatric Nutritional Risk Index , a simplified nutritional screening index , is a significant predictor of mortality in chronic dialysis patients. 2010;1417–20.
53. Takahashi S, Suzuki K, Kojima F, Tanaka Y, Nitta K. Geriatric Nutritional Risk Index as a Simple Predictor of Mortality in Maintenance Hemodialysis Patients : A Single Center Study. 2015;(May):354–62.

