

BAB IV

DAFTAR PUSTAKA

- Alasil T, Wang K, Keane PA, Lee H, Baniasadi N, de Boer JF, Chen TC. Analysis of normal retinal nerve fiber layer thickness by age, sex, and race using spectral domain optical coherence tomography. *J Glaucoma*. 2013 Sep; 22(7):532-1. doi:10.1097/IJG.0b013e318255bb4a. PMID: 22549477.
- Aldebasi YH. Prevalence of amblyopia in primary school children in Qassim province, Kingdom of Saudi Arabia. *Middle East Afr J Ophthalmol*. 2015 Jan-Mar; 22(1):86-91.
- Altintas O, Yuksel N, Ozkan B, Caglar Y. Thickness of the retinal nerve fiber layer, macular thickness, and macular volume in patients with strabismic amblyopia. *J Pediatr Ophthalmol Strabismus* 2005; 42:216 –21.
- Andalib D, Javadzadeh A, Nabai R, Amizadeh Y. Macular and retinal nerve fiber layer thickness in unilateral anisometropic or strabismic amblyopia. *J Pediatr Ophthalmol Strabismus*. 2013 Jul-Aug; 50(4):218-21. doi: 10.3928/01913913-20130319-02. Epub 2013 Mar 26. PMID: 23521028.
- Araki S, Miki A, Yamashita T, et al. A comparison between amblyopic and fellow eyes in unilateral amblyopia using spectral-domain optical coherence tomography. *Clin Ophthalmol*. 2014; 8:2199-2207. Published 2014 Nov 7. doi:10.2147/OPHTH.S69501
- Archer SM. Amblyopia? *J AAPOS*. 2000; 4:257.
- Baddini-Caramelli C, Hatanaka M, Polati M, Umino AT, Susanna R Jr. Thickness of the retinal nerve fiber layer in amblyopic and normal eyes: a scanning laser polarimetry study. *J AAPOS*. 2001; 5:82-84.
- Baddini-Caramelli C, Hatanaka M, Polati M, Umino AT, Susanna R Jr. Thickness of the retinal nerve fiber layer in amblyopic and normal eyes: a scanning laser polarimetry study. *J AAPOS*. 2001; 5:82-84.
- Balazsi AG, Rootman J, Drance SM, et al. The effect of age on the nerve fiber population of the human optic nerve. *Am J Ophthalmol* 1984; 97:760–6.

- Bendschneider D, Tornow RP, Horn FK, et al. Retinal nerve fiber layer thickness in normals measured by spectral domain OCT. *J Glaucoma* 2010;19:475-82.
- Bentaleb ZM, Jouffroy E, Rabilloud M, Grange JD, Kodjikian L. Comparison of Central Macular Thickness Measured by Three OCT Models and Study of Interoperator Variability. *The Scientific World Journal*. Volume 2012, Article ID 842795, 6 pages.
- Bhoiwala DL, Simon JW, Raghu P, Krishnamoorthy M, Todani A, Gandham SB, Simmons S. Optic nerve morphology in normal children. *Journal of American Association for Pediatric Ophthalmology and Strabismus*. Volume 19, Issue 6, December 2015, Pages 531-534.
- Bourne RR, Medeiros FA, Bowd C, et al. Comparability of retinal nerve fiber layer thickness measurements of optical coherence tomography instruments. *Invest Ophthalmol Vis Sci*. 2005;46:1280–1285.
- Bowd C, Weinreb RN, Williams JM, et al. The retinal nerve fiber layer thickness in ocular hypertensive, normal, and glaucomatous eyes with optical coherence tomography. *Arch Ophthalmol*. 2000;118:22–26.
- Bozkurt B, Irkeç M, Orhan M, Karaağaoğlu E. Thickness of the retinal nerve fiber layer in patients with anisometropic and strabismic amblyopia. *Strabismus*. 2003.11:1-7.1119-24.
- Branislav Stanković. Clinical aspects of different types of amblyopia. Faculty of Medicine, University of Belgrade, Belgrade, Serbia; Clinical Center of Serbia, Institute of Ophthalmology, Belgrade, Serbia. 2010
- Budenz DL, Anderson DR, Varma R, et al. Determinants of normal retinal nerve fiber layer thickness measured by Stratus OCT. *Ophthalmology*. 2007;114(6):1046e1052.
- Caprioli J. The contour of the juxtapapillary nerve fiber layer in glaucoma. *Ophthalmology*. 1990;97:358–365; discussion 365–356.
- Carpineto P, Ciancaglini M, Aharrh-Gnama A, et al. Custom measurement of retinal nerve fiber layer thickness using STRATUS OCT in normal eyes. *Eur J Ophthalmol*. 2005;15: 360–366.

- Cheung N, Huynh S, Wang JJ, et al. Relationships of retinal vessel diameters with optic disc, macular and retinal nerve fiber layer parameters in 6-year-old children. *Invest Ophthalmol Vis Sci* 2008;49:2403-8.
- Chow KL. Failure to demonstrate changes in the visual system of monkeys kept in darkness or in colored lights. *J Comp Neurol*. 1955;102:597-606.
- Colen T, de Faber JT, Lemij HG. Retinal nerve fiber layer thickness in human strabismic amblyopia. *Binocul Vis Strabismus Q* 2000;15:141–146. [PubMed: 10893456]
- Demer J, Clark RA, Lim KH, Engle ED. Magnetic resonance imaging of innervational and extraocular muscle abnormalities in Duane-radial ray syndrome. *Invest Ophthalmol Vis Sci* 2007;48:5505–5511. [PubMed: 18055799]
- Dickmann A, Petroni S, Salerni A, et al. Unilateral amblyopia: An optical coherence tomography study. *J AAPOS*. 2009;13:148–150.
- Duranoglu Y. Optic nerve head topographic analysis and retinal nerve fiber layer thickness in strabismic and anisometropic amblyopia. *Ann Ophthalmol (Skokie)* 2007;39:291–295. [PubMed: 18025648]
- El-Dairi MA, Asrani SG, Enyedi LB, Freedman SF. Optical coherence tomography in the eyes of normal children. *Arch Ophthalmol* 2009;127:50-58.
- Fu J, Li SM, Liu LR, et al.; Anyang Childhood Eye Study Group. Prevalence of amblyopia and strabismus in a population of 7th-grade junior high school students in Central China: the Anyang Childhood Eye Study (ACES). *Ophthalmic Epidemiol*. 2014 Jun;21(3):197-203.
- Ganekal S, Jhanji V, Liang Y, Dorairaj S. Prevalence and etiology of amblyopia in Southern India: results from screening of school children aged 5-15 years. *Ophthalmic Epidemiol*. 2013 Aug;20(4):228-31.
- Giordano L, Friedman DS, Repka MX, Katz J, Ibrionke J, Hawes P, Tielsch JM. Prevalence of refractive error among preschool children in an urban population: the Baltimore Pediatric Eye Disease Study. *Ophthalmology*. 2009;116:739-746.
- Greenfield, D.S., Huang, X.-R., Knighton, R.W., 2000. Effect of corneal polarization axis on assessment of retinal nerve fiber layer thickness by scanning laser polarimetry. *Am. J. Ophthalmol.* 129, 715–722.

- Harizman N, Oliveira C, Chiang A, et al. The ISNT rule and differentiation of normal from glaucomatous eyes. *Arch Ophthalmol*. 2006;124:1579–1583.
- Headon MP, Powell TS. Cellular changes in the lateral geniculate nucleus of infant monkeys after suture of eyelids. *J Anat*. 1973;116:135-145.
- Hee MR, Izatt JA, Swanson EA, Huang D, Schuman JS, Lin CP, Puliafito CA, Fujimoto JG. Optic coherence tomography of the human retina. *Arch Ophthalmol*. 1995;113:325-332.
- Henrietta Ho, FRCOphth, Yih-Chung Tham, PhD, Miao Li Chee, BSc, Yuan Shi, PhD, Nicholas Y.Q. Tan, MA, BM BCh, Kah-Hie Wong, MBBS, Shivani Majithia, OD, Carol Y. Cheung, PhD, Tin Aung, FRCOphth, PhD, Tien Yin Wong, FRCS(Ed), PhD, Ching-Yu Cheng, MD, PhD. Retinal Nerve Fiber Layer Thickness in a Multiethnic Normal Asian Population The Singapore Epidemiology of Eye Diseases Study. *American academy of Ophthalmology*. December, 4 2018: 702-711.
- Hess RF, Field DJ. Is the spatial deficit in strabismic amblyopia due to loss of cells or an uncalibrated disarray of cells? *Vision Res*. 1994;34:3397-3406.
- Holmes JM, Clarke MP. Amblyopia. *Lancet*. 2006;367:1343–1351.
- Hougaard JL, Ostensfeld C, Heijl A, et al. Modelling the normal retinal nerve fibre layer thickness as measured by Stratus optical coherence tomography. *Graefes Arch Clin Exp Ophthalmol*. 2006;244:1607–1614.
- Huynh SC, Samarawickrama C, Wang XY, Rohtchina E, Wong TY, Gole GA, Rose KA, Mitchell P. Macular and nerve fiber layer thickness in amblyopia: the Sydney Childhood Eye Study. *Ophthalmology*. 2009;116:1604-1609.
- Huynh SC, Wang XY, Rohtchina E, Mitchell P. Distribution of macular thickness by optical coherence tomography: findings from a population-based study of 6-year-old children. *Invest Ophthalmol Vis Sci* 2006;47:2351-7.
- Huynh SC, Wang XY, Rohtchina E, Mitchell P. Peripapillary retinal nerve fiber layer thickness in a population of 6-year-old children: findings by optical coherence tomography. *Ophthalmology* 2006;113:1583-92.

- Jonas JB, Gusek GC, Naumann GO. Optic disc, cup and neuroretinal rim size, configuration and correlations in normal eyes. *Invest Ophthalmol Vis Sci.* 1988;29:1151–1158
- Jonas JB, Nguyen NX, Naumann GO. The retinal nerve fiber layer in normal eyes. *Ophthalmology* 1989;96:627–32.
- Kanamori A, Kusuhara A, Tatsumi Y, et al. [Correlations among GDx-variable corneal compension, optical coherence tomography, and Heidelberg retina tomograph and relationships between these structural parameters and visual field indices]. *Nihon Ganka Gakkai Zasshi.* 2006;110(3):180–187. Japanese.
- Kasem MA, Badawi AE. Changes in macular parameters in different types of amblyopia: optical coherence tomography study. *Clin Ophthalmol.* 2017 Aug 4;11:1407-1416. doi: 10.2147/OPHTH.S143223. PMID: 28831240; PMCID: PMC5552145.
- Kavitha V, Heralgi MM, Harishkumar PD, Harogoppa S, Shivaswamy H M, Geetha H. Analysis of Macular, Foveal, and Retinal Nerve Fiber Layer Thickness in Children with Unilateral Anisometric Amblyopia and Their Changes Following Occlusion Therapy. *Indian J Ophthalmol* 2019;67:1016-22.
- Kee SY, Lee SY, Lee YC. Thicknesses of the fovea and retinal nerve fiber layer in amblyopic and normal eyes in children. *Korean J Ophthalmol.* 2006;20:177-181.
- Kim YW, Kim SJ, Yu YS. Spectral-domain optical coherence tomography analysis in deprivational amblyopia: a pilot study with unilateral pediatric cataract patients. *Graefes Arch Clin Exp Ophthalmol.* 2013;251(12):2811–2819.
- Kiorpes L, Kiper DC, O'Keefe LP, Cavanaugh JR, Movshon JA. Neuronal correlates of amblyopia in the visual cortex of macaque monkeys with experimental strabismus and anisometropia. *J Neurosci.*1998;18:6411-6424.
- Lam DS, Leung KS, Mohamed S, et al. Regional variations in the relationship between macular thickness measurements and myopia. *Invest Ophthalmol Vis Sci.* 2007;48:376–382.
- Larsson E, Nuija E, Alm A. The optic nerve head assessed with HRT in 5-16-year-old normal children: normal values, repeatability and interocular difference. *Acta Ophthalmol.* 2011; 89:755–758.

- Leguire LE, Rogers GL, Bremer DL. Amblyopia: the normal eye is not normal. *J Pediatr Ophthalmol Strabismus*. 1990;27:32-38.
- Lempert P. Optic nerve hypoplasia and small eyes in presumed amblyopia. *J AAPOS*. 2000;4:258-266.
- Lempert P. Retinal area and optic disc rim area in amblyopic, fellow, and normal hyperopic eyes: a hypothesis for decreased acuity in amblyopia. *Ophthalmology*. 2008;115:2259-2261.
- Lempert P. The axial length/disc area ratio in anisometropic hyperopic amblyopia: a hypothesis for decreased unilateral vision associated with hyperopic anisometropia. 2004;111:304-308
- Lempert P. Axial length-disc area ratio in esotropic amblyopia. 2003;121:821-824
- Leung CK, Mohamed S, Leung KS, et al. Retinal nerve fiber layer measurements in myopia: an optical coherence tomography study. *Invest Ophthalmol Vis Sci*. 2006;47: 5171–5176
- Levi DM. Linking assumptions in amblyopia. *Vis Neurosci*. 2013 Nov;30(5-6):277-87.
- Li J, Ji P, Yu M (2015) Meta-analysis of retinal changes in unilateral amblyopia using optical coherence tomography. *Eur J Ophthalmol* 25:400-409
- Liu H, Zhong L, Zhou X, Jin QZ. Macular abnormality observed by OCT in children with amblyopia failing to achieve normal visual acuity after long-term treatment. *J Pediatr Ophthalmol Strabismus* 2010;47:17-23
- Lu AT, Wang M, Varma R, et al. Combining nerve fiber layer parameters to optimize glaucoma diagnosis with optical coherence tomography. *Ophthalmology*. 2008;115:1352.e2–1357.e2.
- Martinez A, Proupim N, Sanchez M. Retinal nerve fibre layer thickness measurements using optical coherence tomography in migraine patients. *Br J Ophthalmol*. 2008;92:1069–1075.
- Miki A, Shirakashi M, Yaoeda K, et al. Retinal nerve fiber layer thickness in recovered and persistent amblyopia. *Clin Ophthalmol*. 2010;4:1061-1064. Published 2010 Sep 20. doi:10.2147/opth.s13145

- Mistlberger A, Liebmann JM, Greenfield DS, et al. Heidelberg retina tomography and optical coherence tomography in normal, ocular hypertensive, and glaucomatous eyes. *Ophthalmology* 1999;106:2027–32.
- Mistlberger A, Liebmann JM, Greenfield DS, et al. Heidelberg retina tomography and optical coherence tomography in normal, ocular-hypertensive, and glaucomatous eyes. *Ophthalmology*. 1999;106:2027–2032.
- Neelam P, et al. Retinal Nerve Fiber Layer Thickness and Optic Disc Parameters in Anisometropic Amblyopia by Cirrus HD OCT in Pediatric Population. *J Ophthalmol* 2017, 2(1): 000112
- Oscar A, Cherninkova S, Haykin V, et al. Amblyopia screening in Bulgaria. *J Pediatr Ophthalmol Strabismus*. 2014 Sep-Oct;51(5):284-8
- Yi Pang, Geoffrey W. Goodfellow, Christine Allison, Sandra Block, Kelly A. Frantz; A Prospective Study of Macular Thickness in Amblyopic Children with Unilateral High Myopia. *Invest. Ophthalmol. Vis. Sci.* 2011;52(5):2444-2449. doi: <https://doi.org/10.1167/iovs.10-5550>.
- Papastathopoulos KI, Jonas JB, Panda-Jonas S. Large optic discs in large eyes, small optic discs in small eyes. *Exp Eye Res*. 1995;60:459-461.
- Parikh RS, Parikh SR, Sekhar GC, et al. Normal age-related decay of retinal nerve fiber layer thickness. *Ophthalmology*. 2007;114(5):921e926.
- Park KA, Park DY, Oh SY. Analysis of spectral-domain optical coherence tomography measurements in amblyopia: A pilot study. *Br J Ophthalmol* 2011;95:1700-1706.
- Pascual M, Huang J, Maguire MG, et al; Vision in Preschoolers (VIP) Study Group. Risk factors for amblyopia in the vision in preschoolers study. *Ophthalmology*. 2014 Mar;121(3):622-9.
- Paunescu LA, Schuman JS, Price LL, et al. Reproducibility of nerve fiber thickness, macular thickness, and optic nerve head measurements using Stratus OCT. *Invest Ophthalmol Vis Sci*. 2004;45:1716–1724.
- Pineles SL, Demer JL. Bilateral abnormalities of optic nerve size and eye shape in unilateral amblyopia. *Am J Ophthalmol*. 2009;148(4):551-557.e2. doi:10.1016/j.ajo.2009.05.007.

Poinosawmy D, Fontana L, Wu JX, et al. Variation of nerve fibre layer thickness measurements with age and ethnicity by scanning laser polarimetry. *Br J Ophthalmol* 1997;81:350–4

Potts RA, Dreher B, Bennett MR. The loss of ganglion cells in the developing retina of the rat. *Brain Res.* 1982;255:481–6.

Provis, J.M., van Driel, D., Billson, F.A., Russell, P., 1965. Human fetal optic nerve: overproduction and elimination of retinal axons during development. *J. Comp. Neurol.* 238, 92–100.

Provis, J.M., van Driel, D., Billson, F.A., Russell, P., 1985. Development of the human retina: patterns of cell distribution and redistribution in the ganglion cell layer. *J. Comp. Neurol.* 233, 429–451.

Quigley HA, Coleman AL, Dorman-Pease ME. Larger optic nerve heads have more nerve fibers in normal monkey eyes. *Arch Ophthalmol.* 1991;109:1441-1443.

Quoc EB, Delepine B, Tran TH. Thickness of retinal nerve fiber layer and macular volume in childrens and adults with strabismic and anisometropic amblyopia. *J Fr Ophtalmol.* 2009;32:488-495.

Rajavi Z, Moghadasifar H, Feizi M, Haftabadi N, Hadavand MR, Yaseri M, et al. Macular thickness and amblyopia. *J Ophthalmic Vis Res* 2014;9:478-83.

Rasch E, Swift H, Riesen AH, Chow KL. Altered structure and composition of retinal cells in dark reared mammals. *Exp Cell Res.* 1961;25:348-363.

Repka MX, Goldenberg-Cohen N, Edwards AR. Retinal nerve fiber layer thickness in amblyopic eyes. *Am J Ophthalmol* 2006;142:247-251

Repka MX, Kraker RT, Tamkins SM, Suh DW, Sala NA, Beck RW; Pediatric Eye Disease Investigator Group. Retinal nerve fiber layer thickness in amblyopic eyes. *Am J Ophthalmol.* 2009;148:143-147.

Savini G, Zanini M, Carelli V, et al. Correlation between retinal nerve fibre layer thickness and optic nerve head size: an optical coherence tomography study. *Br J Ophthalmol.* 2005;89:489–492.

Schuman JS, Hee MR, Puliafito CA, et al. Quantification of nerve fiber layer thickness in normal and glaucomatous eyes using optical coherence tomography. *Arch Ophthalmol* 1995;113:586–96.

- Schuman JS, Pedut-Kloizman T, Hertzmark E, et al. Reproducibility of nerve fiber layer thickness measurements using optical coherence tomography. *Ophthalmology* 1996;103:1889–98.
- Schuman JS, Wollstein G, Farra T, et al. Comparison of optic nerve head measurements obtained by optical coherence tomography and confocal scanning laser ophthalmoscopy. *AmJ Ophthalmol.* 2003;135:504–512.
- Sehi M, Ume S, Greenfield DS. Scanning laser polarimetry with enhanced corneal compensation and optical coherence tomography in normal and glaucomatous eyes. *Invest Ophthalmol Vis Sci.* 2007;48:2099–2104.
- Siddiqui AH, Raza SA, Ghazipura A, Hussain MA, Iqbal S, Ahsan K, Nadeem A. Analysis of association between type of amblyopia and gender at a tertiary care hospital in Karachi. *J Pak Med Assoc.* 2016 May;66(5):545-8. PMID: 27183933.
- Sihota R, Srinivasan G, Dada T, et al. Is the ISNT rule violated in early primary open-angle glaucoma—a scanning laser tomography study. *Eye (Lond).* 2008;22:819–824.
- Silva F, Alves S, Pina S, Azevedo A, Pêgo P, Santos MJ, et al. Comparison of macular thickness and volume in amblyopic children using time domain optical coherence tomography. *Oftalmologia* 2012;36:231-236.
- Sung KR, Wollstein G, Bilonick RA, et al. Effects of age on optical coherence tomography measurements of healthy retinal nerve fiber layer, macula, and optic nerve head. *Ophthalmology* 2009;116:
- Syunsuke Araki, Atsushi Miki, Tsutomu Yamashita, Katsutoshi Goto, Kazuko Haruishi, Yoshiaki Ieki, Junichi Kiryu. A comparison between amblyopic and fellow eyes in unilateral amblyopia using spectral-domain optical coherence tomography. Department of Ophthalmology, Kawasaki Medical School, Kurashiki, Japan; Department of Sensory Science, Faculty of Health Science and Technology, Kawasaki University of Medical Welfare, Kurashiki, Japan. 2014.
- Szigeti A, Ta'raí E, Szamosi A, Vargha P, Nagy ZZ, et al. (2014) A Morphological Study of Retinal Changes in Unilateral Amblyopia Using Optical Coherence

- Tomography Image Segmentation. PLoS ONE 9(2): e88363. doi:10.1371/journal.pone.0088363
- Tan JC, Poinosawmy D, Fitzke FW, Hitchings RA. Magnification Changes In Scanning Laser Tomography. J Glaucoma. 2004; 13:137–41. [PubMed: 15097259]
- Tariq YM MBBS, Li H PhD, Burlutsky G MStat and Mitchell P PhD. Ethnic Differences In Macular Thickness. Centre for Vision Research, Department of Ophthalmology and Westmead Millennium Institute, University of Sydney, Sydney, New South Wales, Australia. Clinical and Experimental Ophthalmology. 2011
- Varma R, Quigley HA, Pease ME. Changes in optic disc characteristics and number of nerve fibers in experimental glaucoma. *Am J Ophthalmol.* 1992;114(5):554–559.
- Varma R, Skaf M, Barron E. Retinal nerve fiber layer thickness in normal human eyes. *Ophthalmology.* 1996;103: 2114–2119.
- Von Noorden GK, Crawford ML, Middleditch PR. Effect of lid suture on retinal ganglion cells in Macaca Mulatta. *Brain Res.* 1977;122:437-444.
- Von Noorden GK, Crawford ML. The lateral geniculate nucleus in human strabismic amblyopia. *Invest Ophthalmol Vis Sci.* 1992;33:2729-2732.
- Von Noorden GK, Middleditch PR. Histology of the monkey lateral geniculate nucleus after unilateral lid closure and experimental strabismus: further observations. *Invest Ophthalmol.* 1975;14:674-683.
- Von Noorden GK. Histological studies of the visual system in monkeys with experimental amblyopia. *Invest Ophthalmol.* 1973;12:727-738.
- Walker RA, Rubab S, Voll AR, Erraguntla V, Murphy PH. Macular and peripapillary retinal nerve fibre layer thickness in adults with amblyopia. *Can J Ophthalmol.* 2011;46:425-427.
- Wendell-Smith CP. Effect of light deprivation on the postnatal development of the optic nerve. *Nature.* 1964;204:707.

- Wollstein G, Ishikawa H, Wang J, et al. Comparison of three optical coherence tomography scanning areas for detection of glaucomatous damage. *Am J Ophthalmol*. 2005;139:39–43.
- Wu PC, Chen YJ, Chen CH, et al. Assessment of macular retinal thickness and volume in normal eyes and highly myopic eyes with third-generation optical coherence tomography. *Eye (Lond)*. 2008; 22:551–555.
- Yen MY, Cheng CY, Wang AG. Retinal nerve fiber layer thickness in unilateral amblyopia. *Invest Ophthalmol Vis Sci*. 2004;45:2224-2230.
- Yoon SW, Park WH, Baek SH, Kong SM. Thicknesses of macular retinal layer and peripapillary retinal nerve fiber layer in patients with hyperopic anisometropic amblyopia. *Korean J Ophthalmol* 2005;19:62–7.
- Yücel YH, Gupta N, Kalichman MW, et al. Relationship of optic disc topography to optic nerve fiber number in glaucoma. *Arch Ophthalmol*. 1998;116(4):493–497.
- Zhu H, Pan C, Sun Q, et al. Prevalence of amblyopia and strabismus in Hani school children in rural southwest China: a crosssectional study. *BMJ Open* 2019;9:e025441. doi:10.1136/ bmjopen- 2018-025441