

## DAFTAR PUSTAKA

- Anderson CM, Ralph JL, Johnson L, Scheett A, Wright ML, Taylor JY, Ohm JE, Uthus E. First trimester vitamin D status and placental epigenomics in preeclampsia among Northern Plains primiparas. *Life Sci.* 2015;129:10–5.
- Audibert F, Luo ZC, Nuyt AM, Masse B, Julien P, Fraser WD, Group MS. Maternal plasma 25-hydroxyvitamin D levels, angiogenic factors, and preeclampsia. *Am J Obstet Gynecol.* 2013;208(5):390 e391–6.
15. Woodham PC, Brittain JE, Baker AM, Long DL, Haeri S, Camargo CA Jr, Ardawi MS, Nasrat HA, BA'Aqueel HS. Calcium-regulating hormones and parathyroid hormone-related peptide in normal human pregnancy and postpartum: a longitudinal study. *Eur J Endocrinol.* 1997;137(4):402–9.
- Achkar M, Dodds L, Giguere Y, Forest JC, Armson BA, Woolcott C, Agellon S, Spencer A, Weiler HA. Vitamin D status in early pregnancy and risk of preeclampsia. *Am J Obstet Gynecol.* 2015;212(4):511 e511–7.
- Abedi P, Mohaghegh Z, Afshary P, Latifi M. The relationship of serum vitamin D with pre-eclampsia in the Iranian women. *Matern Child Nutr.* 2014;10(2):206–12.
- August P, Marcaccio B, Gertner JM, Druzin ML, Resnick LM, Laragh JH. Abnormal 1,25-dihydroxyvitamin D metabolism in preeclampsia. *Am J Obstet Gynecol.* 1992;166(4):1295–9.
- Alvarez-Fernandez I, Prieto B, Rodriguez V, Ruano Y, Escudero AI, Alvarez FV. Role of vitamin D and sFlt-1/PIGF ratio in the development of early- and late-onset preeclampsia. *Clin Chem Lab Med.* 2015;53(7):1033–40.
- Arngrimsson R, Bjornsson S, Geirsson RT, Bjornsson H et al. (1990) Genetic and familial predisposition to eclampsia and pre-eclampsia in a defined population. *Br J Obstet Gynaecol* 97, 762-769.
- Arngrimsson R, Sigurdar ttir S, Frigge ML, Bjarnad ttir RI et al. (1999) A genome-wide scan reveals a maternal susceptibility locus for pre- eclampsia on chromosome 2p13. *Hum Mol Genet* 8, 1799-1805
- Baxter, JK., 2007. Hypertensive Disorder in: Berghella, V., 2007. *Maternal-Fetal Evidence Based Guidelines*. Informa UK Ltd Chapter 1. p.3-22

- Brown MA, Hague WM, Higgins J, Lowe S, McCowan L, Oats J, Peek MJ, Rowan JA, Walters BN. The Australasian Society of the Study of Hypertension in Pregnancy. The detection, investigation and management of hypertension in pregnancy: full consensus statement. *Aust N Z J Obstet Gynaecol* 2000; 40 (2):139 – 155.
- Baum M, Schiff E, Kreiser D, Dennerly PA, Stevenson DK, Rosenthal T, Seidman DS. End-tidal carbon monoxide measurements in women with pregnancy-induced hypertension and preeclampsia. *Am J Obstet Gynecol.* 2000;183:900–903.
- Boucher BJ. The 2010 recommendations of the American Institute of Medicine for daily intakes of vitamin D. *Public Health Nutr.*2011;14(4):740.
- Badria LF, Amarin ZO. Pre-eclampsia: is it a different disease in primiparous and multiparous women? *Arch Gynecol Obstet.* 2005;273(1):26–31.
- Burris HH, Rifas-Shiman SL, Huh SY, Kleinman K, Litonjua AA, Oken E, Rich- Edwards JW, Camargo CA Jr, Gillman MW. Vitamin D status and hypertensive disorders in pregnancy. *Ann Epidemiol.* 2014;24(5):399–403. e391
- Bodnar LM, Simhan HN, Catov JM, Roberts JM, Platt RW, Diesel JC, Klebanoff MA. Maternal vitamin D status and the risk of mild and severe preeclampsia. *Epidemiology.* 2014;25(2):207–14.
- Baker AM, Haeri S, Camargo CA Jr, Espinola JA, Stuebe AM. A nested case- control study of midgestation vitamin D deficiency and risk of severe preeclampsia. *J Clin Endocrinol Metab.* 2010;95(11):5105–9.
- Belizan JM, Villar J, Gonzalez L, Campodonico L, Bergel E. Calcium supplementation to prevent hypertensive disorders of pregnancy. *N Engl J Med.* 1991;325(20):1399–405.
- Brodowski L, Burlakov J, Myerski AC, von Kaisenberg CS, Grundmann M, Hubel CA, von Versen-Hoyneck F. Vitamin D prevents endothelial progenitor cell dysfunction induced by sera from women with preeclampsia or conditioned media from hypoxic placenta. *PLoS One.* 2014;9(6):e98527.
- Barrera D, Noyola-Martinez N, Avila E, Halhali A, Larrea F, Diaz L. Calcitriol inhibits interleukin-10 expression in cultured human trophoblasts under normal and inflammatory conditions. *Cytokine* 2012;57(3):316–21.
- Bodnar LM, Catov JM, Simhan HN, Holick MF, Powers RW, Roberts

- JM. Maternal vitamin D deficiency increases the risk of preeclampsia. *J Clin Endocrinol Metab.* 2007;92(9):3517–22.
- Boggess KA, Stuebe AM. Midgestation maternal serum 25-hydroxyvitamin D level and soluble fms-like tyrosine kinase 1/placental growth factor ratio as predictors of severe preeclampsia. *Hypertension.* 2011;58(6):1120–5.
- Bamfo, J.E.; Odibo, A.O. Diagnosis and management of fetal growth restriction. *J. Pregnancy* 2011, 2011, doi:10.1155/2011/640715.
- Cunningham, FG., Leveno KJ., Bloom, SL., 2010. *Pregnancy Hypertension. Williams Obstetrics 23<sup>rd</sup> edition Vol 2 Chapter 34.* McGraw-Hill Companies, New York, P.761-803.
- Cottrell, E.C.; Ozanne, S.E. Early life programming of obesity and metabolic disease. *Physiol. Behav.* 2008, 94, 17–28.
- Cardus A, Parisi E, Gallego C, Aldea M, Fernandez E, Valdivielso JM. 1,25-Dihydroxyvitamin D<sub>3</sub> stimulates vascular smooth muscle cell proliferation through a VEGF-mediated pathway. *Kidney Int.* 2006;69(8):1377–84.
- Chan SY, Susarla R, Canovas D, Vasilopoulou E, Ohizua O, McCabe CJ, Hewison M, Kilby MD. Vitamin D promotes human extravillous trophoblast invasion in vitro. *Placenta.* 2015;36(4):403–9.
- Cross NA, Hillman LS, Allen SH, Krause GF, Vieira NE. Calcium homeostasis and bone metabolism during pregnancy, lactation, and postweaning: a longitudinal study. *Am J Clin Nutr.* 1995;61(3):514–23.
- Cudmore M, Ahmad S, Al-Ani B, Fujisawa T, Coxall H, Chudasama K, Devey LR, Wigmore SJ, Abbas A, Hewett PW, Ahmed A. Negative regulation of soluble Flt-1 and soluble endoglin release by heme oxygenase-1. *Circulation.* 2007;115:1789–1797.
- Chua S, Wilkins T, Sargent I, Redman C. Trophoblast deportation in pre-eclamptic pregnancy. *Br J Obstet Gynaecol.* 1991;98:973–979.
- Caniggia I, Taylor CV, Ritchie JW, Lye SJ, Letarte M. 1997. Endoglin regulates trophoblast differentiation along the invasive pathway in human placental villous explants. *Endocrinology* 138:4977–88
- Data rekam medis. BLUD H. Padjongga Daeng Ngalle Kabupaten Takalar. Tahun 2018-Tahun 2020.

- Douglas KA, Redman CWG. Eclampsia in the United Kingdom. *Br Med J* 1994; 309 (6966):1395–1400.
- De-Regil LM, Palacios C, Ansary A, Kulier R, Peña-Rosas JP: Vitamin D supplementation for women during pregnancy (Version 2). *Cochrane Database Syst Rev.* 2012;(2):1-50.
- Diaz L, Arranz C, Avila E, Halhali A, Vilchis F, Larrea F. Expression and activity of 25-hydroxyvitamin D-1 alpha-hydroxylase are restricted in cultures of human syncytiotrophoblast cells from preeclamptic pregnancies. *J Clin Endocrinol Metab.* 2002;87(8):3876–82.
- Dror DK, Allen LH. Vitamin D inadequacy in pregnancy: biology, outcomes, and interventions. *Nutr Rev.* 2010;68(8):465–77.
- Datta S, Alfaham M, Davies DP, Dunstan F, Woodhead S, Evans J, Richards B. Vitamin D deficiency in pregnant women from a non-European ethnic minority population—an interventional study. *BJOG.* 2002;109(8):905–8.
- Duley L. The global impact of pre-eclampsia and eclampsia. *Semin Perinatol.* 2009;33(3):130–7.
- Easterling TR, The Maternal Hemodynamics of Preeclampsia, *Clinic Obstet and Gynecol.* 1992; V. 35, No. 2.
- Easterling TR. 1992. The maternal hemodynamics of preeclampsia. *Clin. Obstet. Gynecol.* 35:375–86
- Esplin MS, Fausett MB, Fraser A, Kerber R et al. (2001) Paternal and maternal components of the predisposition to preeclampsia. *N Engl J Med* 344, 867-872.
- England L, Zhang J. Smoking and risk of preeclampsia: a systematic review. *Front Biosci.* 2007;12:2471–83.
- Evans KN, Bulmer JN, Kilby MD, Hewison M. Vitamin D and placental-decidual function. *J Soc Gynecol Investig.* 2004;11(5):263–71.
- Fischer D, Schroer A, Ludders D, Cordes T, Bucker B, Reichrath J, Friedrich M. Metabolism of vitamin D3 in the placental tissue of normal and preeclampsia complicated pregnancies and premature
- Farina A, Sekizawa A, De Sanctis P, Purwosunu Y, Okai T, et al. 2008. Gene expression in chorionic villous samples at 11 weeks' gestation from women destined to develop preeclampsia. *Prenat. Diagn.* 28:956–61
- Fernandez-Alonso AM, Dionis-Sanchez EC, Chedraui P, Gonzalez-Salmeron MD, Perez-Lopez FR, Spanish Vitamin D, Women's Health Research G. First- trimester maternal serum 25-hydroxyvitamin D(3) status and pregnancy outcome. *Int J*

- Gynaecol Obstet.2012;116(1):6–9.
- Fudge NJ, Kovacs CS. Pregnancy up-regulates intestinal calcium absorption and skeletal mineralization independently of the vitamin D receptor. *Endocrinology*. 2010;151(3):886–95.
- Faulkner JL, Amaral LM, Cornelius DC, Cunningham MW, Ibrahim T, Heep A, Campbell N, Usry N, Wallace K, Herse F, et al. Vitamin D supplementation reduces some AT1-AA-induced downstream targets implicated in preeclampsia including hypertension. *Am J Physiol Regul Integr Comp Physiol*. 2017;312(1):R125–31.
- Finch PJ, Ang L, Colston KW, Nisbet J, Maxwell JD. Blunted seasonal variation in serum 25-hydroxy vitamin D and increased risk of osteomalacia in vegetarian London Asians. *Eur J Clin Nutr*.1992;46(7):509–15.
- Gardosi, J.; Madurasinghe, V.; Williams, M.; Malik, A.; Francis, A. Maternal and fetal risk factors for stillbirth: Population based study. *BMJ* 2013, 346, doi:10.1136/bmj.f108.
- Gynecol. 2013;209(6):544 e541–12. 11. Irgens HU, Reisaeter L, Irgens LM, Lie RT. Long term mortality of mothers and fathers after preeclampsia: population based cohort study. *BMJ*. 2001;323(7323):1213–7. 12.
- Gray TK, Lester GE, Lorenc RS. Evidence for extra-renal 1 alpha-hydroxylation of 25-hydroxyvitamin D3 in pregnancy. *Science*. 1979;204(4399):1311–3.
- Grundmann M, Haidar M, Placzko S, Niendorf R, Darashchonak N, Hubel CA, von Versen-Hoyneck F. Vitamin D improves the angiogenic properties of endothelial progenitor cells. *Am J Physiol Cell Physiol*.2012;303(9):C954–62.
- Gidlof S, Silva AT, Gustafsson S, Lindqvist PG. Vitamin D and the risk of preeclampsia—a nested case-control study. *Acta Obstet Gynecol Scand*. 2015;94(8):904–8.
- Gerber HP, Condorelli F, Park J, Ferrara N. Differential transcriptional regulation of the two vascular endothelial growth factor receptor genes: Flt-1, but not Flk-1/KDR, is up-regulated by hypoxia. *J Biol Chem*. 1997;272:23659–23667.
- Girardi G, Yarin D, Thurman JM, Holers VM, Salmon JE. Complement activation induces dysregulation of angiogenic factors and causes fetal rejection and growth restriction. *J Exp Med*. 2006;203: 2165–2175.

- Graves JA (1998) Genomic imprinting, development and disease--is pre-eclampsia caused by a maternally imprinted gene? *Reprod Fertil Dev* 10, 23-29.
- Gant NF, Chand S, Whalley PJ, MacDonald PC. 1974. The nature of pressor responsiveness to angiotensin II in human pregnancy. *Obstet. Gynecol.* 43:854
- Gant NF, Daley GL, Chand S, Whalley PJ, MacDonald PC. 1973. A study of angiotensin II pressor response throughout primigravid pregnancy. *J. Clin. Investig.* 52:2682–89
- GarovicVD, WagnerSJ, TurnerST, RosenthalDW, WatsonWJ, et al. 2007. Urinary podocyte excretion as a marker for preeclampsia. *Am. J. Obstet. Gynecol.* 196:320e1–7
- HincheyJ, ChavesC, AppignaniB, BreenJ, PaoL, et al. 1996. A reversible posterior leukoencephalopathy syndrome. *N. Engl. J. Med.* 334:494–500
- Homer CS, Brown MA, Mangos G, Davis GK. Non-proteinuric pre-eclampsia: a novel risk indicator in women with gestational hypertension. *J Hypertens* 2008; 26: 295–302.
- Helewa ME, Burrows RF, Smith J, Williams K, Brain P, Rabkin SW. Report of the Canadian Hypertension Society Consensus Conference: 1: definitions, evaluation and classifications of hypertensive disorders in pregnancy. *Can Med Assoc J* 1997; 157:715 – 725.
- Harrison GA, Humphrey KE, Jones N, Badenhop R et al. (1997) A genomewide linkage study of preeclampsia/eclampsia reveals evidence for a candidate region on 4q. *Am J Hum Genet* 60, 1158-1167.
- Hayward C, Livingstone J, Holloway S, Liston WA et al. (1992) An exclusion map for pre-eclampsia: assuming autosomal recessive inheritance. *Am J Hum Genet* 50, 749-757.
- Hubel CA. Oxidative stress in the pathogenesis of preeclampsia. *Proc SocExp Biol Med.* 1999;222:222–235.
- Hubel CA, Wallukat G, Wolf M, Herse F, Rajakumar A, Roberts JM, Markovic N, Thadhani R, Luft FC, Dechend R. Agonistic angiotensin II type 1 receptor autoantibodies in postpartum women with a history of preeclampsia. *Hypertension.* 2007;49:612– 617.
- Hiby SE, Walker JJ, O'Shaughnessy KM, Redman CW, Carrington M, Trowsdale J, Moffett A. Combinations of maternal kir and fetal HLA-C genes influence the risk of preeclampsia and reproductive success. *J Exp Med.* 2004;200:957–965.

- Hanna J, Goldman-Wohl D, Hamani Y, Avraham I, Greenfield C, Natanson-Yaron S, Prus D, Cohen-Daniel L, Arnon TI, Manaster I, Gazit R, Yutkin V, Benharroch D, Porgador A, Keshet E, Yagel S, Mandelboim O. Decidual NK cells regulate key developmental processes at the human fetal-maternal interface. *Nat Med.* 2006;12:1065–1074.
- Huang SJ, Chen CP, Schatz F, Rahman M, Abrahams VM, Lockwood CJ. Pre-eclampsia is associated with dendritic cell recruitment into the uterine decidua. *J Pathol.* 2008;214:328–336.
- Halhali A, Diaz L, Avila E, Ariza AC, Garabedian M, Larrea F. Decreased fractional urinary calcium excretion and serum 1,25-dihydroxyvitamin D and IGF-I levels in preeclampsia. *J Steroid Biochem Mol Biol.* 2007;103(3–5):803–6.
- Halhali A, Villa AR, Madrazo E, Soria MC, Mercado E, Diaz L, Avila E, Garabedian M, Larrea F. Longitudinal changes in maternal serum 1,25-dihydroxyvitamin D and insulin like growth factor I levels in pregnant women who developed preeclampsia: comparison with normotensive pregnant women. *J Steroid Biochem Mol Biol.* 2004;89-90(1–5):553–6.
- Haugen M, Brantsaeter AL, Trogstad L, Alexander J, Roth C, Magnus P, Meltzer HM. Vitamin D supplementation and reduced risk of preeclampsia in nulliparous women. *Epidemiology.* 2009;20(5):720–6.
- Hofmeyr GJ, Lawrie TA, Atallah AN, Duley L: Calcium supplementation during pregnancy for preventing hypertensive disorders and related problems (Version 3). *Cochrane Database Syst Rev.* 2010;(8):1-87.
- Hofmeyr GJ, Duley L, Atallah A. Dietary calcium supplementation for prevention of pre-eclampsia and related problems: a systematic review and commentary. *BJOG.* 2007;114(8):933–43.
- Hobbs RD, Habib Z, Alromaihi D, Idi L, Parikh N, Blocki F, Rao DS. Severe vitamin D deficiency in Arab-American women living in Dearborn, Michigan. *Endocrine Pract.* 2009;15(1):35–40.
- Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr.* 2004;80(6 Suppl):1678S–88S.
- Hollis BW, Wagner CL. Assessment of dietary vitamin D requirements during pregnancy and lactation. *Am J Clin Nutr.* 2004;79(5):717–26.
- Halhali A, Tovar AR, Torres N, Bourges H, Garabedian M, Larrea F. Preeclampsia is associated with low circulating levels of insulin-like

- growth factor I and 1,25-dihydroxyvitamin D in maternal and umbilical cord compartments. *J Clin Endocrinol Metab.* 2000;85(5):1828–33.
- IOM (Institute of Medicine): Dietary Reference Intakes for Calcium and Vitamin D. Washington, DC; 2011.
- Irani RA, Xia Y. The functional role of the renin-angiotensin system in pregnancy and preeclampsia. *Placenta.* 2008;29(9):763–71.
- Irgens HU, Reisæter L, Irgens LM, Lie RT. Long term mortality of mothers and fathers after pre-eclampsia: population based cohort study. *Br Med J* 2001; 323:1213 – 1216.
- Jeyabalan A, Powers RW, Durica AR, Harger GF, Roberts JM, Ness RB. Cigarette smoke exposure and angiogenic factors in pregnancy and preeclampsia. *Am J Hypertens.* 2008;21(8):943–7.
- Javaid MK, Crozier SR, Harvey NC, Gale CR, Dennison EM, Boucher BJ, Arden NK, Godfrey KM, Cooper C, Princess Anne Hospital Study G. Maternal vitamin D status during pregnancy and childhood bone mass at age 9 years: a longitudinal study. *Lancet.* 2006;367(9504):36–43.
- Jensen CL. Effects of n-3 fatty acids during pregnancy and lactation. *Am J Clin Nutr.* 2006;83(6 Suppl):1452S–7S.
- Kementerian Kesehatan Republik Indonesia. Profil Kesehatan Indonesia Tahun 2015. Jakarta: Kementerian Kesehatan RI:2015
- Khan F, Belch JJ, MacLeod M, Mires G. 2005. Changes in endothelial function precede the clinical disease in women in whom preeclampsia develops. *Hypertension* 46:1123–28
- Keiser SD, Boyd KW, Rehberg JF, Elkins S, Owens MY, Sunesara I, et al.  
A high LDH to AST ratio helps to differentiate pregnancy-associated thrombotic thrombocytopenic purpura (TTP) from HELLP syndrome. *J Matern Fetal Neonatal Med* 2012;25:1059–63.
- Koga K, Osuga Y, Tajima T, Hirota Y, Igarashi T, et al. 2009. Elevated serum soluble fms-like tyrosine kinase 1 (sFlt1) level in women with hydatidiform mole. *Fertil. Steril.* In press
- Kuo VS, Koumantakis G, Gallery ED. Proteinuria and its assessment in normal and hypertensive pregnancy. *Am J Obstet Gynecol* 1992;167:723–8.
- Karumanchi SA, Levine RJ. How does smoking reduce the risk of



- preeclampsia? *Hypertension*. 2010;55:1100–1101.
- Kanasaki K, Palmsten K, Sugimoto H, Ahmad S, Hamano Y, Xie L, Parry S, Augustin HG, Gattone VH, Folkman J, Strauss JF, Kalluri R. Deficiency in catechol-O-methyltransferase and 2-methoxyoestradiol is associated with pre-eclampsia. *Nature*. 2008;453:1117–1121.
- Kovacs CS. Vitamin D in pregnancy and lactation: maternal, fetal, and neonatal outcomes from human and animal studies. *Am J Clin Nutr*. 2008;88(2):520S–8S. 31.
- Kristal-Boneh E, Fromm P, Harari G, Ribak J. Association of calcitriol and blood pressure in normotensive men. *Hypertension*. 1997;30(5):1289–94.
- Knight M, Redman CW, Linton EA, Sargent IL. Shedding of syncytiotrophoblast microvilli into the maternal circulation in pre-eclamptic pregnancies. *Br J Obstet Gynaecol*. 1998;105:632–640.
- Kamen DL, Tangpricha V. Vitamin D and molecular actions on the immune system: modulation of innate and autoimmunity. *J Mol Med*. 2010;88(5):441–50.
- Kaufmann P, Black S, Huppertz B. Endovascular trophoblast invasion: implications for the pathogenesis of intrauterine growth retardation and preeclampsia. *Biol Reprod*. 2003;69(1):1–7. 10.
- Lam C, Lim KH, Karumanchi SA. Circulating angiogenic factors in the pathogenesis and prediction of preeclampsia. *Hypertension*. 2005;46: 1077–1085
- Lisonkova S, Joseph KS. Incidence of preeclampsia: risk factors and outcomes associated with early- versus late-onset disease. *Am J Obstet*
- Langer B, Grima M, Coquard C, Bader AM, Schlaeder G, Imbs JL. Plasma active renin, angiotensin I, and angiotensin II during pregnancy and in preeclampsia. *Obstet Gynecol*. 1998;91(2):196–202.
- YC, Qiao G, Uskokovic M, Xiang W, Zheng W, Kong J. Vitamin D: a negative endocrine regulator of the renin-angiotensin system and blood pressure. *J Steroid Biochem Mol Biol*. 2004;89-90(1–

- 5):387–92.
- Lechtermann C, Hauffa BP, Herrmann R, Schundeln MM, Gellhaus A, Schmidt M, Grasmann C. Maternal vitamin D status in preeclampsia: seasonal changes are not influenced by placental gene expression of vitamin D metabolizing enzymes. *PLoS One*.2014;9(8):e105558.
- Lockwood CJ, Matta P, Krikun G, Koopman LA, Masch R, Toti P, Arcuri F, Huang ST, Funai EF, Schatz F. Regulation of monocyte chemoattractant protein-1 expression by tumor necrosis factor-alpha and interleukin-1beta in first trimester human decidual cells: implications for preeclampsia. *Am J Pathol*. 2006;168:445– 452.
- Lok CA, Boing AN, Sargent IL, Sooranna SR, van der Post JA, Nieuwland R, Sturk A. Circulating platelet-derived and placenta-derived microparticles expose Flt-1 in preeclampsia. *Reprod Sci*. 2008;15: 1002–1010.
- Lachmeijer AM, Arngrimsson R, Bastiaans EJ, Frigge ML et al. (2001) A genome-wide scan for preeclampsia in the Netherlands. *Eur J Hum Genet* 9, 758-764.
- Laivuori H, Lahermo P, Ollikainen V, Widen E et al. (2003) Susceptibility loci for preeclampsia on chromosomes 2p25 and 9p13 in Finnish families. *Am J Hum Genet* 72, 168-177.
- Lie RT, Rasmussen S, Brunborg H, Gjessing HK et al. (1998) Fetal and maternal contributions to risk of pre-eclampsia: population based study. *Bmj* 316, 1343-1347.
- Liston WA and Kilpatrick DC (1991) Is genetic susceptibility to pre-eclampsia conferred by homozygosity for the same single recessive gene in mother and fetus? *Br J Obstet Gynaecol* 98, 1079-1086.
- Lopez-Llera M, Horta JLH. Pregnancy after eclampsia. *Am J ObstetGynecol* 1974; 119:193 – 198.
- Levine RJ, Maynard SE, Qian C, Lim KH, England LJ, et al. 2004.Circulating angiogenic factors and the risk of preeclampsia. *N. Engl. J. Med.* 350:672–83
- Lafayette RA, Druzin M, Sibley R, Derby G, Malik T, et al. 1998. Nature of glomerular dysfunction in pre-eclampsia. *Kidney Int.* 54:1240–49
- MaharajAS,WalshTE,SaintGeniezM,VenkateshaS,MaldonadoAE,etal.2008.VEGFandTGF-β are required for the maintenance of the choroid plexus and ependyma. *J. Exp. Med.* 205:491–501
- Meekins JW, Pijnenborg R, Hanssens M, McFadyen IR, van Asshe

- A.1994. A study of placental bed spiral arteries and trophoblast invasion in normal and severe pre-eclamptic pregnancies. *Br. J. Obstet. Gynaecol.* 101:669–74
- Matsuo K, Kooshesh S, Dinc M, Sun CC, Kimura T, Baschat AA. 2007. Late postpartum eclampsia: report of two cases managed by uterine curettage and review of the literature. *Am. J. Perinatol.* 24:257–66
- Maynard SE, Min JY, Merchan J, Lim KH, Li J, et al. 2003. Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may contribute to endothelial dysfunction, hypertension, and proteinuria in preeclampsia. *J. Clin. Investig.* 111:649–58
- MacKay AP, Berg CJ, Atrash HK. Pregnancy-related mortality from preeclampsia and eclampsia. *Obstet Gynecol* 2001; 97:533–538.
- Moses EK, Fitzpatrick E, Freed KA, Dyer TD et al. (2006) Objective prioritization of positional candidate genes at a quantitative trait locus for pre-eclampsia on 2q22. *Mol Hum*
- Mirzakhani H, Litonjua AA, McElrath TF, O'Connor G, Lee-Parritz A, Iverson R, Macones G, Strunk RC, Bacharier LB, Zeiger R, et al. Early pregnancy vitamin D status and risk of preeclampsia. *J Clin Invest.* 2016;126(12):4702–15.
- Magnus P, Eskild A. Seasonal variation in the occurrence of pre-eclampsia. *BJOG.* 2001;108(11):1116–9.
- Mendel CM. The free hormone hypothesis: a physiologically based mathematical model. *Endocr Rev.* 1989;10(3):232–74.
- March KM, Chen NN, Karakochuk CD, Shand AW, Innis SM, von Dadelszen P, Barr SI, Lyon MR, Whiting SJ, Weiler HA, et al. Maternal vitamin D(3) supplementation at 50 mug/d protects against low serum 25- hydroxyvitamin D in infants at 8 wk of age: a randomized controlled trial of 3 doses of vitamin D beginning in gestation and continued in lactation. *Am J Clin Nutr.* 2015;102(2):402–10.
- Marcoux S, Brisson J, Fabia J. The effect of cigarette smoking on the risk of preeclampsia and gestational hypertension. *Am J Epidemiol.* 1989;130(5):950–7.
- Mohaghegh Z, Abedi P, Dilgouni T, Namvar F, Ruzafza S. The relation of preeclampsia and serum level of 25-hydroxyvitamin D in mothers and their neonates: a case control study in Iran. *Horm Metab Res.* 2015;47(4):284–8.

- Merke J, Hofmann W, Goldschmidt D, Ritz E. Demonstration of 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub> receptors and actions in vascular smooth muscle cells in vitro. *Calcif Tissue Int.* 1987;41(2):112–4.
- Merke J, Milde P, Lewicka S, Hugel U, Klaus G, Mangelsdorf DJ, Haussler MR, Rauterberg EW, Ritz E. Identification and regulation of 1,25- dihydroxyvitamin D<sub>3</sub> receptor activity and biosynthesis of 1,25- dihydroxyvitamin D<sub>3</sub>. Studies in cultured bovine aortic endothelial cells and human dermal capillaries. *J Clin Invest.* 1989;83(6):1903–15.
- Merchant SJ, Narumiya H, Zhang Y, Guilbert LJ, Davidge ST. The effects of preeclampsia and oxygen environment on endothelial release of matrix metalloproteinase-2. *Hypert Pregnancy.* 2004;23(1):47–60.
- Ma R, Gu Y, Zhao S, Sun J, Groome LJ, Wang Y. Expressions of vitamin D metabolic components VDBP, CYP2R1, CYP27B1, CYP24A1, and VDR in placentas from normal and preeclamptic pregnancies. *Am J Physiol Endocrinol Metab.* 2012;303(7):E928–35.
- Marya RK, Rathee S, Manrow M. Effect of calcium and vitamin D supplementation on toxemia of pregnancy. *Gynecol Obstet Investig.* 1987;24(1):38–42.
- McIntire, D.D.; Bloom, S.L.; Casey, B.M.; Leveno, K.J. Birth weight in relation to morbidity and mortality among newborn infants. *N. Engl. J. Med.* 1999, 340, 1234–1238.
- MacKay AP, Berg CJ, Atrash HK. Pregnancy-related mortality from preeclampsia and eclampsia. *Obstet Gynecol.* 2001;97(4):533–8.
- Nesby-O'Dell S, Scanlon KS, Cogswell ME, Gillespie C, Hollis BW, Looker AC, Allen C, Dougherty C, Gunter EW, Bowman BA. Hypovitaminosis D prevalence and determinants among African American and white women of reproductive age: third National Health and nutrition examination survey, 1988-1994. *Am J Clin Nutr.* 2002;76(1):187–92.
- Noyola-Martinez N, Diaz L, Avila E, Halhali A, Larrea F, Barrera D. Calcitriol downregulates TNF-alpha and IL-6 expression in cultured placental cells from preeclamptic women. *Cytokine.* 2013;61(1):245–50.
- Nagamatsu T, Fujii T, Kusumi M, Zou L, Yamashita T, Osuga Y, Momoeda M, Kozuma S, Taketani Y. Cytotrophoblasts up-regulate soluble fms- like tyrosine kinase-1 expression under reduced oxygen: an implication for the placental vascular development

- and the pathophysiology of preeclampsia. *Endocrinology*. 2004;145:4838–4845.
- North RA, Ferrier C, Long D, Townend K, Kincaid Smith P. 1994. Uterine artery Doppler flow velocity waveforms in the second trimester for the prediction of preeclampsia and fetal growth retardation. *Obstet. Gynecol.* 83:378–86
- Ozcan C, Wong SJ, Hari P. 2006. Reversible posterior leukoencephalopathy syndrome and bevacizumab. *N. Engl. J. Med.* 354:980–82
- Opsjon SL, Wathen NC, Tingsulstad S, Wiedswang G, Sundan A, Waage A, Austgulen R. TNF, IL-1 and IL-6 in normal human pregnancy. *Am J Obstet Gynecol* 1993; 169:397–404.
- Olsen SF, Secher NJ, Tabor A, Weber T, Walker JJ, Gluud C. Randomised clinical trials of fish oil supplementation in high risk pregnancies. Fish oil trials in pregnancy (FOTIP) team. *BJOG*. 2000;107(3):382–95.
- Olsen SF, Secher NJ. A possible preventive effect of low-dose fish oil on early delivery and pre-eclampsia: indications from a 50-year-old controlled trial. *Br J Nutr.* 1990;64(3):599–609.
- Pedoman Nasional Pelayanan Kedokteran. Diagnosis dan Tatalaksana Preeklampsia. PNPk. Jakarta. 2016.
- Prawirohardjo, S. 2014. Ilmu Kebidanan. Edisi Keempat ed. Jakarta: P.T. Bina Pustaka Sarwono Prawirohardjo.
- Poston L, Briley AL, Seed PT, Kelly FJ, Shennan AH. Vitamin C and vitamin E in pregnant women at risk for pre-eclampsia (VIP trial): randomised placebo-controlled trial. *Lancet.* 2006;367:1145–1154
- Peters RM, Flack JM. Hypertensive disorders of pregnancy. *J Obstet Gynecol Neonatal Nurs.* 2004;33(2):209–20.
- Pena HR, de Lima MC, Brandt KG, de Antunes MM, da Silva GA. Influence of preeclampsia and gestational obesity in maternal and newborn levels of vitamin D. *BMC Pregnancy Childbirth.* 2015;15:112.
- Powe CE, Seely EW, Rana S, Bhan I, Ecker J, Karumanchi SA, Thadhani R. First trimester vitamin D, vitamin D binding protein, and subsequent preeclampsia. *Hypertension.* 2010;56(4):758–63.
- Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, et al. Recommendations for blood pressure measurement in humans and experimental animals: Part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of

- Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Hypertension* 2005;45:142–61.
- Roberts JM, Taylor RN, Musci TJ, Rodgers GM, Hubel CA, McLaughlin MK. 1989. Preeclampsia: an endothelial cell disorder. *Am. J. Obstet. Gynecol.* 161:1200–4 Annu. Rev. Pathol. Mech. Dis. 2010.5:173-192. Downloaded from www.annualreviews.org by HINARI on 11/09/11. For personal use only.
- Redman CW, Sacks GP, Sargent IL. 1999. Preeclampsia: an excessive maternal inflammatory response to pregnancy. *Am. J. Obstet. Gynecol.* 180:499–506
- Roberts JM, Lain KY. 2002. Recent insights into the pathogenesis of pre-eclampsia. *Placenta* 23:359–72
- Roberts JM, Gammill HS. 2005. Preeclampsia: recent insights. *Hypertension* 46:1243–49
- Rodgers GM, Taylor RN, Roberts JM. 1988. Preeclampsia is associated with a serum factor cytotoxic to human endothelial cells. *Am. J. Obstet. Gynecol.* 159:908–14
- Report of the National High Blood Pressure Education Program Working Group on High Blood Pressure in Pregnancy. *Am J Obstet Gynecol* 2000;183:S1–S22.
- Roberts JM. Pre-eclampsia and cardiovascular disease later in life: who is at risk? *Br Med J* 2001; 323:1217.
- Report of the National High Blood Pressure Education Program Working Group on High Blood Pressure in Pregnancy. *Am J Obstet Gynecol* 2000; 183:S1–S22.
- Reprod 12, 505-512. Moses EK, Lade JA, Guo G, Wilton AN et al. (2000) A genome scan in families from Australia and New Zealand confirms the presence of a maternal susceptibility locus for pre-eclampsia, on chromosome 2. *Am J Hum Genet* 67, 1581-1585.
- Redman CW, Sargent IL. Latest advances in understanding pre-eclampsia. *Science.* 2005;308:1592–1594.
- Rumbold AR, Crowther CA, Haslam RR, Dekker GA, Robinson JS. Vitamins C and E and the risks of preeclampsia and perinatal complications. *N Engl J Med.* 2006;354:1796–1806.

- Raijmakers MT, Dechend R, Poston L. Oxidative stress and pre-eclampsia: rationale for antioxidant clinical trials. *Hypertension*. 2004;44:374–380.
- Roberts JM, Myatt L, Spong CY, Thom EA, Hauth JC, Leveno KJ, Pearson GD, Wapner RJ, Varner MW, Thorp JM Jr, Mercer BM, Peaceman AM, Ramin SM, Carpenter MW, Samuels P, Sciscione A, Harper M, Smith WJ, Saade G, Sorokin Y, Anderson GB. Vitamins C and E to prevent complications of pregnancy-associated hypertension. *N Engl J Med*. 2010;362:1282–1291.
- Rinsler MG, Rigby B. Function of aldosterone in the metabolism of sodium and water in pregnancy. *BMJ*. 1957;2:966–969.
- Rosario GX, Konno T, Soares MJ. Maternal hypoxia activates endothelial trophoblast cell invasion. *Dev Biol*. 2008;314:362–375. 131.
- Ritchie LD, Fung EB, Halloran BP, Turnlund JR, Van Loan MD, Cann CE, King JC. A longitudinal study of calcium homeostasis during human pregnancy and lactation and after resumption of menses. *Am J Clin Nutr*. 1998;67(4): 693–701.
- Resnick LM, Nicholson JP, Laragh JH. Calcium metabolism in essential hypertension: relationship to altered renin system activity. *Fed Proc*. 1986;45(12):2739–45.
- Reddy GS, Norman AW, Willis DM, Goltzman D, Guyda H, Solomon S, Philips DR, Bishop JE, Mayer E. Regulation of vitamin D metabolism in normal human pregnancy. *J Clin Endocrinol Metab*. 1983;56(2):363–70.
- Robinson CJ, Alanis MC, Wagner CL, Hollis BW, Johnson DD. Plasma 25-hydroxyvitamin D levels in early-onset severe preeclampsia. *Am J Obstet Gynecol*. 2010;203(4):366 e361–6.
- Roberts JM, Hubel CA. The two stage model of preeclampsia: variations on the theme. *Placenta*. 2009;30(Suppl A):S32–7.
- Roberts JM, Balk JL, Bodnar LM, Belizan JM, Bergel E, Martinez A. Nutrient involvement in preeclampsia. *J Nutr*. 2003;133(5 Suppl 2):1684S–92S.
- Sibai B, Dekker G, Kupferminc M. Pre-eclampsia. *Lancet* 2005, 365,785–799.
- Sibai BM. Maternal and Uteroplacental Hemodynamics for the Classification and Prediction of Preeclampsia. *Journal of the American Heart Association. Hypertension*. 2008; 52;805-806.

- Sibai BM. Diagnosis, prevention, and management of eclampsia. *Obstet Gynecol.* 2005;105(2):402–10.
- Srinivas SK, Edlow AG, Neff PM, Sammel MD, Andrela CM, Elovitz MA. Rethinking IUGR in preeclampsia: dependent or independent of maternal hypertension? *J Perinatol: official journal of the California Perinatal Association.* 2009;29(10):680–4. 9.
- Smith GC, Pell JP, Walsh D. Pregnancy complications and maternal risk of ischaemic heart disease: a retrospective cohort study of 129,290 births. *Lancet.* 2001;357(9273):2002–6. 13. Redman CW. Immunological aspects of pre-eclampsia. *Baillieres Clin Obstet Gynaecol.* 1992;6(3):601–15. 14. Wei SQ,
- Sachan A, Gupta R, Das V, Agarwal A, Awasthi PK, Bhatia V. High prevalence of vitamin D deficiency among pregnant women and their newborns in northern India. *Am J Clin Nutr.* 2005;81(5):1060–4.32.
- Shand AW, Nassar N, Von Dadelszen P, Innis SM, Green TJ. Maternal vitamin D status in pregnancy and adverse pregnancy outcomes in a group at high risk for pre-eclampsia. *BJOG.* 2010;117(13):1593–8.
- Skull SA, Ngeow JY, Biggs BA, Street A, Ebeling PR. Vitamin D deficiency is common and unrecognized among recently arrived adult immigrants from the horn of Africa. *Intern Med J.* 2003;33(1–2):47–51.
- Sanchez PA, Idrisa A, Bobzom DN, Airede A, Hollis BW, Liston DE, Jones DD, Dasgupta A, Glew RH. Calcium and vitamin D status of pregnant teenagers in Maiduguri, Nigeria. *J Natl Med Assoc.* 1997;89(12):805–11.
- Seely EW, Brown EM, DeMaggio DM, Weldon DK, Graves SW. A prospective study of calciotropic hormones in pregnancy and post partum: reciprocal changes in serum intact parathyroid hormone and 1,25-dihydroxyvitamin D. *Am J Obstet Gynecol.* 1997;176(1 Pt 1):214–7.
- Sankaralingam S, Arenas IA, Lalu MM, Davidge ST. Preeclampsia: current understanding of the molecular basis of vascular dysfunction. *Expert Rev Mol Med.* 2006;8(3):1–20.
- Stumpf WE, O'Brien LP. 1,25 (OH)<sub>2</sub> vitamin D<sub>3</sub> sites of action in the brain. An autoradiographic study. *Histochemistry.* 1987;87(5):393–406.
- Scholl TO, Chen X, Stein TP. Vitamin D, secondary hyperparathyroidism, and preeclampsia. *Am J Clin Nutr.* 2013;98(3):787–93.



- Singla R, Gurung P, Aggarwal N, Dutta U, Kochhar R. Relationship between preeclampsia and vitamin D deficiency: a case control study. *Arch Gynecol Obstet*. 2015;291(6):1247–51.
- Shakiba M, Iranmanesh MR. Vitamin D requirement in pregnancy to prevent deficiency in neonates: a randomised trial. *Singap Med J*. 2013;54(5):285–8.
- Schneuer FJ, Roberts CL, Guilbert C, Simpson JM, Algert CS, Khambalia AZ, Tasevski V, Ashton AW, Morris JM, Nassar N. Effects of maternal serum 25- hydroxyvitamin D concentrations in the first trimester on subsequent pregnancy outcomes in an Australian population. *Am J Clin Nutr*. 2014; 99(2):287–95.
- Signore C, Mills JL, Qian C, Yu K, Lam C, Epstein FH, Karumanchi SA, Levine RJ. Circulating angiogenic factors and placental abruption. *Obstet Gynecol*. 2006;108:338 –344.
- Signore C, Mills JL, Qian C, Yu KF, Rana S, Karumanchi SA, Levine RJ. Circulating soluble endoglin and placental abruption. *Prenat Diagn*. 2008;28:852– 858.
- Spinnato JA II, Freire S, Pinto ESJL, Cunha Rudge MV, Martins-Costa S, Koch MA, Goco N, Santos Cde B, Cecatti JG, Costa R, Ramos JG, Moss N, Sibai BM. Antioxidant therapy to prevent preeclampsia: a randomized controlled trial. *Obstet Gynecol*. 2007;110:1311–1318.
- Skjaerven R, Vatten LJ, Wilcox AJ, Ronning T et al. (2005) Recurrence of pre-eclampsia across generations: exploring fetal and maternal genetic components in a population based cohort. *Bmj* 331, 877.
- Sutherland A, Cooper DW, Howie PW, Liston WA et al. (1981) The incidence of severe pre- eclampsia amongst mothers and mothers- in-law of pre-eclamptics and controls. *Br J Obstet Gynaecol* 88, 785-791.
- Sibai BM, Mercer B, Sarinoglu C. Severe preeclampsia in the second trimester: recurrence risk and long-term prognosis. *Am J Obstet Gynecol* 1991; 165(5 Pt. 1): 1408 – 1412.
- Sibai BM, el Nazer A, Gonzalez-Ruiz A. Severe preeclampsia-eclampsia in young primigravid women: subsequent pregnancy outcome and remote prognosis. *Am J Obstet Gynecol* 1986; 155(5):1011–1016.
- Sibai BM, Sarinoglu C, Mercer BM. Eclampsia. VII. Pregnancy outcome after eclampsia and long-term prognosis. *Am J Obstet Gynecol* 1992;166(6 Pt. 1):1757– 1761.
- Smith GCS, Pell JP, Walsh D. Pregnancy complications and maternal risk of ischaemic heart disease: a retrospective cohort study of 129290

- births. *Lancet* 2001; 357:2002 – 2006.
- Sibai BM, Mercer BM, Schiff E, Friedman SA. Aggressive versus expectant management of severe preeclampsia at 28 to 32 weeks' gestation: a randomized controlled trial. *Am J Obstet Gynecol* 1994;171:818–822.
- Stone P, Cook D, Hutton J, Purdie G, Murray H, Harcourt L. Measurements of blood pressure, oedema and protein-uria in a pregnant population of New Zealand. *Aust N Z J Obstet Gynaecol* 1995;35:32–7
- Sibai BM. 2005. Diagnosis, prevention, and management of eclampsia. *Obstet. Gynecol.* 105:402–10
- Shembrey MA, Noble AD. 1995. An instructive case of abdominal pregnancy. *Aust. N.Z. J. Obstet. Gynaecol.* 35:220–1
- Salafia CM, Pezzullo JC, Ghidini A, Lopez-Zeno JA, Whittington SS. 1998. Clinical correlations of patterns of placental pathology in preterm pre-eclampsia. *Placenta* 19:67–72
- Spargo B, Mc CC, Winemiller R. 1959. Glomerular capillary endotheliosis in toxemia of pregnancy. *Arch. Pathol.* 68:593–99
- Tabesh, M., Salehi-Abargouei, A.; 2012. Maternal vitamin D status and risk of pre-eclampsia: a systemic review and meta-analysis. Food security research center, Isfahan University Of Medical Sciences, Isfahan 81745-151, Iran. PubMed-indexed for Medicine.
- Tal R, Shaish A, Barshack I, Polak-Charcon S, Afek A, Volkov A, Feldman B, Avivi C, Harats D. Effects of hypoxia-inducible factor-1alpha overexpression in pregnant mice: possible implications for pre- eclampsia and intrauterine growth restriction. *Am J Pathol.* 2010;177: 2950 –2962.
- Tarcin O, Yavuz DG, Ozben B, Telli A, Ogunc AV, Yuksel M, Toprak A, Yazici D, Sancak S, Deyneli O, et al. Effect of vitamin D deficiency and replacement on endothelial function in asymptomatic subjects. *J Clin Endocrinol Metab.* 2009;94(10):4023–30.
- Thorne-Lyman A, Fawzi WW. Vitamin D during pregnancy and maternal, neonatal and infant health outcomes: a systematic review and meta-analysis. *Paediatr Perinat Epidemiol.* 2012;26(Suppl 1):75–90.
- Ullah MI, Koch CA, Tamanna S, Rouf S, Shamsuddin L. Vitamin D deficiency and the risk of preeclampsia and eclampsia in Bangladesh. *Horm Metab Res.* 2013;45(9):682–7.

- van der Meer IM, Karamali NS, Boeke AJ, Lips P, Middelkoop BJ, Verhoeven I, Wuister JD. High prevalence of vitamin D deficiency in pregnant non-western women in The Hague, Netherlands. *Am J Clin Nutr.* 2006;84(2):350–3. quiz 468-359
- Velloso EP, Vieira R, Cabral AC, Kalapothakis E, Santos RA. Reduced plasma levels of angiotensin-(1-7) and renin activity in preeclamptic patients are associated with the angiotensin I- converting enzyme deletion/deletion genotype. *Brazilian J Med Biol Res.*2007;40(4):583–90.
- Vaisbuch E, Whitty J, Hassan S, Romero R, Kusanovic J, Cotton D, Sorokin Y, Karumanchi SA. Circulating angiogenic and anti-angiogenic factors in women with eclampsia. *Am J Obstet Gynecol.*2011;204: 152.e1–152.e9.
- Villar J, Purwar M, Merialdi M, Zavaleta N, Thi Nhu Ngoc N, Anthony J, De Greeff A, Poston L, Shennan A. World Health Organisation multicentre randomised trial of supplementation with vitamins C and E among pregnant women at high risk for pre-eclampsia in populations of low nutritional status from developing countries. *BJOG.* 2009;116: 780 –788.
- Vince GS, Starkey PM, Austgulen R, Kwiatowski D, Redman CWG. Interleukin-6, tumour necrosis factor and soluble tumour necrosis factor in women with pre- eclampsia. *Br J Obstet Gynaecol* 1995;
- 1 Magee LA, Ornstein MP, von Dadelszen P. Fortnightly review: management of hypertension in pregnancy. *Br Med J* 1999;318:1332–1336. 02:20–25.
- von Dadelszen P, Magee LA, Lee SK, Stewart SD, Simone C, Koren G, Walley KR, Russell JA. Activated protein C in normal human pregnancy and pregnancies complicated by severe preeclampsia: a therapeutic opportunity? *Crit Care Med* 2002; 30:1883–1892.
- Venkatesha S, Toporsian M, Lam C, Hanai J, Mammoto T, et al. 2006.  
Soluble endoglin contributes to the pathogenesis of preeclampsia. *Nat. Med.* 12:642–49
- WHO. Maternal deaths fell 44% since 1990-UN: Report from WHO, UNICEF, UNFPA, World Bank Group and the United NATIONS POPULATION Division highlights progress 2015. Tersedia dari: <http://www.who.int/mediacentre/news/releases/2015/maternal->

[mortality/en/](#).

- Wulan sari. Hubungan kenaikan berat badan ibu bersalin selama hamil dengan kejadian preeklampsia di Rumah Sakit Umum Daerah Panembahan Senapati. Tahun 2019.
- Wheeler TL 2nd, Blackhurst DW, Dellinger EH, Ramsey PS. Usage of spot urine protein to creatinine ratios in the evaluation of preeclampsia. *Am J Obstet Gynecol* 2007; 196:465.e1–4.
- Wallukat G, Homuth V, Fischer T, Lindschau C, Horstkamp B, Jupner A, Baur E, Nissen E, Vetter K, Neichel D, Dudenhausen JW, Haller H, Luft FC. Patients with preeclampsia develop agonistic autoantibodies against the angiotensin AT1 receptor. *J Clin Invest*. 1999;103:945–952.
- Wimalasundera RC, Larbalestier N, Smith JH, de Ruiter A, McG Thom SA, Hughes AD, Poulter N, Regan L, Taylor GP. Preeclampsia, anti-retroviral therapy, and immune reconstitution. *Lancet*. 2002;360:1152–1154.
- Weir RJ, Paintin DB, Robertson JL, Tree M, Fraser R, Young J. Renin, angiotensin and aldosterone relationships in normal pregnancy. *Proc R Soc Med*. 1970;63:1101–1102
- Weir RJ, Brown JJ, Fraser R, Kraszewski A, Lever AF, McIlwaine GM, Morton JJ, Robertson JL, Tree M. Plasma renin, renin substrate, angiotensin II, and aldosterone in hypertensive disease of pregnancy. *Lancet*. 1973;1:291–294.
- Walters MR, Wicker DC, Riggle PC. 1,25-Dihydroxyvitamin D3 receptors identified in the rat heart. *J Mol Cell Cardiol*. 1986;18(1):67–72.
- Williams MA, Zingheim RW, King IB, Zebelman AM. Omega-3 fatty acids in maternal erythrocytes and risk of preeclampsia. *Epidemiology*. 1995;6(3):232–7.
- Wetta LA, Biggio JR, Cliver S, Abramovici A, Barnes S, Tita AT. Is midtrimester vitamin D status associated with spontaneous preterm birth and preeclampsia? *Am J Perinatol*. 2014;31(6):541–6.
- Webb AR, Kline L, Holick MF. Influence of season and latitude on the cutaneous synthesis of vitamin D3: exposure to winter sunlight in Boston and Edmonton will not promote vitamin D3 synthesis in human skin. *J Clin Endocrinol Metab*. 1988;67(2):373–8.
- Wei SQ, Audibert F, Hidiroglou N, Sarafin K, Julien P, Wu Y, Luo ZC, Fraser WD. Longitudinal vitamin D status in pregnancy and the risk of pre-eclampsia. *BJOG*. 2012;119(7):832–9.
- Weiler H, Fitzpatrick-Wong S, Veitch R, Kovacs H, Schellenberg J, McCloy U, Yuen CK. Vitamin D deficiency and whole-body and

- femur bone mass relative to weight in healthy newborns. *CMAJ*.2005;172(6):757–61.
- Weisman Y, Harell A, Edelstein S, David M, Spirer Z, Golander A. 1 alpha,25- Dihydroxyvitamin D3 and 24,25-dihydroxyvitamin D3 in vitro synthesis by human decidua and placenta. *Nature*.1979;281(5729):317–9.
- World Health Organization. Fact Sheet - Maternal Mortality. <http://www.who.int/mediacentre/factsheets/fs348/en/>. Accessed 23 July 2016.
- Xiong X, Demianczuk NN, Buekens P, Saunders LD. Association of preeclampsia with high birth weight for age. *Am J Obstet Gynecol* 2000; 183:148–155.
- Xiong X, Demianczuk NN, Saunders LD, Wang FL, Fraser WD. Impact of preeclampsia and gestational hypertension on birth weight by gestational age. *Am J Epidemiol* 2002; 155:203–209. Xu L, Lee M, Jeyabalan A, Roberts JM. The relationship of hypovitaminosis D and IL-6 in preeclampsia. *Am J Obstet Gynecol*. 2014;210(2):149 e141–7.
- Young BC, Levine RJ, Karumanchi SA. Pathogenesis of preeclampsia. *Annu Rev Pathol*. 2010;5:173–92.
- Yu CK, Ertl R, Skyfta E, Akolekar R, Nicolaides KH. Maternal serum vitamin D levels at 11-13 weeks of gestation in preeclampsia. *J Hum Hypertens*. 2013;27(2):115–8.
- Zhou Y, Damsky CH, Chiu K, Roberts JM, Fisher SJ. 1993. Preeclampsia is associated with abnormal expression of adhesion molecules by invasive cytotrophoblasts. *J. Clin. Investig.* 91:950–60
- Zhou Y, Fisher SJ, Janatpour M, Genbacev O, Dejana E, et al. 1997. Human cytotrophoblasts adopt a vascular phenotype as they differentiate. A strategy for successful endovascular invasion? *J. Clin. Investig.* 99:2139–51
- Zhou Y, Damsky CH, Fisher SJ. 1997. Preeclampsia is associated with failure of human cytotrophoblasts to mimic a vascular adhesion phenotype. One cause of defective endovascular invasion in this syndrome? *J. Clin. Investig.* 99:2152–64
- Zhou Y, McMaster M, Woo K, Janatpour M, Perry J, et al. 2002. Vascular endothelial growth factor ligands and receptors that regulate human cytotrophoblast survival are dysregulated in severe preeclampsia and hemolysis, elevated liver enzymes, and

- low platelets syndrome. *Am. J. Pathol.* 160:1405–23
- Zhou J, Su L, Liu M, Liu Y, Cao X, Wang Z, Xiao H. Associations between 25-hydroxyvitamin D levels and pregnancy outcomes: a prospective observational study in southern China. *Eur J Clin Nutr.* 2014;68(8):925–30.
- Zhou CC, Zhang Y, Irani RA, Zhang H, Mi T, Popek EJ, Hicks MJ, Ramin SM, Kellems RE, Xia Y. Angiotensin receptor agonistic autoantibodies induce pre-eclampsia in pregnant mice. *Nat Med.* 2008;14: 855– 862.
- Zhou A, Carrell RW, Murphy MP, Wei Z, Yan Y, Stanley PL, Stein PE, Broughton Pipkin F, Read RJ. A redox switch in angiotensinogen modulates angiotensin release. *Nature.* 2010;468:108–111.
- Zhang J, Zeisler J, Hatch MC, Berkowitz G. Epidemiology of pregnancy-induced hypertension. *Epidemiol Rev.* 1997;19(2):218–32.