

## DAFTAR PUSTAKA

- Akhundov, K. *et al.* (2012) ‘Development of a cost-effective method for platelet-rich plasma (prp) preparation for topical wound healing’, *Annals of Burns and Fire Disasters*, 25(4), pp. 207–213.
- Aleckovic M, Simon C. 2008. Is teratoma formation in stem cell research a characterization tool or a window to developmental biology? *Reprod Biomed Online*, 17(2): 270
- Aslı Duran, ŞirinYaşar, SemaAytekin, et al.,. 2018. Clinical and histopathological evaluation of the effects of platelet rich plasma, platelet poor plasma and topical serum physiologic treatment on wound healing caused by radiofrequency electrosurgery in rats. *Turkderm-Turk Arch Dermatol Venereology* 2018;52:44-50
- Baglioni, S., Francalanci, M., Squecco, R., Lombardi, A., Cantini, G., Angeli, R., Luconi, M. (2009). Characterization of human adult stem-cell populations isolated from visceral and subcutaneous adipose tissue. *The FASEB Journal*, 23(10), 3494–3505. doi:10.1096/fj.08-126946
- Bakacak, M., Bostanci, M. S., İnanc, F., Yaylali, A., Serin, S., Attar, R., Yildirim, O. K. (2015). Protective Effect of Platelet Rich Plasma on Experimental Ischemia/Reperfusion Injury in Rat Ovary. *Gynecologic and Obstetric Investigation*, 81(3), 225–231. doi:10.1159/000440617
- Barret J, Herndon D., 2005. *Principles and Practice of Burn Surgery*. New York, USA. Marcel Dekker.
- Borrione, P., Gianfrancesco, A. D., Pereira, M. T., &Pigozzi, F. (2010). Platelet-Rich Plasma in Muscle Healing. *American Journal of Physical Medicine & Rehabilitation*, 89(10), 854–861.
- Bourin P, Bunnell BA, Casteilla L, Dominici M, Katz AJ, March KL, Redl H, Rubin JP, Yoshimura K, Gimble JM. Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International Society for Cellular Therapy (ISCT). *Cytotherapy*. 2013;15:641–8.
- Braund, R., Hook, S. and Medlicott, N. J. (2007) ‘The role of topical growth factors in chronic wounds.’, *Current drug delivery*. United Arab Emirates, 4(3), pp. 195–204. doi: 10.2174/156720107781023857.
- Cervelli V, et al. (2010). Tissue regeneration in loss of substance on the lower limbs through use of platelet-rich plasma, stem cells from adipose tissue, and hyaluronic acid. *Adv Skin Wound Care*. 2010; 23:262-72.)
- Chierigato K, Castegnaro S, Madeo D, Astori G, Pegoraro M, Rodeghiero F: Epidermal growth factor, basic fibroblast growth factor and platelet-derived growth factor-bb can substitute for fetal bovine serum and compete with human platelet-rich plasma in the ex vivo expansion of mesenchymal stromal cells derived from adipose tissue. *Cytotherapy* 2011, 13:933–943. doi: 10.3109/14653249.2011.583232.

- Choi, J., Minn, K. W., & Chang, H. (2012). The Efficacy and Safety of Platelet-Rich Plasma and Adipose-Derived Stem Cells: An Update. *Archives of Plastic Surgery*, 39(6), 585. doi: 10.5999/aps.2012.39.6.585
- Cohen IK, Degelman RF, Crossland MC. Wound healing and wound care. In: Schwartz SI, ed. *Principles of surgery*, 6 th ed. New York: McGraw-Hill, 1994
- Comella, K., Silbert, R., &Parlo, M. (2017). Effects of the intradiscal implantation of stromal vascular fraction plus platelet rich plasma in patients with degenerative disc disease. *Journal of Translational Medicine*, 15(1).
- Darinskas, A., Paskevicius, M., Apanavicius, G., Vilkevicius, G., Labanauskas, L., Ichim, T. E., & Rimdeika, R. (2017). Stromal vascular fraction cells for the treatment of critical limb ischemia: a pilot study. *Journal of Translational Medicine*, 15(1). doi:10.1186/s12967-017-1243-3
- Deng, C. *et al.* (2018) ‘Treatment of human chronic wounds with autologous extracellular matrix/stromal vascular fraction gel: A STROBE-compliant study’, *Medicine (United States)*, 97(32), pp. 1–7. doi: 10.1097/MD.00000000000011667.
- El-Sharkawy, H., Kantarci, A., Deady, J., Hasturk, H., Liu, H., Alshahat, M., & Van Dyke, T. E. (2007). Platelet-Rich Plasma: Growth Factors and Pro- and Anti-Inflammatory Properties. *Journal of Periodontology*, 78(4), 661–669.
- EMS (Emergency Management of Severe Burns) Course Manual, 17th Edition, 2013. ANZBA Ltd.
- Eppley, B. L., Pietrzak, W. S. and Blanton, M. (2006) ‘Platelet-rich plasma: a review of biology and applications in plastic surgery.’, *Plastic and reconstructive surgery*. United States, 118(6), pp. 147e-159e. doi: 10.1097/01.prs.0000239606.92676.cf.
- Eyuboglu, A. A. *et al.* (2018) ‘The effect of adipose derived stromal vascular fraction on stasis zone in an experimental burn model’, *Burns*. Elsevier Ltd and International Society of Burns Injuries, 44(2), pp. 386–396. doi: 10.1016/j.burns.2017.08.016.
- Ferraro, G. A., Mizuno, H., &Pallua, N. (2016). Adipose Stem Cells: From Bench to Bedside. *Stem Cells International*, 2016, 1–2. doi:10.1155/2016/6484038
- Fotouhi, A. *et al.* (2018) ‘Platelet rich plasma, stromal vascular fraction and autologous conditioned serum in treatment of knee osteoarthritis’, *Biomedicine and Pharmacotherapy*. Elsevier, 104(April), pp. 652–660. doi: 10.1016/j.biopha.2018.05.019
- Fouber, P. *et all.* (2016) ‘Adipose-derived regenerative cell therapy for burn wound healing: A comparison of two delivery methods’, *Advances in Wound Care*, 5(7), pp. 288-298. doi: 10.1089/wound.2015.0672.
- Fu Y-S, Cheng Y-C, Lin M-Y A, Chu P-M, Chou S-C, Shih Y-H, Ko M-H, Sung M-S. (2006). Conversion of Human Umbilical Cord Mesenchymal Stem Cells in Wharton's Jelly to Dopaminergic Neurons In Vitro: Potential Therapeutic Application for Parkinsonism. *Stem Cells*; 24(1):115-124. doi: 10.1634/stemcells.2005-0053.

- Gentile, P., Scioli, M. G., Bielli, A., Orlandi, A., & Cervelli, V. (2016). Concise Review: The Use of Adipose-Derived Stromal Vascular Fraction Cells and Platelet Rich Plasma in Regenerative Plastic Surgery. *STEM CELLS*, 35(1), 117–134. doi: 10.1007/978-1-4939-7799-4\_9.
- Ghieh, F., Jurjus, R., Ibrahim, A., Geagea, A. G., Daouk, H., El Baba, B., ... Jurjus, A. (2015). The Use of Stem Cells in Burn Wound Healing: A Review. *BioMed Research International*, 2015, 1–9. doi:10.1155/2015/684084
- Grigore, T.V. and Cozma, C. (2018). “Platelet Rich-Plasma as a site-tergeted approach in wound healing: a molecular perspective”, *Discoveries*,6(4), pp. 1-8. doi: 10.15190/d.2018.8.
- Gurtner, GC,. (2007). Wound Healing. Normal and Abnormal. In *Grab and Smith's Plastic Surgery 6<sup>th</sup> edition* (pp. 15 – 22). Philadelphia : Elseviers.
- Halim D., Murti H. Sandra F., Boediono A., et al., 2010. Stem Cell dasar teori dan aplikasi klinis. Erlangga press. Jakarta. 1-136p
- Han J, Koh YJ, Moon HR, Ryoo HG, Cho CH, Kim I, Koh GY (2010). Adipose tissue is an extramedullary reservoir for functional hematopoietic stem and progenitor cells. *Blood* 115:957–64. doi: 10.1182/blood-2009-05-219923.
- Harrison, D. A. (2012). The JAK/STAT Pathway. *Cold Spring Harbor Perspectives in Biology*, 4(3), a011205–a011205.
- Hausman, G. J. and Richardson, R. L. (2004) ‘Adipose tissue angiogenesis.’, *Journal of animal science*. United States, 82(3), pp. 925–934. doi: 10.2527/2004.823925x.
- Hu, Z. *et al.* (2009) ‘Platelet-rich plasma induces mRNA expression of VEGF and PDGF in rat bone marrow stromal cell differentiation.’, *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics*. United States, 107(1), pp. 43–48. doi: 10.1016/j.tripleo.2008.06.023.
- Josh F, et al., 2012. Accelerated and safe proliferation of human adipose-derived stem cells in medium supplemented with human serum. Tokyo, Japan. *J Nippon Med* (79): pp. 444-452. edition. Denver, USA. RC Press.
- Kalangi, S.J.R., 2011, *Peran Integrin pada Angiogenesis Penyembuhan Luka*, *Cermin Dunia Kedokteran*, 38(3): 177 – 181.
- Karagergou, E., Dionyssopoulos, A., Karayannopoulou, M., Psalla, D., Theodoridis, A., Demiri, E., & Koliakos, G. (2018). Adipose-derived stromal vascular fraction aids epithelialisation and angiogenesis in an animal model. *Journal of Wound Care*, 27(10), 637–644.
- Karina *et all.* (2019). Combination of the stromal vascular fraction and platelet-rich plasma accelerates the wound healing process: pre-clinical study in a Sprague-Dawley rat model. *Stem Cell Investigation*. AME Publishing Company, 6, p.18. doi: 10.21037/sci.2019.06.08.
- Kazakos, K. *et al.* (2009) ‘The use of autologous PRP gel as an aid in the management of acute trauma wounds’, *Injury*, 40(8), pp. 801–805. doi: 10.1016/j.injury.2008.05.002.

- Keyv S, Jacobson M, Benoit P. The biology of platelet concentrate as prepared by the Harvest Technologies SmartPRP System. Proceedings of the 3rd Annual Meeting of Techvest Conference on Tissue Repair, Replacement, and Regeneration; 2001 Oct 23; New York, USA
- Kim Yeol, et al., 2014. Effects of Platelet-Rich Plasma, Adipose-Derived Stem Cells and Stromal Vascular Fraction on the Survival of Human Transplanted Adipose Tissue. Seoul, Republic of Korea. *J Korean Med Science* (29): pp. 193-200. doi: 10.3346%2Fjkms.2014.29.S3.S193.
- Kuwada, S. K. and Li, X. (2000) ‘Integrin alpha5/beta1 mediates fibronectin-dependent epithelial cell proliferation through epidermal growth factor receptor activation’, *Molecular biology of the cell*. The American Society for Cell Biology, 11(7), pp. 2485–2496. doi: 10.1091/mbc.11.7.2485.
- Lawrence, W. T. (1998) ‘Physiology of the acute wound.’, *Clinics in plastic surgery*. United States, 25(3), pp. 321–340.
- L. Dailey, D. Ambrosetti, A. Mansukhani, and C. Basilico. 2005. “Mechanisms underlying differential responses to FGF signaling,” *Cytokine and Growth Factor Reviews*, vol. 16, no. 2, pp. 233–247. doi: 10.1016/j.cytogfr.2005.01.007.
- Li H, Liu D, Yu Y, Wu T. 2009. Experimental research of the promotion effect of autogeneic PRP on osteogenic differentiation of human adipose-derived stem cells in vitro. *ZhongguoXiu Fu Chong Jian Wai KeZaZhi*: 23:732–736
- Liang, Z. J. *et al.* (2018) ‘Precise Intradermal Injection of Nanofat-Derived Stromal Cells Combined with Platelet-Rich Fibrin Improves the Efficacy of Facial Skin Rejuvenation’, *Cellular Physiology and Biochemistry*, 47(1), pp. 316–329. doi: 10.1159/000489809.
- MacLeod, A. S., & Mansbridge, J. N. (2015). The Innate Immune System in Acute and Chronic Wounds. *Advances in Wound Care*, 5(2), 65–78. doi: 10.1089%2Fwound.2014.0608.
- McGee, G. S. *et al.* (1988) ‘Recombinant basic fibroblast growth factor accelerates wound healing.’, *The Journal of surgical research*. United States, 45(1), pp. 145–153. doi: 10.1016/0022-4804(88)90034-0.
- Moenadjat Y, dkk., 2011. *Luka Bakar: Masalah dan Tatalaksana*, edisi 4. Jakarta, Indonesia. Balai Penerbit FKUI.
- Mohammed, M. O. *et al.* (2019) ‘Role of Bovine Fetal Platelet-rich Plasma (PRP) on Skin Wound Healing in Mice’, *International Journal of Pharmacology*, 16(1), pp. 18– 26. doi: 10.3923/ijp.2020.18.26
- Motamed, S. *et al.* (2017) ‘Cell-based skin substitutes accelerate regeneration of extensive burn wounds in rats’, *American Journal of Surgery*, 214(4), pp. 762–769. doi: 10.1016/j.amjsurg.2017.04.010.
- Park, J., Hwang, S., & Yoon, I.-S. (2017). Advanced Growth Factor Delivery Systems in Wound Management and Skin Regeneration. *Molecules*, 22(8), 1259. doi: 10.3390/molecules22081259.

- Randy, A., (2018). *The Effect of Platelet Rich Plasma (PRP) and Stromal Vascular Fraction Cell (SVFs) and the combination of PRP+SVFs on Burn Wound Healing in the Model of Burn Wound (Grade IIB) of Wistar Rats.*(Tesis). Makassar: Universitas Hasanuddin
- Rantam FA, Ferdiansyah, Nasronudin, Purwati. 2009. Stem Cell Exploration Method of Isolation and Culture. Airlangga University Press, Surabaya
- Raposo E, et al., 2016. Adipose-derived Stem Cells Added to Platelet-Rich Plasma for Chronic Skin Ulcer Therapy. Parma, Italy. *Wound Research Journal* Vol. 28: pp.434-442.
- Rigotti G, Marchi A., 2009. Adipose-derived Mesenchymal Stem Cells: Past, Present and Future. Verona, Italy. *Aesthetic Plastic Surgery Journal* (33): pp. 271–273.
- Rosenstrauch D, Ponglajen G, Zidar N, Gregoric I. 2005. Stem Cell Therapy for Ischemic Heart Failure. *ex Heart Inst J*;32:339-347
- Rumalia VK, Borah GL: Cytokine, growth factor and plastic surgery. *PlastReconstrSurg* 108:719-733, 2001; doi: 10.1097/00006534-200109010-00019.
- Schaffer, C.J.; Nanney, L.B. Cell biology of wound healing. *Int. Rev. Cytol.* 1996, 169, 151–181. doi: 10.1016/S0074-7696(08)61986-5.
- Schöler, H.R. (2016). *The Potential of Stem Cells: An Inventory.*
- Sethi A, Kaur T, Malhotra S K, Gambhir M L. Moisturizers: The slippery road. *Indian J Dermatol* 2016;61:279-87. doi: 10.4103%2F0019-5154.182427.
- Singh, V. K., Saini, A., Kalsan, M., Kumar, N., & Chandra, R. (2016). Describing the Stem Cell Potency: The Various Methods of Functional Assessment and In silico Diagnostics. *Frontiers in Cell and Developmental Biology*, 4. doi:10.3389/fcell.2016.00134.
- Sjamsuhidajat, de Jong., 2016. *Buku Ajar Ilmu Bedah, Sistem Organ dan Tindak Bedahnya.* edisi 4 Vol. 2. Jakarta, Indonesia. EGC.
- Soenarso WS, Rahayu EC, Sriharjo S. 2007. Prospek dan Tinjauan Bioetik Pengembangan Teknologi Kloning Sel Punca untuk Terapi di Bidang Kedokteran. Deputi Bidang Pengembangan Sistem Iptek Nasional. Kementrian Negara Riset dan Teknologi, Jakarta
- Sun, M. *et al.* (2017) ‘Adipose Extracellular Matrix/Stromal Vascular Fraction Gel Secretes Angiogenic Factors and Enhances Skin Wound Healing in a Murine Model’, *BioMed Research International*, 2017. doi: 10.1155/2017/3105780.
- Tajima, S. *et al.* (2014) ‘Direct and Indirect Effects of a Combination of Adipose- Derived Stem Cells and Platelet-Rich Plasma on Bone Regeneration’, *Tissue engineering. Part A*, 21. doi: 10.1089/ten.TEA.2014.0336.
- Tantuway V, et al., Autologus grafting of non manipulated freshly isolated-adipose tissue derived stromal vascular fraction in single surgical sitting for treatment of knee osteoarthritis. *Int J Res Orthop.* 2017 Jan;3(1):107-115
- Tan S., Dosan R., (2019).Lessons From Epithelization: *The Reason Behind Moist Wound Environment.* The Open Dermatology Journal. DOI: 10.2174/1874372201913010034,2019,13,34-40.

- T Velnar, T Bailey, V Smrkolj, (2009). The Wound Healing Process : an Overview of Cellular and Molecular Mechanism, *The J of International Medical Research*, p. 1528 – 42. doi: 10.1177/147323000903700531.
- Vaghardoost, R. *et al.* (2018) ‘The Healing Effect of Sesame Oil, Camphor and Honey on Second Degree Burn Wounds in Rat.’, *World journal of plastic surgery*, 7(1), pp.67–71.
- Van Pham, P., Bui, K. H.-T., Ngo, D., Vu, N., Truong, N., Phan, N. L.-C., Phan, N. (2013). Activated platelet-rich plasma improves adipose-derived stem cell transplantation efficiency in injured articular cartilage. *Stem Cell Research & Therapy*, 4(4), 91.
- Widjajakusumah, Tanzil A., 2014. *Guyton and Hall Buku Ajar Fisiologi Kedokteran*, edisi 12. Singapore. Elsevier.
- Witte MB, Barbul A: General principles of wound healing. *Surg Clin North Am* 77:509–528, 1997.
- Wobus AM, Boheler KR. 2005. Embryonic Stem Cells: Prospects for Developmental Biology and Cell Therapy. *Physiol Rev.* 85: 635–678. doi: 10.1152/physrev.00054.2003.
- Yun, Y.-R., Won, J. E., Jeon, E., Lee, S., Kang, W., Jo, H., ... Kim, H.-W. (2010). Fibroblast Growth Factors: Biology, Function, and Application for Tissue Regeneration. *Journal of Tissue Engineering*, 1(1), 218142.
- Zhang YS, He JH, Xiao GY, Li QM: Effect of platelet-rich plasma on the proliferation and adipogenic differentiation of human adipose-derived stem cells in vitro. *Nan Fang Yi Ke Da XueXue Bao* 2011, 31:525–528.
- Ziegler, T. R., Pierce, G. F., & Herndon, D. N. (Eds.). (1997). *Growth Factors and Wound Healing*. doi:10.1007/978-1-4612-1876-0
- Zuk PA, Zhu M, Ashjian P, De Ugarte DA, Huang JI, Mizuno H, et al. (2002). Human adipose tissue is a source of multi-potent stem cells. *Mol Biol Cell.* 13:4279–95. [PubMed: 12475952]. doi: 10.1091/mbc.e02-02-0105.



KEMENTERIAN PENDIDIKAN DAN KEBUDAYAAN  
UNIVERSITAS HASANUDDIN FAKULTAS KEDOKTERAN  
KOMITE ETIK PENELITIAN KESEHATAN  
RSPTN UNIVERSITAS HASANUDDIN

RSUP Dr. WAHIDIN SUDIROHUSODO MAKASSAR

Sekretariat : Lantai 2 Gedung Laboratorium Terpadu

JL.PERINTIS KEMERDEKAAN KAMPUS TAMALANREA KM.10 MAKASSAR 90245.

Contact Person: dr. Agussalim Bukhari, M.Med.,PhD, SpGK TELP. 081241850858, 0411 5780103, Fax : 0411-581431



**REKOMENDASI PERSETUJUAN ETIK**

Nomor : 1127/UN4.6.4.5.31/ PP36/ 2019

Tanggal: 26 Nopember 2019

Dengan ini Menyatakan bahwa Protokol dan Dokumen yang Berhubungan Dengan Protokol berikut ini telah mendapatkan Persetujuan Etik :

No Protokol	UH19110983	No Sponsor Protokol	
Peneliti Utama	dr. Andi Sinapati Palissei	Sponsor	
Judul Peneliti	Efek Penambahan Platelet-Rich Plasma (PRP) Dan Stromal Vascular Fraction (SVFs) Terhadap Proses Epitelisasi Penyembuhan Pada Luka Bakar Deep Dermal Tikus Wistar		
No Versi Protokol	1	Tanggal Versi	15 Nopember 2019
No Versi PSP		Tanggal Versi	
Tempat Penelitian	Laboratorium Hewan dan Laboratorium Penelitian Fakultas Kedokteran Universitas Hasanuddin Makassar		
Jenis Review	<input type="checkbox"/> Exempted <input checked="" type="checkbox"/> Expedited <input type="checkbox"/> Fullboard Tanggal	Masa Berlaku 26 Nopember 2019 sampai 26 Nopember 2020	Frekuensi review lanjutan
Ketua Komisi Etik Penelitian Kesehatan FKUH	Nama Prof.Dr.dr. Suryani As'ad, M.Sc.,Sp.GK (K)	Tanda tangan	
Sekretaris Komisi Etik Penelitian Kesehatan FKUH	Nama dr. Agussalim Bukhari, M.Med.,Ph.D.,Sp.GK (K)	Tanda tangan	

Kewajiban Peneliti Utama:

- Menyerahkan Amandemen Protokol untuk persetujuan sebelum di implementasikan
- Menyerahkan Laporan SAE ke Komisi Etik dalam 24 Jam dan dilengkapi dalam 7 hari dan Laporan SUSAR dalam 72 Jam setelah Peneliti Utama menerima laporan
- Menyerahkan Laporan Kemajuan (progress report) setiap 6 bulan untuk penelitian resiko tinggi dan setiap setahun untuk penelitian resiko rendah
- Menyerahkan laporan akhir setelah Penelitian berakhir
- Melaporkan penyimpangan dari prokol yang disetujui (protocol deviation / violation)
- Mematuhi semua peraturan yang ditentukan