

DAFTAR PUSTAKA

- Achutan S, Singh I, Varthya SB, Srinivasan A, Chakrabarti A, Dota D. Gabapentin prophylaxis for postoperative nausea and vomiting in abdominal surgeries: a quantitative analysis of evidence from randomized controlled clinical trials. *Brith J Anaesth*. 2015;114(4): 588-97.
- Agustine R, Oktaliansyah E, Surahman E. Perbandingan gabapentin 600 mg dengan pregabalin 150 mg preoperatif terhadap nilai *numeric rating scale* dan pengurangan kebutuhan opioid pascaoperasi modifikasi radikal mastektomi. *J Anest Periop*. 2014; 2(2): 117-23.
- Arbel R, Stanleigh J, Ioscovich A. Pain management following abdominal hysterectomy : novel approaches and review of the literature. *J Clin Gynecol Obstet*. 2013;2(2):51-55.
- Arumugam S, Lau CSM, Chamberlain RS. Use of preoperative gabapentin significantly reduce postoperative opiod consumption: a meta-analysis. *J Pain Res*. 2016;9: 631-40.
- Braga AFA, Frias JAF, Braga FSS, Pereira RIC, Titotto SMMC. Spinal anesthesia for elective ceasarean section. Use of different doses of hyperbaric bupivacaine associated with morphine and clonidine. *Act Cir Bras*. 2013;28(1):23-32.
- Bryson GL, MacNeil R, Jeyaraj LM, Rosaeg OP. Small dose spinal bupivacaine for cesarean delivery does not reduce hypotension but accelerates motor recovery. *Can J Anesth*. 2007;57(7):531-7.
- Butterworth JF, Mackey DC, Wasnick JD. Spinal, epidural & caudal blocks. In : Morgan & mikhail's clinical anesthesiology, 5th Ed. New York: McGraw Hill Education; 2013. p. 937-58.
- Costigan M, Scholz J, Woolf CJ. Neuropathic pain: a maladaptive response of the nervous system to damage. *Annu Rev Neurosci*. 2009; 32:1–32.
- Fabritius ML, Geisler A, Petersen PL, Nikolajsen L, Hansen MS, Kontinen V, et al. Gabapentin for post-operative pain management – a systematic review with meta-analyses and trial sequential analyses. *Act Anaest Scan*. 2016;60: 1188-1208.
- Gottschalk A, Smith DS. New consepts in acute pain therapy: preemptive analgesia. *Am Fam Physic*. 2001;63(1):1979-87.

- Grant MC, Lee HW, Page AJ, Hobson D, Wick E, and Wu CL. The effect of preoperative gabapentin on postoperative nausea and vomiting : a meta- analysis. *Anaesth Analg*. 2016;122(4):976-85.
- Hafez MH, Abdelhamid MH, Youssef MMI, Abdelrahim. Randomized controlled trial of two oral regimens of gabapentin versus placebo in patients for cesarean section under spinal anesthesia regarding postoperative pain, sedation, nausea and vomiting. *Egyp J Anaes*. 2017.; 33: 59-65.
- International Association for the Study of Pain. IASP's proposed new definition of pain released for comment. 2019 Aug 7 [diunduh 22 Februari 2020]; Tersedia dari <http://www.iasp-pain.org>.
- Kandappan G, Raju TS. To evaluate the role of gabapentin as a preemptive analgesic in patients undergoing total abdominal hysterectomy under spinal anaesthesia. *Kour Evid Based Med Healthc*. 2016; 3(49): 2487-91.
- Katz J, Clarke H. Preventive analgesia and beyond: current status, evidence and future direction. In: Macintyre P, editor. *Clinical pain management acute pain*. London : Hodder Arnold Ltd.2008. p.154-98.
- Katzung BG. Obat-obat *antiseizure*. Dalam : *Farmakologi dasar dan klinik*. Jakarta: Salemba Medika; 2002. hal. 104-5.
- Khabi PM, Yaghooti AA, Marashi SH. Effect of pre-emptive gabapentin on postoperative pain following lower extremity orthopedic surgery under spinal anaesthesia. *Sing Med J*. 2011;52(12): 879-82.
- Kjaer K, Kim J. Spinal and epidural blockade for gynecology surgery. In : Wong CA, editor. *Spinal and epidural anesthesia*. New York : McGraw-Hill; 2007. p. 247-56.
- Matthews EA, Dickenson AH. Combination of gabapentin and morphine mediates enhanced inhibitory effects on dorsal horn neuronal responses in a rat model of neuropathy. *Anesthesiology*. 2002; 96(3): 633–40.
- McCracken, G., Houston, P. and Lefebvre, G. Guideline for the management of postoperative nausea and vomiting. *J Obstet Gynaecol Can* 2008; 30(7), 600-7.
- Schmidt PC, Ruchelli G, Mackey SC, Carrol IR. Perioperative gabapentinoids. : choice of agent, dose, timing, and effects on chronic postsurgical pain. *The American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins. Anaesthesiology*. 2013; 119:1215-21.

- Stoelting RK, Hillier SC. Antiepileptic drugs. In : Pharmacology and physiology in anesthetic practice. Philadelphia : Wolters Kluwer Health; 2015. p. 345-63.
- Suhandoko, Pradian E, Maskoen TT. Reliabilitas dan validitas penilaian skala sedasi *richmond agitation sedation scale* (RASS) dan ramsay pada pasien kritis dengan ventilasi mekanik di ruang perawatan intensif. J Anest Periop. 2014; 2(3):186-93.
- Tanra AH, Rehatta NM, Musba AMT. Nyeri pascabedah. Dalam : Penatalaksanaan nyeri. Makassar : Departemen Anestesi, Terapi Intensif, dan Manajemen Nyeri Fakultas Kedokteran Universitas Hasanuddin; 2013. p.44-9.
- Waldman SD. Functional anatomy of the nociceptors. In : Pain review. 2nd ed. Philadelphia: Elsevier; 2009. p.181-2.
- Woolf CJ, Max MB. Mechanism-based pain diagnosis : issues for analgesic drug development. Anesthesiology. 2001;95(1):241-9.
- Yacob MC, Ihsan MH, Wijaya DW. Perbandingan efektivitas antara gabapentin 600 mg dan gabapentin 900 mg kombinasi dengan ketorolak 30 mg/8 jam sebagai analgesia pascabedah pada total abdominal histerektomi dengan anestesi umum. J Anest Periop. 2016; 4(3): 147-53.
- Yilmaz M, Wong CA. Technique of neuraxial anesthesia. In : Wong CA, editor. Spinal and epidural anesthesia. New York : McGraw-Hill; 2007. p. 33-44.
- Zulfariansyah A, Nawawi AM, Bisri T. Perbandingan gabapentin 600 mg dengan 1.200 mg per oral preoperatif terhadap nilai *visual analogue scale* dan pengurangan kebutuhan petidin pascaoperasi pada modifikasi mastektomi radikal. J Anes Perio. 2013;1(3): 174-82.