

DAFTAR PUSTAKA

1. Gibbon L, Belizan JM, Lauer JA, Betran AP, Merialdi M, Althabe F. The global numbers and costs of additionally needed and unnecessary caesarean sections performed per year: overuse as a barrier to universal coverage [Internet]. Vol 30, World health report. Geneva, Switzerland; 2010. 1-31 p. Available from: <http://www.who.int/healthsystems/topics/financing/healthreport/30C-sectioncost.pdf>
2. Badan Pusat Statistik, BKKBN, Kementerian Kesehatan, Macro Inc. Laporan Survei Demografi dan Kesehatan Indonesia 2007. BKKBN, Departemen Kesehatan, Macro Calverton Mary Land. 2008.
3. Kementerian Kesehatan. Laporan Nasional Riset Kesehatan Dasar 2013 [Internet]. Jakarta: Badan Litbang Kesehatan; 2013. Available from: <http://www.litbang.depkes.go.id/sites/download/rkd2013/LaporanRisksdas2013.pdf>
4. Kementerian Kesehatan. Laporan Nasional Riset Kesehatan Dasar 2018 [Internet]. Jakarta: Badan Litbang Kesehatan; 2018. Available from: <http://www.litbang.depkes.go.id/sites/download/rkd2018/LaporanRisksdas2018.pdf>
5. Roofthoof E, Joshi G.P, Rawal N, Van de Velde M. PROSPECT guideline for elective caesarean section: update systematic review and procedure-specific postoperative pain management recommendations. *Anesthesia* 2021. 2020 Nov. 76: 665-680.
6. PROSPECT. Recommendations for postoperative analgesia after caesarean section 2014. www.postoppain.org (accessed 4/11/2020).
7. Southworth S, Peters J, Rock Amy, Pavliv L. A Multicentre, Randomized, Double-Blind, Placebo-Controlled Trial of Intravenous Ibuprofen 400 and 800 mg every 6 Hours in the Management of Postoperative Pain. *Clinical Therapeutics*. 2009. Vol 31. No 9. 1922-1935.
8. Thybo K, Pederson D, Dahl J, Nersesjan M, Jakobsen J, Pedersos N, et al. Effect of Combination of Paracetamol (Achetaminophen) and Ibuprofen vs

- Either alone on Patient-Controlled Morphine Consumption in the First 24 Hours After Total Hip Arthroplasty The PANSOID Randomized Clinical Trial. *JAMA*. 2019; 321(6):562-571.
9. Apfelbaum JL, Chen C, Mehta SS, Gan TJ. Postoperative pain experience: results from a national survey suggest postoperative pain continues to be undermanaged. *Anesth Analg*. 2003;9:534-40.
 10. Thota RS, Ramkiran S, Garg R, Goswami J, Baxi V, Thomas M. Opioid free onco-anesthesia: Is it time to convict opioids. A systematic review of literature. *J Anesthesiol Clin Pharmacol* 2019;35:447.
 11. Carvalho B, Chu L, Cohen S, Riley E. Valdecoxib for Postoperative Pain Management After Caesarean Delivery: A Randomized, Double-Blind, Placebo-Controlled Study. *International Anesthesia Research Society*. Sept 2006. Vol 103, No 3. 664.
 12. Kawasaki Y, Zhang L, Cheng JK, Ji RR. Cytokine mechanisms of central sensitization: distinct and overlapping role of interleukin-1, interleukin-6, and tumor necrosis factor- in regulating synaptic and neuronal activity in the superficial medulla spinalis. *J. Neurosci* 2008;28:5189-94.
 13. Rettig TCD, Verwijmeren L, Dijkstra IM, Boerma D, Garde EMW, Noordzij PG. Postoperative interleukin-6 level and early detection of complication after elective major abdominal surgery. *Ann Surg*. 2016;263: 1207-12.
 14. Ali S, Sofi K, Dar AQ. Comparison of intravenous infusion of tramadol alone with combination of tramadol and paracetamol for postoperative pain after major abdominal surgery in children. 2017;11(2):472-6.
 15. Kiliçaslan A, Tuncer S, Yüceaktaş A, Uyar M, Reisli R. The effects of intravenous paracetamol on postoperative analgesia and tramadol consumption in cesarean operations. *Agri*. 2010;22(1):7-12.
 16. Praveen Rao PN, Knaus EE. Evolution of nonsteroidal anti-inflammatory drugs (NSAIDs): cyclooxygenase (COX) inhibition and beyond. *J pharmaceut sci* 2008; 11 (2): 81-110.

17. Derry S, Derry CJ, Moore RA. Single dose oral ibuprofen plus oxycodone for acute postoperative pain in adults. *Cochrane Database Systemic Reviews*.2013; 6: 1-32.
18. Ehsani M, Moghadamnia A, Zahedpasha S, Maliji G, Haghanifar S, Aghajanpour S, Kani N. The role of prophylactic ibuprofen and N-acetylcysteine on the level of cytokines in periapical exudates and the post-treatment pain. *DARU J Pharm Sc*. 2012: 20(30); 1-6.
19. Vadivelu N, Whitney CJ, Sinatra RS. Pain pathways and acute pain processing. In: Sinatra RS, de Leon-Cassasola, Viscusi ER, eds. *Acute pain management*. Cambridge: Cambridge University Press; 2007. p.3-20.
20. Tanra AH, Rehatta NM, Musba AMT. *Penatalaksanaan nyeri*. Makassar: Departemen Anestesi, Terapi Intensif dan Manajemen Nyeri Fakultas Kedokteran Universitas Hasanuddin; 2013. hal.1-64.
21. Ashburn MA, Ready LB. Postoperative pain. In: Loeser JD, Butler SH, Chapman CR, Turk DC, eds. *Bonica's management of pain*, 3rd Ed. New York: Lippincott Williams & Wilkins; 2001;p.67-72.
22. Ghorri MK, Zhang YF, Sinatra RS. Pathophysiology of acute pain.In: Sinatra RS, de Leon-Cassasola, Viscusi ER, eds. *Acute pain management*. Cambridge: Cambridge University Press; 2007. p.21-32.
23. Vadivelu N, Mitra S, Narayan D. Review recent advances in postoperative pain management. *Yale J Biol Med*. 2010;83(1):11-25.
24. Gottschalk A, Smith DS. New concepts in acute pain therapy: preemptive analgesia. *Am Fam Physician* 2001; 63: 1979-84.
25. Rao M. Acute post operative pain. *Indian J Anesth*. 2006; 50 (5) : 340-4
26. Pyati S, Gan TJ. Perioperative pain management. *CNS Drugs*. 2007;21(3):185-211.
27. Dhillon S. Tramadol-paracetamol fixed-dose combination: a review of its use in the management of moderate to severe pain. *Clin Drug Investig*. 2010;30(10):711-738.
28. Gudin J. Opioid therapies and cytochrome p450 interactions. *J Pain Symptom Manage*. 2012; 44(6):4-14.
29. Gritsenko K. Multimodal therapy in perioperative analgesia *Best Pract Res Clin Anaesthesiol*. 2014;28(1):59-79.

30. Argoff CE. Recent management advances in acute postoperative pain. *Pain Pract.* 2014;14(5):477-87.
31. Gerbershagen HJ, Aduckathil S, Wijck AJM, Peelen LM, Kalkman CJ, Meissner W. Pain intensity on the first day after surgery: a prospective cohort study comparing 179 surgical procedures. *Anesthesiol.* 2013;118(4):934-44.
32. Koh W, Nguyen KP, Jahr JS. Intravenous non-opioid analgesia for peri- and postoperative pain management: a scientific review of intravenous acetaminophen and ibuprofen. *Korean J Anesthesiol.* 2015;68(1):3-12.
33. Joshi GP. Procedure-specific pain management and outcome strategies. *Best Pract Res Clin Anaesthesiol.* 2014;28(2):191-201.
34. Weinbroum AA. Non-opioid IV adjuvants in the perioperative period: pharmacological and clinical aspects of ketamine and gabapentinoids. *Pharmacol Res.* 2012;65(4):411-29.
35. Graham GG, Davies MJ, Day RO, Mohamudally A, Scott KF. The modern pharmacology of paracetamol: therapeutic actions, mechanism of action, metabolism, toxicity and recent pharmacological findings. *Inflammopharmacol.* 2013;21(3):201-32.
36. Bonnefont J, Courade JP, Alloui A, Eschaliere A. Antinociceptive mechanism of action of paracetamol. *Drugs.* 2003;63(2):1-4.
37. Tsen LC. Anesthesia for obstetric care and gynecologic surgery. In: Longnecker DE, Mackey SC, Newman MF, Sandberg WS, Zapol WM, eds. *Anesthesiology*, 3rd Ed. New York: Mc-Graw Hill Medical; 2018. p.1063-82.
38. Ahmad MR, Marwoto. Pengelolaan nyeri pasca seksio caesarea. Dalam: Bisri T, Wahjoeningsih S, Suwondo BS, penyunting. *Anestesi obstetri.* Bandung: Saga Olahcitra; 2013. hal.301-14.
39. Freddy WP. Analgesik-antipiretik, analgesik anti inflamasi non steroid. *Farmakologi dan terapi.* 3th ed. Jakarta: Balai penerbit FKUI. 2008: 183-197.
40. Payan DG, Katzung BG. Obat anti-inflamasi nonsteroid. Analgesic non opioid. *Katzung BG.* 4th ed. Jakarta : EGC,1998 : 558-574.
41. Jongh R, Vissers KC, Meert TF, Booij L, Deyne C, Heylen RJ. Role of interleukin-6 in nociception and pain. *Anesth Analg* 2003;96:1096-103.
42. Tekieh E, Manaheji H, Zaringhalam J, Maghsoudi N, Alani A, Zardoof H. Increased serum interleukin-6 level time-dependently regulates hyperalgesia

- and spinal mu opioid reseptor expression during CFA-induced arthritis. *ECLI J* 2011;10: 23-33.
43. Jun-hua Z. Yu-guang H. Immune system: a new look at pain. *Chin Med J* 2006;119:930-38.
44. Hamza, M & Dionne, RA. 2009.Mechanisms of non-opioid analgesics beyond cyclooxygenase enzyme inhibition. *Curr Mol Pharm*, 2, 1-14
45. Breivik H, Borchgrevink PC, Allen SM, Rosseland LA, Romundstad L, Hals EKB, et al. Assessment of pain. *Br J Anaesth*. 2008;101(1):17-24.
46. Park, A. Ibuprofen use is associated with reduced C-reactive protein and interleukin-6 levels in chronic spinal injury. *J. Spinal Cord Med*. 2020.
47. Sharrawy E, Hakim I, Sameeh E. Attenuation of C-Reactive Protein Increases After Exodentia by Tramadol and Ibuprofen. *Anesth Prog*. 2006 June. 53:78-82.