

DAFTAR PUSTAKA

- Akkamahdevi P, Srinivas HT, Siddesh A, Kaadli N. Comparision of efficacy of sufentanil and fentanyl with low-concentration bupivacaine for combined spinal epidural labour analgesia. Indian Journal of Anaesthesia. Jul Aug 2012;56(4);365-9
- Althaus J, Wax J. Analgesia and Anaesthesia in labour. Obstet Gynecol Clin N Am.2005;32; 231– 44
- Attri J, Makhni R, Sethi S. Levobupivacaine for labor analgesia. Anesth Essays Res. 2016; 10(3); 493-6.
- Baldissari MR: Cardiovascular and Endocrinologic Changes Associated with Pregnancy. In: Textbook of Critical Care Medicine, 5th edition, Fink MP, Abraham E, Vincent Jean-Louis, Kochanek P (eds), Elsevier, Inc, Philadelphia, PA, 2005.
- Ballantyne JC, Cousins MJ, Glamberardino MA, Jamison RN, Mc Grath PA, Rajagopal MR. Modern Labor Analgesia. IASP. August 2011;19(4);1-4
- Birnbach DJ, Browne IM. Anaesthesia for Obstetrics.In Miller RD editor;Miller's Anaesthesia 6th eds.San Fransisco;Elsevier Churchill Livingstone;2006.p. 2306-33
- Cambic CR, Wong CA. Labour analgesia and obstetric outcomesBr. J. Anaesth. 2010; 105 (suppl 1): 50-60
- Camorrcia M, Campogna G, Berritta C, Columb M. The Relative Potencies for Motor Block After Intrathecal Ropivacaine, Levobupivacaine, and Bupivacaine. Anesthesia & Analgesia. April 2007;104(4);904-907
- Chan SY, Chiu JW. Intrathecal labor analgesia using Levobupivacaine 2.5 mg with Fentanyl 25 µg – would half the dose suffice?. Med Sci Monit. 2004; 10(10): 110-4
- Chernecky CC, Berger BJ. Cortisol serum and plasma. In Laboratory and diagnostic procedure 5th eds. Philadelphia;Saunders;2008.p. 123
- Costanzo LS. Endocrine physiology.:Physiology 3rd ed. New York;Saunders;2006.p.408-19
- Ebirim LN, Buowari OY, Ghosh S. Physical and Psychological Aspects of Pain in Obstetrics.In. Pain in perspective. River State; University of Port Hartcourt Teaching Hospital;2012..p. 219-32

Eltschig HK, Lieberman ES, Camann WR. Regional Anesthesia and Analgesia for Labor and Delivery. N Eng J Med 2003 Jan 23;348(4):319-30

Engel NMAA, Van De Velde M, Nijjhuis JG, Marcus MAE. Labour analgesia effects on foetal heart rate. A mini-review. Open Journal of Obstetrics and Gynecology.2011;1; 113-20

Hughes D, Hill D, Fee JPH. Intrathecal ropivacaine or bupivacaine with fentanyl for labour. British Journal of Anaesthesia.2001;87(5);733-7

Jones L, Othman M, Dowswell T, Alfirevic Z, Gates S, Newburn M. Pain management for women in labour: an overview of systematic reviews (Review). The Cochrane Collaboration.2012;(3);1-140

Kim KM, Kim YM. The comparison of clinically relevant doses of intrathecal ropivacaine and levobupivacaine with fentanyl for labor analgesia. Korean J Anesthesiol. Dec 2013; 65(6);525-530

Kono H, Furuhashi N, Shinkawa O, Takahashi T, Tsujiei M, Yajima A. The maternal serum cortisol levels after onset of labor. Tohoku J Exp Med.1987 Jun;152(2):133-7

Labor S, Maguire S. The Pain of Labour. Br J Pain. 2008;2:15-19.

Labor S, Maguire S.The Pain of Labour. British Journal of Pain. December 2008;2(2):15-9

Leone S, Cianni SD, Casati A, Fanelli G. Pharmacology, toxicology, and clinical use of new long acting local anesthetics, ropivacaine and levobupivacaine. Acta Biomed. 2008;79(1):92-105

Long JB, Suresh S. Neonatal anesthesia. In: Barash PG, Cullen BF, Stoelting RK, Cahalan MK, Stock MC, Ortega R, et al, editors. Clinical anesthesia. 8th ed. Philadelphia: Wolters Kluwer. 2008. p.2957.

Loubert C, Hinova A, Fernando R. Update on modern neuraxial analgesia in labour: a review of the literature of the last 5 years. Anaesthesia.2011;66;191-212

Manizech P, Leila P. Perceived Environmental Stressors and Pain Perception During Labor Among Primiparous and Multiparous Women. J Reprod Infertil.2009;10(3):217-23.

Pandya ST. Labour Analgesia;Recent advance.Indian Jornal Anaesthesia.2010;54(5);400-8

Patel NP, Armstrong SL, Fernando R, Columb MO, Bray JK, Sodhi V, Lyons GR. Combined spinal epidural vs epidural labour analgesia: does initial intrathecal analgesia reduce the subsequent minimum local analgesic concentration of epidural bupivacaine?. *Anaesthesia*. June 2012;67(6):584-93

Randall M. Physiology of Stress;Cortisol and the Hypothalamic-Pituitary-Adrenal Axis. February 2011(cited 2013 November 22). Available from:www.dujs.com

Reynolds F. Labour analgesia and the baby: good news is no news. *International Journal of Obstetric Anesthesia* (2011) 20; 38–50

Rudra A. Pain relief in labour-Review article. Update in *Anaesthesia*.2009;6-10

Schrock SD, Smith CH. Labor Analgesia. *American Family Physician*. 2012 March ;85(5):447-54

Vercauteren M, Hans G, De Decker K, Andreassen H. Levobupivacaine combine with sufentanil and ephineprine for intrathecal labor analgesia : a comparison with racemic bupivacaine. *Anesth Analg*. 2001; 93(4) ; 996-1000

Viscomi CM, Rathmell JP, Pace NL. Duration of intrathecal labor analgesia: early versus advanced labor.*Anesthesia and Analgesia*. May 1997;84(5):1108-12

Wilkin I, Pygon B, Peter B, Tyrol D. Analgesia and anaesthesia for obstetrical patients. *Obstetrics Guidelines*. Chicago; University of Illinois Medical Center;2008.p. 1-11

Wong CA. Analgesia and Anesthesia for Labor and Delivery. October 2009 (cited 2013 October 12). Available from:www.The global library of women's medicine.com

Yarnell RW, McDonald JS. *Obstetric Analgesia and Anesthesia*. In DeCherney AH, Nathan L editors; *Current Obstetric & Gynecologic Diagnosis & Treatment*, 9th eds. California;Mc Graw-Hill;2003.p. 291-303