

DAFTAR PUSTAKA

- Choi SJ, Kim MH, Jeong HY, Lee JJ. Effect of intraoperative Lidocaine on anesthetic consumption, and bowel function, pain intensity, analgesic consumption and hospital stay after breast surgery. *Korean J Anesthesiol* 2012; 62(5): 429-34.
- Ferrini R, Paice JA. How to initiate and monitor infusional Lidocaine for severe and/or neuropathic pain. *J Support Oncol* 2004; 2(1): 90–4.
- Frizelle H. Mechanism of postoperative pain-nociceptive in postoperative pain management: an evidence-based guide to practice. Philadelphia: Saunder Elseiver, 2006;27-33.
- Gottschalk A, Smith DS. New concepts in acute pain therapy: preemptive analgesia. *Am Fam Physician* 2001; 63: 1979-84.
- Hans G A, Lauwick S M, Kab A, Bonhome V, Struys M R F, Hans P C, Lamy M L, Loris J L. Intravenous lidocaine infusion reduces bispectral index guided requirements of propofol only during surgical stimulation. *British J of An.*2010. 105(4): 471-9
- Ivan. Pengaruh pemberian lidokain intravena dalam mengurangi kebutuhan opioid pada penanganan nyeri pascabedah laparotomi histerektomi. MKes [tesis]. Makassar: Universitas Hasanuddin; 2015.
- Jensen EW, et al. Monitoring hypnotic effect and nociception with two EEG-derived indices, qCON and qNOX, during general anaesthesia. *Acta Anaesthesiol Scand.* 2014; 58: 933–41.
- Jensen EW. New Findings And Trends For Depth Of Anesthesia Monitoring. *Korean J Anesthesiol* 2018; 71(5): 343-4.
- Kaba A, Laurent SR, Detroz BJ, Sesler DI, Duriex ME, Lamy ML. Intravenous lidocaine infusion fascilitates acutre rehabilitation after laparoscopic colectomy. *Anesthesiology.* 2007;106(1):11-8
- Khan JS, Yousuf M, Victor C, Sharma A, Siddiqui N. An estimation for an appropriate end time for an intraoperative intravenous Lidocaine infusion in bowel surgery: a comparative meta-analysis. *JCA* 2016; 28: 95-104.
- Koppert W, Weigand M, Neumann F, et al. Perioperative intravenous Lidocaine has preventive effects on postoperative pain and morfin consumption after major abdomen surgery. *Anesth Analg* 2004; 98(4): 1050-5.

- Kranke P, Jokinen J, Pace NL, Schnabel A, Hollmann MW, Hahnenkamp K, et al. Continuous intravenous perioperative Lidocaine infusion for postoperative pain and recovery. *Cochrane Database of Systematic Review* 2015; Issue 7. Art. No: CD009642. DOI: 10.1002/14651858.CD009642.pub2.
- Lauretti GR. Mechanisms of analgesia of intravenous Lidocaine. *Rev Bras Anesthesiol* 2008; 58(3): 280-6.
- Lauwick S, Kim DJ, Michelagnoli G, Mistracetti G, Feldman L, Fried G, et al. Intraoperative infusion of Lidocaine reduces postoperative fentanyl requirements in patients undergoing laparoscopic cholecystectomy. *Can J Anesth* 2008; 55 (11): 754-60.
- McCracken, G., Houston, P. and Lefebvre, G. Guideline for the Management of Postoperative Nausea and Vomiting. *Journal of Obstetrics and Gynaecology Canada* 2008; 30, 600-607. [https://doi.org/10.1016/S1701-2163\(16\)32895-X](https://doi.org/10.1016/S1701-2163(16)32895-X)
- Morgan GE, Mikhail MS, Murray, MJ. *Inhalation Anesthetics*, eds. *Clinical anesthesiology*. 5th Ed. New York: McGraw-Hill Companies .2013. pp. 153-73
- Nakhli M S, Kahloul M, Guizani T, Zedini C, Chaouch A, Naija W. Intravenous lidocaine as adjuvant to general anesthesia in renal surgery. *Libyan J of Med*. 2018; 13(1): 1433418
- Navia J T, Lopez J T, Guerrero JJ, Arenas A V, Gutierrez T V. Effect of single dose of lidocaine and ketamine on intraoperative opioid requirements in patients undergoing elective gynecological laparatomies under general anesthesia. A randomized, placebo controlled pilot study. *Farm Hosp*. 2016;40(1):44-51
- Oliveira CMB, Issy AM, Sakata RK. Intraoperative intravenous Lidocaine. *Rev Bras Anesthesiol* 2010; 60(3): 325-33.
- Omar A M, Aboushanab O H, Effect of intravenous lidocaine infusion on sevoflurane requirements as monitored by bispectral index : A randomized double-blinded controlled study. *Egyptian J An*. 2013; 29: 235-39
- Sipahutar TC, Fuadi I, Bisri T. Efek lidokain intravena terhadap nilai Numeric Rating Scale dan kebutuhan fentanyl pascaoperasi dengan anestesi umum. *JAP*. 2013;1(3):167-73
- Stoelting RK. Local Anesthetics. In: *Pharmacology and Physiology in Anesthetic Practice*. 3rd ed. USA: Lippincot-Raven; 2015. p. 199-224

- Tanra AH, Rehatta NM, Musba MT. Lintasan nyeri. Dalam: Penatalaksanaan nyeri. Edisi 1. Makassar: Bagian ilmu anestesi perawatan intensif dan manajemen nyeri fakultas kedokteran universitas hasanuddin,2013;2-10.
- The American Society of Anesthesiologist task force on acute pain management. Practice guidelines for acute pain management in the perioperatif setting. *Br J Anesth* 2004;100:1574-81.
- Ventham NT, Kennedy ED, Brady RR, Paterson HM, Speake D, Foo I, et al. Efficacy of intravenous Lidocaine for postoperative analgesia following laparoscopic surgery: A Meta-Analysis. *World J Surg* 2015; 39(9): 2220-34. DOI: 10.1007/s00268-015-3105-6.
- Weinberg L, Peake B, Tan C, Nikfarjam M. Pharmacokinetics and pharmacodynamics of lignocaine: A review. *World J Anesthesiol* 2015; 4(2): 17-29.
- Wellington J, Chia YY. Patient variables influencing acute pain management. In: Sinatra RS, Leon-casasola OA, Ginsberg B, Viscusi ER, McQuay H. Eds. Acute pain management. Cambridge University press. 2009. pp 33-41
- Yang SY, Kang H, Choi GJ, Shin HY, Baek CW, Jung YH, et al. Efficacy of intraperitoneal and intravenous Lidocaine on pain relief after laparoscopic cholecystectomy. *Journal of International Medical Research* 2013; 0(0): 1-13.
- Yon J H, Choi G J, Kang H, Park J M, Yang H S. Intraoperative systemic lidocaine for pre-emptive analgesics in subtotal gastrectomy : a prospective, randomized, double-blind, placebo-controlled study. *Can J Surg.* 2014: 57(3); 175-82