

## Daftar Pustaka

1. Ono Nelson et al. (2010). *Hemiarthroplasty in the Treatment Fractures of the Femoral Neck*. Rev Bras Ortop. 2010;45(4):382-8
2. Abreu Eduardo et al (2015). Evaluation of the quality of patients undergoing hemiarthroplasty of the hip. Rev Bras Ortop. 2015; 50 (5): 530-536.
3. Baba Tomonori et al (2012). Bipolar hemiarthroplasty for femoral neck fracture using the direct anterior approach. *World J Orthop* 2013; 4(2): 85-89 Available from: URL: <http://www.wjgnet.com/2218-5836/full/v4/i2/85.htm> DOI: <http://dx.doi.org/10.5312/wjo.v4.i2.85>
4. Robertson AG et al (2018). Hip hemi-arthroplasty for neck of femur fracture: what is the current evidence?. *World J Orthop* 2018; 9(11): 235-244 Available from: URL: <http://www.wjgnet.com/2218-5836/full/v9/i11/235.htm> DOI: <http://dx.doi.org/10.5312/wjo.v9.i11.235>
5. Yurdakul et al (2015). Is cemented bipolar hemiarthroplasty a safe treatment for femoral neck fracture in elderly patient?. *Clinical Interventions in Aging* 2015; 10 1063-1067.
6. Moretti VM Post ZD. Surgical approaches for total hip arthroplasty. *Indian Journal Orthop* [serial online] 2017 [cited 2020 Jan 31]; 51: 368-76. Available from: <http://www.ijoonline.com/text.asp?2017/51/4/368/209946>
7. Nilsson Lars et al (1994). Functional after primary hemiarthroplasty and secondary total hip arthroplasty in Femoral Neck Fracture. *The journal of Arthroplasty* Vol. 9 no.4 1994.
8. Noda Mitsuki et al (2017). Comparison of post-operative muscular strength between gamma nailing and hemiarthroplasty system in femoral intertrochanteric fractures. *The open orthopedics journal*, 2017, 11, 255-262.
9. Nagai Takako, Okawa Akihiro (2016). Risk factor affecting postoperative walking ability following hip fracture surgery in the elderly. *Orthop muscular syst* 2016, 5:2
10. Salpakoski Anu et al (2013). Walking recovery after a hip fracture: A

prospective follow-up study among community-dwelling over 60 years old men and women. *BioMed Research International* Volume 2014, Article ID 289549, 11 pages <http://dx.doi.org/10.1155/2014/289549>

11. Snyder Caryn et al (2012). Serum albumin in relation to change in muscle mass, muscle strength, and muscle power in older men. *J Am Geriatr Soc.* 2012 September; 60(9): 1663–1672. doi:10.1111/j.1532-5415.2012.04115.x.
12. Bower Emily et al (2016). Fear of falling after hip fracture: prevalence, course and relationship with one-year functional recovery. *Am J Geriatr Psychiatry.* 2016 December ; 24(12): 1228–1236. doi:10.1016/j.jagp.2016.08.006.
13. Kamel, H. K., Iqbal, M. A., Mogallapu, R., Maas, D., & Hoffmann, R. G. (2003). Time to Ambulation After Hip Fracture Surgery: Relation to Hospitalization Outcomes. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 58(11), M1042–M1045. <https://doi:10.1093/gerona/58.11.m1042>
14. Kuru, T., & Olçar, H. A. (2020). Effects of early mobilization and weight bearing on postoperative walking ability and pain in geriatric patients operated due to hip fracture: a retrospective analysis. *Turkish journal of medical sciences*, 50(1), 117–125. <https://doi.org/10.3906/sag-1906-57>
15. Horikawa, A., Miyakoshi, N., Shimada, Y., & Kodama, H. (2014). Comparison of activities of daily living after osteoporotic hip fracture surgery in patients admitted from home and from geriatric health service facilities. *Clinical interventions in aging*, 9, 1847–1851. <https://doi.org/10.2147/CIA.S69244>
16. Hagino Tetsuo (2007). A simple scoring system to predict ambulation prognosis after hip fracture in the elderly. *Arch Orthop Trauma Surg*, 2007, 127:603–606, DOI 10.1007/s00402-007-0330-y.
17. Rosenbaum Lola Sicard, Kathye E. Light, and Andrea L. Behrman (2002). Gait, Lower Extremity Strength, and Self-Assessed Mobility After Hip

Arthroplasty. *Journal of Gerontology: MEDICAL SCIENCES*, 2002, Vol. 57A, No. 1, M47–M51

18. Mukka Sebastian *et al* (2017). The influence of cognitive status on outcome and walking ability after hemiarthroplasty for femoral neck fracture: a prospective cohort study. *Eur J Orthop Surg Traumatol*, 27:653–658, DOI 10.1007/s00590-016-1873-9.
19. Rasch Anton *et al* (2010). Muscle strength, gait, and balance in 20 patients with hip osteoarthritis followed for 2 years after THA. *Acta Orthopaedica* 2010; 81 (2): 183–188.
20. Holstege Marije S. *et al* (2011). Preoperative Quadriceps Strength as a Predictor for Short-Term Functional Outcome After Total Hip Replacement. *American Congress of Rehabilitation Medicine*.
21. Shim D. –G. *et al* (2017). Rectus femoris muscle atrophy and pretibial recovery caused by preoperative traction in femoral shaft fractures-comparison traction between traction period. *Orthopaedics & Traumatology: Surgery & Research* 103 (2017) 691–695.
22. Nankaku Manabu *et al* (2016). Preoperative gluteus medius muscle atrophy as a predictor of walking ability after total hip arthroplasty. *Phys Ther Res* 19: 8-12.
23. Kamimura Akiko *et al* (2014). Preoperative Predictors of Ambulation Ability at Different Time Points after Total Hip Arthroplasty in Patients with Osteoarthritis. *Rehabilitation Research and Practice* Volume 2014, Article ID 861268, 7 pages <http://dx.doi.org/10.1155/2014/861268>.