

DAFTAR PUSTAKA

1. Frolich MA. Obstetric Anesthesia. In: Morgan G. Morgan & Mikhail's Clinical Anesthesiology. 6th Ed. Butterworth J, Mackey D, Wasnick J, eds. Philadelphia: McGraw-Hill Education; 2018.p.1358-410.
2. Turnbull JH, Aleshi P. Spinal and Epidural Anesthesia. In: Basic Clinical Anesthesia. Sikka PK, Beaman ST, Street JA, eds. New York: Springer; 2015.p.211-232.
3. Oh AY, Hwang JW, Song IA, Kim MH, Ryu JH, Park HP, et al. Influence of the timing of administration of crystalloid on maternal hypotension during spinal anesthesia for cesarean delivery: preload versus coload. *BMC Anesthesiol.* 2014;14:1-5.
4. Skillman CA, Plessinger MA, Woods JR, Clark KE. Effect of graded reductions in uteroplacental blood flow on the fetal lamb. *Am J Physiol.* 1985;249(6):H1098-105.
5. Mercier FJ, Augè M, Hoffmann C, Fischer C, Le Gouez A. Maternal hypotension during spinal anesthesia for caesarean delivery. *Minerva Anesthesiol.* 2013;79(1):62-73.
6. Klohr S, Roth R, Hofmann T, Rossaint R, Heesen M. Definitions of hypotension after spinal anaesthesia for caesarean section: literature search and application to parturients. *Acta Anaesthesiologica Scandinavica* 2010; 54: 909–21.
7. Eckstein KL, Marx GF. Aortocaval compression and uterine displacement. *Anesthesiology.*1974; 40:92–6.
8. Lee A, Ngan Kee WD, Gin T. A quantitative, systematic review of randomized controlled trials of ephedrine versus phenylephrine for the management of hypotension during spinal anesthesia for cesarean delivery. *Anesth Analg* 2002;94:920–6.
9. Lewar EI, Sanjaya DA, Agus IG, Putra S. Kriteria pasien yang menggunakan

- cairan kristaloid dan koloid pada *sectio Caesaria*. *J Kesehatan Prim*. 2018;3(2):124–30.
10. Soma-pillay P, Nelson-piercy C, Tolppanen H, Mebazaa A. Physiological changes in pregnancy. *Cardiovasc J Afr*. 2016;27(2):89–94.
 11. Lopatin J, Jivan K. Spinal Anesthesia. In: *Anesthesiology core review*. Freeman BS, Berger JS, eds. Philadelphia: McGraw-Hill Education; 2016.p.201-204.
 12. Furfaro LL, Chang BJ, Payne MS. Applications for bacteriophage therapy during pregnancy and the perinatal period. *Front Microbiol*. 2017;8:1–14.
 13. Minj SD, Beck R, Kumar A, Tiwari P, Chandan RK, Sen S. The incidence and management of hypotension in the pregnant parturients undergoing Caesarean section following spinal anaesthesia with 0.5% bupivacaine. *Int J Reprod Contraception. Obstet Gynecol*. 2018;7(3):1205-1211.
 14. S M, S A. Effects of general anesthesia vs regional anesthesia on neonatal outcomes: a systemic review and meta analysis. *Int J Anesthesiol Res*. 2016;4:261–71.
 15. American Society of Anesthesiologists. Practice guidelines for obstetric anesthesia: an updated report by the american society of anesthesiologists task force on obstetric anesthesia and the society for obstetric anesthesia and perinatology. *Anesthesiology*. 2016;124:270-300.
 16. Obi VO, Umeora OUI. Anesthesia for emergency cesarean section: a comparison of spinal versus general anesthesia on maternal and neonatal outcomes. *African J Med Heal Sci*. 2018;17:31–4.
 17. Zhang N, He L, Ni J-X. Level of sensory block after spinal anesthesia as a predictor of hypotension in parturient. *Medicine*. 2017;96(25):1-6.
 18. Ayub F, Ahmad A, Aslam KZ, Saleem I. Frequency of headache with 25G or 27G quincke needles after spinal anesthesia in patients undergoing elective

- cesarean section. *Anaesthesia, Pain Intensive Care*. 2017;21(2):170–3.
19. Kementerian Kesehatan Republik Indonesia. Pedoman nasional pelayanan kedokteran anesthesiologi dan terapi intensif. Keputusan Menteri Kesehatan Republik Indonesia Nomor HK.02.02/Menkes/251/2015. 2015:1-139.
 20. Jang W, Cho Y-H, Lee D-H, Kim S-H. Subarachnoid hematoma after spinal anesthesia - a case report. *Anesth Pain Med*. 2018;13(2):154–7.
 21. Tanambel P, Kumaat L, Laleno D. Profil penurunan tekanan darah (hipotensi) pada pasien *sectio Caesarea* yang diberikan anestesi spinal dengan menggunakan bupivakain. *e-CliniC*. 2017;5(1):1–6.
 22. Mohta M. Ropivacaine: Is it a good choice for spinal anesthesia? *J Anaesthesiol Clin Pharmacol*. 2015;31(4):457–8.
 23. Chung C, Choi S, Yeo K, Park H, Lee S, Chin Y. Hyperbaric spinal ropivacaine for cesarean delivery: a comparison to hyperbaric bupivacaine. *Anesth Analg*. 2001;93:157–161.
 24. Riley E, Cohen S, Rubenstein A, Flanagan B. Prevention of hypertension after spinal anesthesia for cesarean section: 6% hetastarch versus lactated ringer's solution. *Anesth Analg*. 1995;81:838–42.
 25. Siddik-Sayyid S, Aouad M, Jalbout M, Zalaket M, Berzina C, Baraka A. Intrathecal versus intravenous fentanyl for supplementation of subarachnoid block during cesarean delivery. *Anesth Analg*. 2002;95:209-13.
 26. Bryson G, Macneil R, Jeyaraj L, Rosaeg O. Small dose spinal bupivacaine for Caesarean delivery does not reduce hypotension but accelerates motor recovery. *Can J Anesth*. 2007;54:531–7.
 27. Bogra J, Arora N, Srivastava P. Synergic effect of intrathecal fentanyl and bupivacaine in spinal anesthesia for cesarean section. *BMC Anesthesiol*. 2005;5:1-5.
 28. Suwardi C. Perbandingan analgesia spinal pada bedah sesar antara kombinasi

- 10 mg bupivakain hiperbarik 0,5% + 12,5 mcg dengan 12,5 mg bupivakain hiperbarik 0,5%. Universitas Indonesia; 2005:1-8.
29. Akmal E. Penyebaran anestetik lokal dan efek hemodinamik pada operasi seksio sesaria dalam anestesia spinal. Universitas Indonesia; 2008:1-5.
 30. Chooi C, Jj C, Rs L, Middleton P, Chemali M, Rs E, et al. Techniques for preventing hypotension during spinal anaesthesia for Caesarean section (Review). *Cochrane Database of Systematic Reviews*. 2017;8:1-335.
 31. Xia F, Feng MDS, Shen PDX. Crystalloid versus colloid in stabilizing hemodynamic of patients undergoing cesarean section with neuraxial anesthesia a randomized controlled trial. *Sci Insigt*. 2019:1-26.
 32. Riley ET, Mangum K, Carvalho B, Butwick AJ. The crystalloid co-load : clinically as effective as colloid preload for preventing hypotension from spinal anaesthesia for Caesarean delivery. *Turkish J Anaesthesiol Reanim*. 2019;(7):35–40.
 33. Mohta M, Dubey M, Malhotra RK, Tyagi A. Comparison of the potency of phenylephrine and norepinephrine bolus doses used to treat post-spinal hypotension during elective Caesarean section. *Int J Obstet Anesth*. 2019;38:25–31.
 34. Cooper D, Carpenter M, Mowbray P, Desira W, Ryall D, Kokri M. Fetal and maternal effects of phenylephrine and ephedrine during spinal anesthesia for cesarean delivery. *Anesthesiology*. 2002;97(6):1582–90.
 35. Ralston D, Shnider S, DeLorimier A. Effects of equipotent ephedrine, metaraminol, mephentermine, and methoxamine on uterine blood flow in the pregnant women. *Anesthesiology*. 1974;4:354–70.
 36. Aziza L, Aqmarina A, Ihsan M. Pedoman pencegahan dan pengendalian coronavirus disease (COVID-19). Kementerian Kesehatan Republik Indonesia. 2020;5:1–214.

37. Vercauteren MP, Hoffmann V, Coppejans HC, Van Steenberge AL, Adriaensen HA. Hydroxyethylstarch compared with modified gelatin as volume preload before spinal anaesthesia for Caesarean section. *Br J Anaesth*. 1996;76(5):731–3.
38. Kinsella SM, Carvalho B, Dyer RA, Fernando R, McDonnell N, Mercier FJ, et al. International consensus statement on the management of hypotension with vasopressors during caesarean section under spinal anaesthesia. *Anaesthesia*. 2018;73(1):71–92.
39. Oh AY, Hwang JW, Song IA, Kim MH, Ryu JH, Park HP, et al. Influence of the timing of administration of crystalloid on maternal hypotension during spinal anesthesia for cesarean delivery: Preload versus coload. *BMC Anesthesiol*. 2014;14.
40. Lirk P, Haller I, Wong CA. Management of spinal anaesthesia-induced hypotension for caesarean delivery: A European survey. *Eur J Anaesthesiol*. 2012;29(9):452–3.
41. Linden EL Van Der, Browne JL, Vissers KM, Antwi E, Agyepong IA, Grobbee DE, et al. Maternal body mass index and adverse pregnancy outcomes: A Ghanaian cohort study. *Obesity*. 2016;24(1):215–22.
42. Chumpathong S, Chinachoti T, Visalyaputra S, Himmunngan T. Incidence and risk factors of hypotension during spinal anesthesia for cesarean section at Siriraj Hospital. *J Med Assoc Thail*. 2006;89(8):1127–32.
43. Basu JK, Jeketera CM, Basu D. Obesity and its outcomes among pregnant South African women. *Int J Gynecol Obstet* [Internet]. 2010;110(2):101–4. Available from: <http://dx.doi.org/10.1016/j.ijgo.2010.02.020>
44. Whiteside JB, Wildsmith JAW. Spinal anaesthesia: An update. *Contin Educ Anaesthesia, Crit Care Pain* [Internet]. 2005;5(2):37–40. Available from: <http://dx.doi.org/10.1093/bjaceaccp/mki017>
45. Jabalameli M, Hashemi J, Soleimani B, Soltani H, Behdad S. Prevention of post-spinal hypotension using crystalloid, colloid and ephedrine with three different combinations: A double blind randomized study. *Adv Biomed Res*. 2012;1(1):36.

46. Iqbal MS, Ishaq M, Masood A, Khan MZ. Optimal dose of prophylactic intravenous ephedrine for spinal-induced hypotension during cesarean section. *Anaesthesia, Pain Intensive Care*. 2010;14(2):71–5.
47. Nishikawa K, Yokoyama N, Saito S, Goto F. Comparison of effects of rapid colloid loading before and after spinal anesthesia on maternal hemodynamics and neonatal outcomes in cesarean section. *J Clin Monit Comput*. 2007;21(2):125–9.
48. Ahmed HO, Hossam M, Adel A. Volume Preload versus Ephedrine Infusion for Prevention of Hypotension Due to Spinal Anesthesia for Cesarean Section. *Open J Anesthesiol*. 2016;31(1):37–44.
49. Furfaro LL, Chang BJ, Payne MS. Applications for bacteriophage therapy during pregnancy and the perinatal period. *Front Microbiol*. 2018;8(JAN):1–14.
50. Tamilselvan P, Fernando R, Bray J, Sodhi M, Columb M. The effects of crystalloid and colloid preload on cardiac output in the parturient undergoing planned cesarean delivery under spinal anesthesia: A randomized trial. *Anesth Analg*. 2009;109(6):1916–21.
51. Braveman F, Scavone B, Bleasing M, Wong C. Obstetric anaesthesia. *Clinical Anesthesia*. 8th ed. Barash PG, Cullen BF, Stoelting RK, Cahalan MK SM, editor. Philadelphia; 2017. 2841–9. p.