

Daftar Pustaka

1. Dooley J, Roy L, Flanagan E, Martin G,. Anesthesia for orthopedic surgery. *In: Mackey Sc, Newman MF, Sanberg WS, Zapol WM, eds. Anesthesiology. edisi 3.* New York : McGraw-Hill; 2018. p 1108-18.
2. *Tarpada SP, Morris MT, Burton DA.* Spinal fusion surgery: a historical perspective. *J Orthopaed.* 2016;14(2):134-6.
3. Carragee EJ, Cheng I, Salari N, Brown AP, Larson CP. Spine surgery. *In: Jaffe RA, schmiesing CA, Golianu B, eds. Anesthesiologist's manual of surgical procedures, edisi 5.* Philadelphia: Wolters Kluwer; 2014. p.1520-46.
4. *Nagappa S, Kalappa S, Sridhara RB,.* Clonidine, as an adjuvant to caudal epidural ropivacaine for lumbosacral spine surgeries. *Anesth Essays Res.* 2018;12(1): 240-5.
5. *Saoud A, Elkabarity R, abdellatif A, .* Efficacy of preemptive caudal analgesia in single level lumbar spine decompression and fusion surgery. *World Spin Column J.* 2012;3(3):7-19.
6. *Vadivelu N, Mitra S, schermer E, Kodumudi V, Kaye AD, Urman RD.* Preventive analgesia for postoperative pain control: a broader concept. *Loc Reg Anesth.* 2014;7:17-22.
7. *Lopez MB, Cadorniga AG, Gonzalez JML, Suarez ED, Carballo CL, Sobrino FP.* Erector Spinae Block. A narrative review. *Central Eur. J. Clinical Res.* 2018;1(1);28-39.
8. *Forero M, Adhikary SD, Lopez H, Tsui C, Chin KJ, .* The Erector Spinae Plane Block A Novel Analgesic Technique in Thoracic Neuropathic Pain. *Reg Anesth Pain Med.* 2016;41:621-7.
9. *Debono B, Wainwright TW, Wang MY, Sigmundsson FG, Yang MMH, Nanninga HS, dkk.* Consensus statement for perioperative care in lumbar spinal fusion: enhanced recovery after surgery society recommendation. *Spine J.* 2021;21(5):729-52.
10. *Waelkens P, Alsabbagh E, Sauter A, joshi GP, Beloeil H.* Pain management after complex spine surgery. *Eur J Aneaesthesiol.* 2021;38:hlm985-994 .
11. *Prolo LM, oklund SA, Zawadzki N, Desai M, Polo DJ.* Uninstrumented posterior lumbal interbody fusion: have technological advances in stabilizing the lumbar spine truly improved outcomes? *Wold neurosurg.* 2018;115;503-4.

12. *Rasa SN, Carr BD, Cohen M, Finnerup NB, Flor H, Gibson S, dkk.* The revised international association for the study of pain definition of pain: concepts, challenges, and compromises. *Pain.* 2020;161(9):1976-82.
13. *IASP's proposed new definition of pain release for comment.* International Association for the study of pain. 2019. [Diunduh 8 Juli 2021]. tersedia dari <https://www.iasp-pain.org/PublicationsNews/NewsDetail.aspx?ItemNumber=10475>.
14. *Joshi GP, Ogunnaike BO.* Consequences of inadequate postoperative pain relief and chronic persistent postoperative pain. *Anesthesiol Clin North Am.* 2005;23(1):21-36.
15. *Macres SM, Moore PG, Fishman SM.* Acute pain management. In: Barash PG, Cullen BF, Stoelting RK, Cahalan MK, Stock MC, Orthega R, Sharar SR, Holt NF. *Clinical anesthesia*, edisi8. 2017. Wolters Kluwer : Philadelphia. p. 3919-32.
16. *Lavand'homme P.* From preemptive to preventive analgesia: time to reconsider the role of perioperative peripheral nerve block? *Reg Anesth Pain med.* 2011;36(1)4-6.
17. *Wahdah SW, Radwan TA, Mohammed MM, Mohamed AA, Salama AK.* Effect of bilateral ultrasound-guided erector spinae blocks on postoperative pain and opioid use after lumbar spine surgery: A prospective randomized controlled trial. *Egypt J Anaesth.* 2020;37(1):100-6.
18. *Altiparmak B, Tker MK, Uysal AI, Demirbilek SG.* Comparison of the efficacy of erector spinae plane block performed with different concentrations of bupivacaine on postoperative analgesia after mastectomy surgery: randomized, prospective, double blinded trial. *BMC Anesthesiol.* 2019;19(1):31.
19. *Chin KJ, Adkhikary SD, Forero M.* Erector spinae plane (ESP) block: a new paradigm in regional anesthesia and analgesia. *Curr Anesthesiol Rep.* 2019;9;271-80.
20. *Apibunyopas Y, Ho B.* Ultrasound-guided erector spinae plane (ESP) block: A novel intervention for mechanical back pain in the emergency department. *CJEM.* 2019;21(2):302-5.
21. *Elsharkawy H, Naguib MA.* Peripherally acting analgesic. In: Flood P, Rathmell JP, Shafer S. *Stoelting's pharmacology & physiology in anesthetic practice*, edisi5. Philadelphia :Wolters Kluwer p282-287.

22. *Chen LL, Mao J.* Nonopioid pain medication. In: Gropper MA, Cohen NH, Eriksson LI, Fleisher LA, Leslie K, Wiener-Kronish JP, eds. *Miller's anesthesia*, edisi 9. Philadelphia : Elsevier. p877-8.
23. *Butterworth JF, Mackey DC, Wasnick JD.* Local anesthetic. in: Butterworth JF, Mackey DC, Wasnick JD, eds. *Morgan & Mikhail's Clinical anesthesiology*, edisi 6. McGraw Hill : New York. p 452-471.
24. *Kwon WJ, bang SU, Sun WY.* Erector spinae plane block for effective analgesia after total mastectomy with sentinel or axillary lymph node dissection: a report of three cases. 2018. *J Korean Med Sci*:33(45);e291.
25. *Analgesic agent.* In: Butterworth JF, Mackey DC, Wasnick JD, eds. *Morgan & Mikhail's clinical anesthesiology*, edisi 6. New York : McGraw Hill; 2017. p189-98.
26. *Cumming KI, Naguib MA.* Opioid agonist and antagonist. In: Flood P, Rathmell JP, Shafer S, eds. *Pharmacology and physiology in anesthetic practice*. edisi 5. Wolters Kluwer : Philadelphia. p217-50.
27. *Daniel M, Weiskopf RB, Noorani M.* Fentanyl augments the blockade of the sympathetic response to incision (MAC-BAR) produced by desflurane and isoflurane : desflurane and isoflurane MAC-BAR without and with fentanyl. *Anesthesiology*. 1998;88:43–49.
28. *Becker LD, Paulson BA, Miller RD, Severinghaus JW, Eger EI.* Biphasic respiratory depression after fentanyl - droperidol or fentanyl alone used to supplement nitrous oxide anesthesia . *Anesthesiology*. 1976; 44:291–295.
29. *Flood P, Shafer S.* Inhaled anesthetics. In: Flood P, Rathmell JP, Shafer S. eds. *Stoelting's pharmacology & physiology in anesthetic practice*, edisi 5. 2015. Wolter Kluwer : Philadelphia. p98-152.
30. *Perouansky M, Pearce RA, Hemmings H, Franks NP.* Inhaled anesthetics: mechanism of action. in: Gropper MA, Cohen NH, Eriksson LI, Fleisher LA, Leslie K, Kronish, JPW, eds. *Miller's anesthesia*. edisi 9. Philadelphia : Elsevier. p487-506.
31. *Ji F, Wang D, Zhang J, Liu H, Peng K.* Effects of propofol anesthesia versus sevoflurane anesthesia on postoperative pain after radical gastrectomy: a randomized controlled trial. *J Pain Res*. 2018;11:1247-1254.
32. *Mishra LD, Rajkumar N, Singh SN, Dubey RK, Yadav G.* A comparative study of propofol and isoflurane anaesthesia using butorphanol in neurosurgery. *Indian J Anaesth*.2009;53(3):324-29.
33. *Farhan NE, Rees DA, Evans C.* Measuring cortisol in serum, urine and saliva - are our assay good enough? *Ann Clin Biochem*.2017;54(3);308-22.

34. *Hall JE, Guyton AC, . Adrenocortical hormones. . .In: Hall, JE, ed. Guyton and hall textbook of medical physiology, edisi 13. bab 78. hal 964-979.*
35. *Ali NS, Elshorbagy HM, Hassanien AAM. Efficacy of ultrasound guided erector spinae plane block on hemodynamic in patient undergoing abdominal surgery. Med J Malaysia. 2020;3(31):107-11.*
36. *Felix SP, Nielsen L, Bak P, Roth D, Fischer M, Bjerring, Zbinden AM. Analgesic effect in human of subanaesthetic isoflurane concentration evaluated by experimentally induced pain. Br J Anaesth.1995;75(1):55-60.*
37. *Mohaghegh T, Yazdi B, Norozi A Fateh S, Modir H, Mohammedbeigi A. Effect of intravenous anesthesia with propofol versus isoflurane inhalation anesthesia in postoperative pain of inguinal herniotomy: a randomized clinical trial. Med Gas.2017;7:86=92.*
38. *Yang M, Yu Y, Liu Q. Analgesic effect of sevoflurane and isoflurane on elderly patients with colon cancer and their influences on immunity and postoperative cognitive function. Iran J Public Health.2019;48(3):444-50.*
39. *Ji FH, Wang D, Zhang, Liu HY, Peng K. effect of propofol anesthesia versus sevoflurane anesthesia on postoperative pain after radical gastrectomy: a randomized controlled trial. J Pain Res. 2018;(22):1247-54.*
40. *Inagaki Y, Mashimo T, Kuzukawa A, Tsuda Y, Yosiya I. Epidural lidocaine delays arousal from isoflurane anesthesia. Anesth Analg.1994;79:68-72.*
41. *Hlgashizawa T, Koga Y. Effect of infraorbital nerve block under general anesthesia on consumption of isoflurane and postoperative pain in endoscopic endonasal maxillary sinus surgery. J Anaesth.2001;15:136-8.*
42. *Casati L, Galinski SF, Barrera E, Pol O, Puig MM. isoflurane requirements during combined general/epidural anesthesia for major abdominal surgery. Anesth Analg.2002;94:1331-7.*
43. *Navaro EM. desfluran-allgemeinanasthesie zur sectio caesarea: vergleich mit isofluran und epiduralanasthesie. AINS.2000;35(4):232-6.*
44. *Liem TH. Hasenbos MAWM. Booij LHDJ. Gielen MJM. Coronary artery bypass grafting using two different anesthetic technique. J.Cardiothorac.Vasc.Anesth. 1992;6(2):156-61.*
45. *Sukmono BB, Rahendra, Sari A Aditianingsih D, Tantri AR. Comparison of intraoperative hemodynamic stability between quadratus lumborum and epidural block in patients who underwent laparoscopic nephrectomy. J. Phys. Conf. Ser. 2019. 1246 012064.*

46. Zhang TJ, Zhang JJ, Zong YQ, Hong YZ, Qiu Y, Hua Z. Bilateral erector spinae plane block for open posterior lumbar surgery. *J Pain Res.* 2020;13:709-17.
47. Javrosky BR, Raff H, Carroll TB, Schimnich AA, Singh RJ, Franco JM. New cutoff for the biochemical diagnosis of adrenal insufficiency after ACTH stimulation using specific cortisol assays. *J. Endocr. Soc.* 2021;5(4):1-11.
48. Sharvika A, Pushkar W, Sonal S, Aditi S, Prathamesh B, Deeisha G. Evaluation of presurgical serum cortisol level in patients undergoing major maxillofacial surgery. *Ann Maxillofac Surg.* 2020; 10:25-30.
49. Naito Y, Tamai S, Shingu K, Shindo K, Matsui T, Segawa H, dkk. Response of plasma adrenocorticotropic hormone, cortisol, and cytokine during and after upper abdominal surgery. *Anesthesiology.* 1992;77:46-31.
50. Forero M, Adhikary SD, Lopez H, Tsui C, Chin KJ. The erector spinae plane block. A novel analgesic technique in thoracic neuropathic pain. 2016. *Reg Anesth Pain Med*;41(5):621-7.
51. Erlangga ME, Sitanggang RH, Bisri T. Perbandingan pemberian deksametason 10 mg dengan 15 mg intravena sebagai adjuvan analgetik terhadap skala nyeri pascabedah pada pasien yang dilakukan radikal mastektomi termodifikasi. *JAP.* 2015;3(3):146-54.
52. Cardwell TW, Zabala V, Mineo J, Ochnerr C. The effects of perioperative peripheral nerve blocks on peri- and postoperative opioid use and pain management. *Am Surg.* 2021;0(0):1-9.
53. Liu Q, Chely JE, William JP, Gold MS. Impact of peripheral nerve block with low dose local anesthetics on analgesia and Functional Outcomes Following Total Knee Arthroplasty: A Retrospective Study. *Pain Med.* 2015. 16(5):998-1006.
54. Liang X, Zhou W, Fan Y. Erector spinae plane block for spinal surgery: a systematic review and meta-analysis. *Korean J Pain.* 2021;34(4):487-500.
55. Yeşiltaş S, Abdallah A, Uysal Ö, Yılmaz S, Çinar İ, Karaaslan K. The efficacy of intraoperative freehand erector spinae plane block in lumbar spondylolisthesis: A randomized controlled study. *Spine.* 2021;46(17):E902-10.
56. Modig J. Regional anaesthesia and blood loss. *Acta anesthesiol Scand.* 1988;32(supp 89):44-8.
57. Dultz LA, Rosalind, Dumas RP, Grant JL, Park C, Alexander JC, dkk. Safety of erector spinae plane block in patients with chest wall trauma on venous thromboembolism prophylaxis. *J Surg Res.* 2021;(263):124-9.

58. *Wang JQ, Chen LY, Jiang BJ, Zhao YM.* Development of a nomogram for predicting blood transfusion risk after hemiarthroplasty for femoral neck fracture in elderly patients. *Med Sci Monit.* 2020;26:e920255.
59. *Misra S, Awal S.* Does erector spinae plane block result in improved postoperative analgesia and enhanced recovery in adult patients after cardiac surgery? *Interact CardioVasc Thorac Surg.* 2021;32:873–7.
60. *Khrisna SN, Kaushal B, Bisoi AK.* Bilateral Erector Spinae Plane block for Acute Post-Surgical pain in Adult Cardiac Surgical patients: a randomized controlled trial. *Cardiothorac Vasc Anesth.* 2018;33(2):368-75.
61. *Macaire P, Ho N, Tan N, Nguyen B, Vu V, Quach C, dkk.* Ultrasound guided continuous thoracic erector spinae plane block within an enhanced recovery program is associated with decreased opioid consumption and improved patient postoperative rehabilitation after open cardiac surgery. *J Cardiothorac Vasc Anesth.* 2019;33(6):1659-67.
62. *Yuan Z, liu J, Jiau K, Fan Y, Zhang Y.* Ultrasound guided erector spinae plane block improve opioid sparing perioperative analgesia in pediatric patients undergoing thoracoscopic lung lesion resection: a prospective randomized controlled trial. *Transl Pediatr.* 2022;11(5):706-14.
63. *Ahiskalioglu A, Tulgar S, Celik M, Ozer Z, Alici HA, Aydin ME.* Lumbar erector spinae plane block as a main anesthetic method for hip surgery in high risk elderly patients : Initial experience with a magnetic resonance imaging. *Eurasian H Med.* 2020;52(1):16-20.
64. *Kim WJ, Lim W.* Efficacy of erector spinae plane block with opioid-sparing analgesic technique in breast-conserving surgery. *Ann Surg Treat Res.* 2022;100(5):253-9.
65. *Zhu L, Wang M, Wang X, Wang Y, Chen L, Li J.* Changes of opioid consumption after lumbar fusion using ultrasound guided lumbar erector spinae plane. *Pain Physician.* 2021 Mar;24(2):E161-E168.
66. *Chitnis SS, Tang R, Mariano ER.* The role of regional analgesia in personalized postoperative pain management. *Korean J Anesthesiol.* 2020;73(5):363-71.
67. *Takemura H, Fujita D, Matsuda M, Fujita K, Sakaguchi M, Amaya F.* Peripheral nerve block combined with general anesthesia for lower extremity amputation in hemodialysis patients: case series. *JA Clin Rep.* 2018;4(7):hlm 1-5.

68. *Pearse RM, Harrison DA, James P, Watson D, Hinds C, Rhodes A.* Identification and characterisation of the high risk surgical population in the united kingdom. *Crit Care.*2006;10(3):R81.
69. *Pestel G, hoeft A.* Hemodynamic monitoring and management in patient undergoing, high risk surgery: a survey among north american and european anesthesiologists. *Critical care.*2011;15:R197.
70. *Daabis M.* American society of anaesthesiologist physical status classification. *Indian J Anaesth.* 2011;55(2):111-5.
71. *Bergmann I, Heetfeld M, Crozier TA, Schafdecker HG, Poschi R, Wiese CH.* Peripheral nerve blocks give greater hemodynamic stability than general anesthesia for ASA III patients undergoing outpatient knee arthroscopy. *Cent. Eur. J. Med.* 2013;8(4): 436-42.
72. *Christos SC, Chiampas G, Offman R, Rifenburg R.* Ultrasound guided three in one nerve block for femur fracture. *West. J. Emerg. Med.* 2010;4(9): hlm 310-3.
73. *Lai HY, Foo LL, Yong CF, Loh PS, Chaw SH, Hasan MS, et al.* The hemodynamic and pain impact of peripheral nerve block versus spinal anesthesia in diabetic patients undergoing diabetic foot surgery. *Clin Auton Res.* 2020;30(1):53-60.
74. *Tonkovic D, Stambolija V, Lozic M, pavlovic DB, Sekulic A, Baronica R, Peric M.* Scalp block for hemodynamic stability during neurosurgery. *Period. Biol.* 2015;2(117):247-50.
75. *Van den Broek RJC, Van de Geer R, Schepel NC, Llu WY, Bouwman RA, Versyck B.* Evaluation of adding the Erector spinae plane block to standard anesthetic care in patients undergoing posterior lumbar interbody fusion surgery. *Sci Rep.* 2021;11.
76. *Singh S, Choudhary NK, Lalin D, Verma V.* Bilateral ultrasound guided erector spinae plane block for postoperative analgesia in lumbar spine surgery; a randomized control trial. *Neurosurg Anesthesiol.*2019;00(00):1-5.
77. *Sinha C, Kumar A, Kumar A, Kumari P, Agrawal P, Arun SK.* Two level bilatral ultrasound guided erector spinae plane block for kyphosis correective surgery: a case report. *Anesth Pain.* 2022;17:327-330.
78. *Ueshima H, inagaki M, Toyone T, Otake H, .* Efficacy of the erecctor spinae plane block for lumbar spinal surgery: a retrospective study. *Asian Spine J.* 2019;13(2):254-7.

79. *Yayik AM, Cesur S, Ozturk F, Ahiskalioglu A, Ay AN, Celik EC, et al.* Postoperative analgesic efficacy of the ultrasound-guided erector spinae plane block in patients undergoing lumbar spinal decompression surgery: a randomized controlled study. *World Neurosurg.* 2019;126:e779-e785.
80. *Steverson M.* Ageing and health . Oktober 2021. diunduh 13 September 2022. dari: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>.
81. *Butterworth JF, Mackey DC, Wasnick JD.* Pediatric anesthesia. In: Butterworth JF, Mackey DC, Wasnick JD, eds. *Morgan & Mikhail's clinical anesthesiology*, edisi 6. New York : McGraw Hill; 2017. p1517-66.
82. *Kessler P, Aybek T, Neidhart G, Dogan, S, Lischke V, Bremerich DH, Byhahn C.* Comparison of three anesthetic techniques for off pump coronary artery bypass grafting: general anesthesia, combined general and high thoracic epidural anesthesia, or high thoracic epidural anesthesia alone. Comparison of three anesthetic techniques for off pump coronary artery bypass grafting: general anesthesia, combined general and high thoracic epidural anesthesia, or high thoracic epidural anesthesia alone.