

COVER LETTER OF EDITED MANUSCRIPT

Date: 08th February 2022

To
The Editor,
VISION JOURNAL

I am enclosing herewith a manuscript entitled “**CHOROIDAL THICKNESS IN CORRELATION WITH AXIAL LENGTH AND MYOPIA DEGREE**”. We would like to inform you that we have revised our manuscript based on academic editor and reviewers’ suggestion. Here are we attached point by point response:

RESPONSE TO REVIEWER 1 COMMENTS

Point 1 = The inclusion criteria should be specified in a more detailed manner. The type of OCT device should be indicated.

Response 1:

We have added an explanation of the inclusion criteria to the main manuscript line 81 - 87, while an explanation of the types of OCT in line 101-103. In this study, our criteria for recruiting subjects were patients aged 20-50 years, had a refractive error $\geq -0.5D$, did not suffer from anterior segment abnormalities during examination, no history of eye infection or eye surgery. The type of OCT used in our study is Heidelberg Spectralis® OCT (3-mode), Germany (line 99-101).

Point 2 = The classification of myopia into degrees is presented differently in the Introduction (line 42 - 43) and in the Results (line 102-103) section.

Response 2:

In the manuscript line 42-43 we cite the references by Kaiti et al. (2021) without changing its contents at all. However, in line 102-103 we refer to the standard of assigning myopia by Okafor et al. (2009) which states that low myopia ($\leq -3.00 D$), moderate (-3.25 to $-6.00 D$), and high ($> -6.00 D$) myopias. However, we agree with your input that all references should be done in line, therefore we replace the references in the introduction section from Kaiti’s statement to Okafor’s statement while in the method section we continue to use the criteria by Okafor as previously written.

Pont 3 = The result is long time known and therefore predictable: higher degrees of myopia and longer axial lengths are associated with thinner choroids. However, the presentation of choroidal thickness according to the quadrant may be of some interest, especially in relation to the axial length. One important interpretation of the results which is revealed in the Discussion section is related to the role of the choroid in modeling the retina and sclera in patients with myopia.

Response 3:

We have added an explanation of the correlation between CT and retinal change on line 230-240, like explained below:

Based on Jin et al. (2016), the myopic retinas were thinner than those of emmetropic or hyperopic subjects, especially in the superior parafoveal and all 4 perifoveal subfields ($p < 0.05$). But, the results of previous studies on factors influencing the thickness of the ganglion cell layer and nerve fiber layer have been conflicting. While some suggested that the thickness of the ganglion cell layer and peripapillary nerve fiber layer is correlated with spherical equivalent refraction and axial length in adults, others did not observe this relationship (line 234-240).

Karahan (2013) reported that choroidal change plays a major role in the development and progression of many retinal disease. Thickening of the choroid could affect the nutrition supply to the retina, because outer retina layer is nourished by the choroidal vasculature. Thus, choroidal thickness provides useful information to clinicians (line 241-244).

RESPONSE TO REVIEWER 2 COMMENTS

Point 1 = Introduction, line 32: Although both statements, that myopia is more likely to occur at young age and more frequent in people who are for example microscopists, it should be formulated differently. As it currently stands, it reads as if young people often work as microscopists.

Response:

Thank you, we agree with your comment. In the introduction, we have made a sentence change from “myopia is more likely to occur at young age and more frequent in people who are for example microscopists” become “Myopia is more likely to occur in several condition such as young age (mostly 8 to 15 years old), hereditary person with myopic parent, and persons who work extensively with the eyes such as microscopists, computer users, or university students” (line 33-35).

Point 2 = Materials and methods section: The authors should add, as stated at the end of the manuscript, that informed consent was obtained, that the study was approved by an IRB and that the study adhered to the Declaration of Helsinki. The authors should also mention which software was used for statistical analysis.

Response:

We have added a sentence regarding the ethical statement according to the reviewer's input on the manuscript in line 109-112.

Point 3 = Table 3: It would be of interest for the readers if a figure showing the most relevant correlations could be provided.

Response:

According to the reviewer's input, we have added "figure 5" as a form of visualization that depicts the correlation between choroidal thickness with axial length and choroidal thickness with myopia degree (line 174-175).

Point 4 = Maybe it would also be of interest to have a control group with no myopia.

Response:

We have already added the examinations result of 10 healthy subjects (20 eyes) as control group. The results are presented in table 1 (line 135-136). We also attach normal OCT photo for figure 1 and figure 2 (line 125 and 130).

RESPONSE TO REVIEWER 3 COMMENTS

The paper of Habibah Setyawati Muhiddinn and coauthors reports relationships between axial length, choroidal thickness, and myopia. Unfortunately, these findings are well-studied and do not require additional confirmation obtained in a study with limited number of patients and I can't recommend this paper for publication.

Point 1 = "It occurs when there is an excessive and continuous expansion of the axial length (AL)" not only in this case "resulting in a change in the secondary fundus leading to visual impairment, choroidal neovascularization, retinal detachment, zonal areas of chorioretinal atrophy, myopic macular schisis and hole." This occurs in minority of cases.

Response:

Thank you very much for your suggestion. We agree with your opinion. We wrote that statement based on the references we quoted from a combination of several manuscripts. However, in the abstract section we have made improvements by writing a general explanation of myopia in line 9-11: "Myopia is a condition in which the visual images come to a focus in front of the retina of the eye. This disease is a major cause of visual disability which present in 108 million persons globally".

Point 2 = The methods are poorly described. Especially CT measurement which was performed in manual manner (instead of semi-automatically manner stated) and which requires two graders to be done.

Response:

We agree with your opinion. However, it is true that the tool we use is a semi-automatic machine, the type of machine we used is Heidelberg Spectralis® OCT, Germany, where the inspection was carried out by 2 technicians (two graders) then the results were blinded confirmed by the research team (authors: HSM and AMI).

For the method section, we have also provided additional detailed information regarding control subjects, inclusion criteria, examination techniques, ethical statements and data analysis techniques that we used (line 77-111).

Point 3 = I see no new data from this manuscript.

Response:

In this study, the novelty value that we propose is the measurement of the degree of choroidal thickness based on the segmentation of the eyeball area. We made measurements at nine examination points (C=Sub Fovea, T6=Temporal 3, T3=Temporal 1.5, N3= Nasal 1.5, N6 = Nasal 3, S6 = Superior 3, S3 = Superior 1.5, I3 = Inferior 1.5, and I6 = Inferior 3) where it was found that CT correlated with the degree of myopia and axial length. Our study found that the thinnest CT measurements were in the nasal area in subjects with mild, moderate and severe myopia and this explanation has been included in the manuscript (line 135-136).

RESPONSE TO REVIEWER 4

Point 1 = This study provides the following information that it is necessary to screen for choroidal thickness in myopic patients using SD-OCT, especially in moderate and high degrees to prevent complications. There was a significant correlation between the degree of myopia and choroidal thickness. Therefore, a higher degree of myopia will cause a decrease in the thickness of the choroid. A significant increase in myopia degree will lead to a decrease in the choroidal thickness. Also, it is necessary to measure the length of the eyeball axis in myopic patients because a decrease in choroidal thickness is significant to the elongation of the eyeball.

Response:

We agree with your comment. In our manuscript we have written suggestions about it is better if the patient performs an examination of the axial length and choroidal thickness to prevent serious complications of myopia. We recommend this examination to prevent the possibility of further complication such as the progressivity of myopic choroidal neovascularization (CNV) which will eventually lead to worsening of vision. We have written this suggestion in the manuscript line 258-260.

We are looking for possible evaluation and also publication in Vision journal.

Thank you very much your kind attention.

Sincerely,

Andi Muhammad Ichsan

Ophthalmology Department, Medical Faculty, Hasanuddin University

Email: am_ichsan@unhas.ac.id

Mobile phone: +6281342280880

Postal address: Jl.Perintis Kemerdekaan Km.10, Tamalanrea, 90245

FIGURES




Figure 1. Choroidal thickness based on degrees of myopia horizontally.
(A) Normal subject (B) Low myopia, (C) Moderate myopia, (D) High myopia.




Figure 2. Choroidal thickness based on degrees of myopia vertically. (A) Normal subject (B) Low myopia, (C) Moderate myopia, (D) High myopia




Figure 3. Choroidal thickness based on axial length horizontally.
 (A) Normal axis (22,47 mm) (B) Medium axis (23,75 mm), (C) Long axis (27,60 mm)




Figure 4. Figure 3. Choroidal thickness based on axial length vertically. (A) Normal axis (22,47 mm) (B) Medium axis (23,75 mm), (C) Long axis (27,60 mm)




Figure 5. (A) Correlation between choroidal thickness and myopia degree (B) Correlation between choroidal thickness and axial length.

CHOROIDAL THICKNESS IN CORRELATION WITH AXIAL LENGTH AND MYOPIA DEGREE

Habibah Setyawati Muhiddin¹, Andi Ratna Mayasari¹, Batari Todja Umar¹, Junaedi Sirajuddin¹, Ilhamjaya [Patellongi](#)², Itzar Chaidir Islam¹, Andi Muhammad Ichsan^{1*}

¹Ophthalmology Department, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia;

²Physiology Department, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia;

* Correspondence: am_ichsan@med.unhas.ac.id; Tel.: +6281342280880

Abstract: Background: Myopia is a condition in which the visual images come to a focus in front of the retina of the eye. This disease is a major cause of visual disability which present in 108 million persons globally. A major cause of visual disability is myopia. It occurs when there is an excessive and continuous expansion of the axial length (AL), resulting in a change in the secondary fundus leading to visual impairment, choroidal neovascularization, retinal detachment, zonal areas of chorioretinal atrophy, myopic macular schisis and hole. Purpose: This study aims to determine the relationship between the degree of myopia, the axial length, and the choroidal thickness (CT). Methods: This is an observational analytical study that made use of a cross-sectional design. A total of 49-59 participants with refractive errors, underwent treatment at Hasanuddin University Hospital and 69-116 eyes were measured and analyzed. The choroidal thickness was measured using the Enhance Depth Imaging OCT (EDI-OCT) tool which is divided into nine observational areas. Furthermore, all data obtained were compared using statistical analysis such as the one-way ANOVA and Pearson correlation test ($p < 0.05$). Results: There was a significant relationship between the choroidal thickness with axial length ($p < 0.05$), and myopia degrees ($p < 0.05$). Conclusions: The thickness of the choroid decreases with an increase in the axial length and degree of myopia which further indicates that the higher the myopia degree the thinner the choroidal vasculature.

Keywords: Choroidal thickness; Myopia; Axial length

Citation: Muhiddin HS et al. CHOROIDAL THICKNESS IN CORRELATION WITH AXIAL LENGTH AND MYOPIA DEGREE. *Vision* 2022, 6, x. <https://doi.org/10.3390/xxxxx>

Received: date
Accepted: date
Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Myopia, also known as shortsightedness, is a major cause of visual disability around the world. In 1972 and 2004, the prevalence of myopia increased from 25 to 44% in the United States while in Asia, the prevalence is approx. >80%.⁽¹⁾ In 2010, it was noted that the uncorrected refractive error was the major cause of vision impairment and the second most frequent cause of blindness affecting 108 million persons globally. Furthermore, the cases of myopia are expected to increase by more than 5000 million by 2050.⁽²⁾

Myopia is more likely to occur in several condition such as young age (mostly 8 to 15 years old), hereditary person with myopic parent, and persons who work extensively with the eyes such as microscopists, computer users, or university students. Myopia is more likely to occur within the ages of 8 and 15 and it is prevalent in persons who work extensively with the eyes such as microscopists.^(1,3) According to the World Health Organization, the major cause of vision impairment and blindness around the world are macular degeneration, vitamin A deficiency, infectious disease, uncorrected refractive error with cataracts, and myopia.⁽⁴⁾

Furthermore, myopia has been classified according to anatomical and pathological features, the age of onset, the rate of progression, the degree and theory of development. Physiological myopia occurs when the refractive components of the eye fail to correlate,

unlike pathologic myopia (alternatively, malignant or degenerative myopia) which occurs when the optical system of the eye lies outside the limit of normal biological variations. According to degree, myopia can be grouped into low (<3.00 D), medium (3.00 D-5.00 D), and high (>5.00 D).(5)

According to Okafor et al. (2009), the degree of myopia is could be divided into 3 categories: very low (<-1.0 D), low (≥ -1 to ≤ -3.00 D), moderate (> -3.00 to < -6.00 D), high (> -6.00 D to ≤ -10 D) and very high (> -10 D) myopias.(6)

Myopia and refractive-error disorder may develop when there are irregular contri-butions of the ocular components to the eye structures. Four structures that contribute to the refractive status of a given human eye are aqueous and vitreous humor, cornea, and lens. When the lens and cornea fail to neutralize the axial length (AL) shortening or elon-gation, hyperopia or myopia may occur. Therefore, some parameters such as the anterior chamber depth (ACD), corneal curvature, vitreous chamber depth, axial length and, lens thickness are widely analyzed when studying eye diseases. However, among these com-ponents, more attention is paid to the axial length which is the major parameter for both hyper myopia and myopia.(7) Furthermore, studies have shown that the alteration of en-vironmental factors and the identification of genetic correlations may play a significant role in axial elongation, myopia progression, and future ocular complications.(8,9)

Myopia is known to be the cause of multiple eye fatalities around the globe and in-vestigation conducted on a particular population in various hospital have shown that when the axial length or refractive error is ~ 26.5 mm or -8.00 D, the parapapillary atrophy and optic disc becomes enlarged gradually. However, when these values are higher, the prevalence of glaucomatous optic neuropathy and myopic retinopathy is increased. My-opia is identified as the excessive and continuous expansion of the axial length, result-ing in a change in the secondary fundus which leads to visual impairment, as well as cho-roidal neovascularization, retinal detachment, zonal areas of chorioretinal atrophy, my-opic macular schisis, and hole.(10)

In high myopic eyes, recent changes start in the choroid therefore, studies have shown that the choroid is a very valuable structure that is required in the pathophysiology of high myopia.(11) The choroidal vasculature helps nourish the outer retina (including the photoreceptors), however, when there is a loss of the vascular tissue and an extreme thinning of the choroid, it leads to visual impairment and damage to the photoreceptors. The thickness of the choroidal is an essential parameter used for studying the pathogene-sis of high myopia.(12) Furthermore, measurement of CT in vivo is suitable for determin-ing the onset of diseases and their progression which causes thinning of the choroidal.(13) The presence of lacquer cracks and choroidal neovascularization (CNV) is seen mostly in eyes with thinner macular choroids.(14) Therefore, this study aims to determine the rela-tionship between the degree of myopia, the axial length, and the choroidal thickness.

2. Materials and Methods

2.1 Study design

This is an observational analytical study that made use of a cross-sectional design. ~~A total of 59 participants were included. There are 49 patients with refractive errors and underwent treatment at Hasanuddin University Hospital and 10 normal subjects as a control group. A total of 116 eyes were measured and analyzed. In this study, our criteria for recruiting subjects were patients aged 20-50 years, had a refractive error ≥ 0.5 D, did not suffer from anterior segment abnormalities during examination, no history of eye infec-tion or eye surgery. A total of 49 participants with refractive errors, underwent treatment at Hasanuddin University Hospital and 69 eyes were measured and analyzed.~~ Each pa-tient who meets the criteria was asked to fill out and sign an informed consent form after then examined according to applicable standards.

2.2 Ophthalmology examination

The examination was carried out by measuring the patient's visual acuity and correction using a Snellen chart projector, trial lens, and retinoscopy. The results obtained from retinoscopy were in the form of a spherical equivalent (SE). The anterior segment of the eye was then examined with a slit lamp biomicroscope, then the patient was made to undergo an indirect funduscopy examination with a 90D ocular lens to view the posterior segment of the eye. Subsequently, the length of the patient's eyeball axis was obtained using A-Scan Ultrasound which measures from the top of the cornea to the posterior segment, and the results were represented in millimeters (mm). The choroidal thickness was measured using the Enhance Depth Imaging OCT (EDI SD-OCT) tool which is divided into nine observation areas. [The type of OCT used is Heidelberg Spectralis® OCT \(3-mode\), Germany. It was carried out by 2 technicians \(two graders\) then the results were blindly confirmed by the research team \(HSM and AMI\).](#) ~~The test~~ was performed semi-automatically by drawing a perpendicular line between the outermost part of the hyper-reflective line that represents the RPE with the hypo-reflective line that represents the choroid-scleral surface using a calipers software tool.

2.3 Data analysis and interpretation


All results obtained were recorded and compared [using SPSS for windows ver. 24.0 using the with](#) One-way ANOVA and ~~Spearman-Pearson~~ correlation test (sig. $p < 0.05$) which was represented in tables and figures. [The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Ethics Committee for Medical Research of Hasanuddin University \(Approval No.: 751/UN 4.6.4.5.31 /PP36/2021\).](#)

3. Results


The degree of myopia is divided into 3 categories; low (≤ -3.00 D), moderate (~~≥ -3.25~~ 0.00 to ≤ -6.00 D), and high (> -6.00 D) myopias. This was compared with the choroidal thickness found in various areas of the macula. The result obtained showed a significant difference except for the 1.5 mm temporal region (T3) (Table 1). This result indicates a significant relationship between the choroidal thickness and the degree of myopia. A high degree myopia showed a thinner choroidal vasculature.

From the horizontal image (Fig.1), the choroid found in the low myopia is thickest in the subfoveal unlike in medium and high myopia where it is thickest in the temporal. The thinnest area in all groups was the nasal area. Vertically (Fig.2), in low and moderate myopia, the choroid is thickest in the superior area while in high myopia it is thickest in the subfoveal area. The thinnest area in each group was the inferior region. Comparison of the choroidal thickness based on the axial length showed a significant difference ($p < 0.05$), except for the superior region (S6).

96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133



134



135

Figure 1. Choroidal thickness based on degrees of myopia horizontally.
(A) Normal subject (B) Low myopia, (C) Moderate myopia, (D) High myopia.
Figure 1. Choroidal thickness based on degrees of myopia horizontally.
(A) Low myopia 1.75D, (B) Moderate myopia 3.75D, (C) High myopia 10.25D

136

137

138

139

140





Figure 2. Choroidal thickness based on degrees of myopia vertically. (A) Normal subject (B) Low myopia, (C) Moderate myopia, (D) High myopia

141

142

143 Formatted: Indent: First line: 0.9 cm

144 Formatted: Centered

145

Figure 2. Choroidal thickness based on degrees of myopia vertically—
(A) Low myopia 1.75D, (B) Moderate myopia 3.75D, (C) High myopia 10.25D

Table 1. Comparison of choroidal thickness and the degrees of myopia

Choroidal Thickness based on area (Mean ± SD) μm	Myopia degree (D)				p
	Normal (n=20)	Low myopia (n=50)	Moderate myopia (n=31)	High myopia (n=15)	
C=Sub Fovea	296.70±69.62	307.52±79.30	253.39±64.84	267.33±104.01	0.010
T6=Temporal 3	273.35±46.82	280.22±76.18	255.06±47.20	231.07±59.52	0.028
T3=Temporal1.5	295.45±66.94	298.12±78.30	262.58±54.02	267.80±83.84	0.076
N3=Nasal 1.5	274.15±71.70	252.50±64.89	208.23±48.70	196.07±89.16	0.002
N6=Nasal 3	219.25±77.72	173.00±50.72	151.06±47.20	132.33±77.61	0.027
S6=Superior 3	319.90±84.54	306.02±72.50	269.13±56.69	258.67±89.28	0.022
S3=Superior 1.5	299.40±76.45	314.18±80.06	261.52±52.95	257.73±93.38	0.003
I3=Inferior 1.5	293.95±77.33	307.12±71.74	249.97±62.72	244.73±76.31	<0.001
I6=Inferior 3	278.50±86.19	282.04±62.99	237.84±47.31	226±65.93 ^{bb}	0.001

*One-way Anova test

The mean choroid thickness of the ~~short-normal~~ (<22.5 mm), medium (22.5 to 25.0 mm) and long (>25.5 mm) axis in the horizontal section (Fig.3) were thickest in the fovea, while the thinnest in each group was found in the nasal area. Furthermore, vertical measurement (Fig.4) on the ~~short-normal~~ axis, showed that the thickest area is on subfoveal area while on the medium and long axis it is thickest in the superior area. The thinnest area in the ~~short-normal~~ and medium axis groups was the inferior region, while for the long axis it was the sub fovea area.

146
147
148
149

Formatted Table

150
151
152
153
154
155
156
157
158
159
160




Figure 3. Choroidal thicknes based on axial length horizontally.

(A) ~~Short axis Normal axis (22,47 mm)~~, (B) Medium axis (23,75 mm), (C) Long axis (27,60 mm)




Figure 4. Figure 3. Choroidal thicknes based on axial length vertically.

(A) ~~Short axis Normal axis (22,47 mm)~~ (B) Medium axis (23,75 mm), (C) Long axis (27,60 mm)

Table 2 showed that there is a significant correlation between choroidal thickness and axial length ($p < 0,05$), except on S6 area. This indicates that the thickness of the choroid decreases with increasing axial length.

161
162
163
164
165

166
167
168
169
170
171
172
173
174

Formatted: Centered

Table 2. Comparison of choroidal thickness and axial length

Coroidal thickness based on area (Mean±SD) µm	Axial length length (mm)			p*
	Short <u>Normal</u> (n=4)	Moderate (n=74)	Long (n=18)	
C=Sub Fovea	345,00±16,31	294,04±79,83	227,89±77,30	0.002
T6=Temporal 3	347,25±73,3	270,20±65,20	222,22±51,0	0.001
T3=Temporal 1,5	328,75±35,76	289,12±75,56	241,83±55,25	0.020
N3=Nasal 1,5	307±39,43	233,09±65,93	196,89±68,55	0.008
N6=Nasal 3	247,25±49,14	160,69±52,17	135,44±55,50	0.001
S6=Superior 3	332,75±49,25	291,73±71,58	255,83±74,45	0.074
S3=Superior 1,5	336±31,76	298,03±76,48	238±76,04	0.006
I3=Inferior 1,5	330,5±41,06	287,24±73,33	233,22±70,31	0.008
I6=Inferior 3	311,25±14,7	264,58±63,82	224,5±52,34	0.012

*One-way ANOVA test

Table 3 shows the correlation coefficient between the axial length of the eyeball and the thickness of the choroid (r= 0.270 – 453) which is higher than the degree of myopia (r= 0.230 – 407).

Table 3. Correlation between axial length, degree of myopia and choroidal thickness in various regions

Choroidal Thickness based on area (Mean±SD) µm	Axial length (mm)		Myopia degree (D)	
	Coefficient correlation (r)	p*	Coefficient correlation (r)	p*
C=Sub Fovea	-0,400	<0,001	-0,274	0.003
T6=Temporal 3	-0,385	<0,001	-0,317	0.001
T3=Temporal1,5	-0,307	<0,001	-0,230	0.012
N3=Nasal 1,5	-0,427	0,001	-0,361	<0,001
N6=Nasal 3	-0,343	<0,001	-0,332	<0,001
S6=Superior 3	-0,270	<0,001	-0,300	0.002

175
176

177
178
179
180
181
182
183
184

S3=Superior 1,5	-0,389	0,004	-0,308	0.001
I3=Inferior 1,5	-0,453	<0,001	-0,388	<0.001
I6=Inferior 3	-0,431	<0,001	-0,407	<0.001

***Pearson correlation test**

The choroidal thickness showed a significant relationship ($p < 0.05$) with the axial length and degree of myopia found in various areas of the macula and this has a negative correlation coefficient; Therefore, means, the longer the axis of the eyeball the higher the degree of myopia which will lead to choroidal thinning (Fig.5).




Figure 5. (A) Correlation between choroidal thickness and myopia degree (B) Correlation between choroidal thickness and axial length.

4. Discussion

Historically, some opticians have thought that myopia is a hereditary abnormality, whereas others have imagined it to be environmentally induced. However, several studies conducted on animals and humans over the last four decades have suggested that the occurrence of myopia is controlled by both genetic and environmental factors.(3)

The comparison between the degrees of myopia and choroidal thickness showed a significant correlation ($p < 0.05$) that indicates that a higher degree of myopia will lead to thinning of the choroidal layer. Moreover, the mean choroidal thickness obtained by OCT examination was based on the degree of myopia found in the horizontal area. The low degree myopia was found to be thickest in the subfoveal area, while the moderate and high degree myopia was thickest in the temporal region. The thinnest choroid was found in the nasal area of all degrees of myopia.

Based on studies conducted, it is known that the thinning of the choroid reduces ischemia and perfusion of the choroid which leads to the upregulation of the angiogenic factors in the eyes and this may also lead to the formation of myopic choroidal neovascularization and other features of macular degeneration.(15)

In a cross-sectional study by El-Shazly *et al.* (2017) Macular CT was measured in different degrees of myopia and in normal control eyes, and a similar result was obtained which is significantly lower in myopes than in emmetropes. Moreover, it varies by location, where the thickest CT in low myopic eyes, is found in the subfoveal area, while the thinnest is located in the nasal region. However, for eyes with moderate myopia, the thickest is found in the temporal region while the thinnest region remained in the nasal direction.(16)

Another study by Deng *et al.* (2018) showed that the mean CT in the perifoveal, parafoveal, and central foveal regions where $215 \pm 50 \mu\text{m}$, $227 \pm 60 \mu\text{m}$, and $229 \pm 65 \mu\text{m}$

185
186
187
188
189
190
191

192
193
194
195

196

197
198
199
200

201
202
203
204
205
206
207

208
209
210
211

212
213
214
215
216
217
218

219
220

respectively while the mean spherical equivalent (SE) of the patient was -1.71 ± 2.22 diopter (D) (range from -7.63 to 4.25 D). Furthermore, the mean global peripapillary choroidal thickness (PPCT) was 136 ± 33 μm .(17)

Based on the result obtained from different studies carried out on animals, an alteration in the thickness of the choroid may occur when maintaining a clear vision. Earlier studies on macaques, marmosets, and chicks have led to the hypothesis which states that the thickening of the choroid may occur when myopic defocus is induced due to changes in the position of the retina when maintaining a clear vision. This is possible because, in myopic defocus, the retina is at the back of the image plane so when thickening of the choroid occurs it moves the retina forward.(11)

Furthermore, this study also indicates that an increase in axial length leads to a decrease in the thickness of the choroid. (Table 2). Moreover, the choroidal thickness showed a significant relationship ($p < 0.05$) between the axial length and the degree of myopia found in various areas of the macula that has a negative correlation coefficient. Therefore, this indicates that the longer the axis of the eyeball and the higher the degree of myopia the more choroidal thinning will occur.

In general, the Axial length increases rapidly at the early stage of life but slowly in adulthood and decreases in old age.(7) According to Lee *et.al* (2020) the best-corrected visual acuity (BCVA), baseline axial length, anterior chamber depth, and age were significantly associated with changes in axial length ($p = 0.005$, $p < 0.001$, $p = 0.006$, and $p = 0.045$ respectively).(18)

Furthermore, the choroid may stimulate axial growth by regulating the remodeling of the scleral extracellular matrix, which is important for emmetropization during eye formation. In experimental animals induced with hyperopia and myopia, a change in the thickness of the choroid exceeds that of axial length and scleral remodeling.(12)

According to histologic, clinical, and population-based investigations, it was shown that an increase in axial elongation led to a significant thinning of the choroid. Furthermore, in emmetropic subjects, the mean CT was 250 μm while in highly axially myopic patients it decreases to < 30 μm . Therefore, this indicates that an increase in the axial elongation led to a decrease in the distance between the Bruch membrane and sclera.(16)

Based on Jin et al. (2016), the myopic retinas were thinner than those of emmetropic or hyperopic subjects, especially in the superior parafoveal and all 4 perifoveal subfields ($P < 0.05$). But, the results of previous studies on factors influencing the thickness of the ganglion cell layer and nerve fiber layer have been conflicting. While some suggested that the thickness of the ganglion cell layer and peripapillary nerve fiber layer is correlated with spherical equivalent refraction and axial length in adults, others did not observe this relationship.(12).

Karahan (2013) reported that choroidal change plays a major role in the development and progression of many retinal disease. Thickening of the choroid could affect the nutrition supply to the retina, because outer retina layer is nourished by the choroidal vasculature. Thus, choroidal thickness provides useful information to clinicians.(19)

An increase in the axial length led to an increase in the retina thickness which aid in blood supply. This, however, caused an increase in length of the ocular axis resulting in a compensatory thickening of the choroidal capillary layer in the fovea centralis and an increase in the number of capillaries and volume. Moreover, an increase in the myopia degree will cause the axial length of the eye to increase as well. However, the retina may fail to compensate causing the capillary layer and choroid to become thinner.(20)

5. Conclusions

221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Justified, Indent: First line: 0.8 cm, Space Before: 0 pt, After: 0 pt, Line spacing: single

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Field Code Changed

There was a significant correlation between the degree of myopia and choroidal thickness. Therefore, a higher degree of myopia will cause a decrease in the thickness of the choroid. Furthermore, it is necessary to screen for choroidal thickness in myopic patients using SD-OCT, especially in moderate and high degrees to prevent complications, because a significant increase in myopia degree will lead to a decrease in the choroidal thickness. Also, it is necessary to measure the length of the eyeball axis in myopic patients because a decrease in choroidal thickness is significant to the elongation of the eyeball. All of these examinations are to prevent some further complication such as the progressivity of myopic choroidal neovascularization (CNV) which will eventually lead to worsening of vision.

Author Contributions: HSM, ARM, AMI: Conception and design of the work, performing the medical examination, analysis and interpretation of data, drafting the work. BTU, JS, IP, AMI: ~~performing the medical examination, giving~~ Giving supervision and quality control of the medical examination as a team, review and editing of the manuscript draft. ICI: project administration, drafting the work and revising it for publication.

Funding: This research received no external funding

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Ethics Committee for Medical Research of Hasanuddin University (Approval No.: 751/UN 4.6.4.5.31 /PP36/2021).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgments: All authors would like to thank and appreciate the team of nurses and staff of Hasanuddin University Hospital who were involved in the preparation, examination and follow up of our patients. We also thank Goodlingua for English preparation review.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Kaur K, Gurnani B, Kannusamy V. Myopia: Current concepts and review of literature. TNOA J Ophthalmic Sci Res. 2020;58(4):280.
2. Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, et al. Global Prevalence of Myopia and High Myopia and Temporal Trends from 2000 through 2050. Ophthalmology [Internet]. 2016;123(5):1036–42. Available from: <http://dx.doi.org/10.1016/j.ophtha.2016.01.006>
3. Cooper J, Tkatchenko A V. A Review of Current Concepts of the Etiology and Treatment of Myopia. Eye Contact Lens. 2018;44(4):231–47.
4. Fredrick DR. Myopia. Br Med J. 2002;324(7347):1195–9.
5. Kaiti R, Shyangbo R, Sharma IP, Dahal M. Review on current concepts of myopia and its control strategies. Int J Ophthalmol. 2021;14(4):606–15.
6. Okafor M, O FO, I EB. Myopia : a Review of Literature Aetiology Progression of Myopia. Niger J Med. 2009;18(2):134–8.
7. Meng W, Butterworth J, Malecaze F, Calvas P. Axial length of myopia: A review of current research. Ophthalmologica. 2011;225(3):127–34.

Formatted: Indent: Left: 5.4 cm

8. Pugazhendhi S, Ambati B, Hunter AA. Pathogenesis and prevention of worsening axial elongation in pathological myopia. *Clin Ophthalmol*. 2020;14:853–73. 320
321
9. Yotsukura E, Torii H, Ozawa H, Hida RY, Shiraishi T, Corso Teixeira I, et al. Axial length and prevalence of myopia among schoolchildren in the equatorial region of brazil. *J Clin Med*. 2021;10(1):1–11. 322
323
324
10. Teberik K, Kaya M. Retinal and Choroidal Thickness in Patients with High Myopia without Maculopathy. *Ocul Immunol Inflamm*. 2017;25(2):1438–9. 325
326
11. Lee GY, Yu S, Kang HC, Kim JS, Lee KW, Lee J-H. Choroidal Thickness Variation According to Refractive Error Measured by Spectral Domain-optical Coherence Tomography in Korean Children. *Korean J Ophthalmol*. 2017;31(2):151. 327
328
329
12. Jin P, Zou H, Zhu J, Xu X, Jin J, Chang TC, et al. Choroidal and Retinal Thickness in Children with Different Refractive Status Measured by Swept-Source Optical Coherence Tomography. *Am J Ophthalmol* [Internet]. 2016;168:164–76. Available from: <http://dx.doi.org/10.1016/j.ajo.2016.05.008> 330
331
332
333
13. Fujiwara T, Imamura Y, Margolis R, Slakter JS, Spaide RF. Enhanced Depth Imaging Optical Coherence Tomography of the Choroid in Highly Myopic Eyes. *Am J Ophthalmol* [Internet]. 2009;148(3):445–50. Available from: <http://dx.doi.org/10.1016/j.ajo.2009.04.029> 334
335
336
337
14. Wang S, Wang Y, Gao X, Qian N, Zhuo Y. Choroidal thickness and high myopia: A cross-sectional study and meta-analysis *Retina*. *BMC Ophthalmol* [Internet]. 2015;15(1):1–10. Available from: <http://dx.doi.org/10.1186/s12886-015-0059-2> 338
339
340
15. Xiong S, He X, Zhang B, Deng J, Wang J, Lv M, et al. Changes in Choroidal Thickness Varied by Age and Refraction in Children and Adolescents: A 1-Year Longitudinal Study. *Am J Ophthalmol* [Internet]. 2020;213:46–56. Available from: <https://doi.org/10.1016/j.ajo.2020.01.003> 341
342
343
344
16. El-Shazly AA, Farweez YA, Elsebaay ME, El-Zawahry WMA. Correlation between choroidal thickness and degree of myopia assessed with enhanced depth imaging optical coherence tomography. *Eur J Ophthalmol*. 2017;27(5):577–84. 345
346
347
17. Deng J, Li X, Jin J, Zhang B, Zhu J, Zou H, et al. Distribution Pattern of Choroidal Thickness at the Posterior Pole in Chinese Children With Myopia. *Invest Ophthalmol Vis Sci*. 2018;59(3):1577–86. 348
349
350
18. Lee MW, Lee SE, Lim H Bin, Kim JY. Longitudinal changes in axial length in high myopia: A 4-year prospective study. *Br J Ophthalmol*. 2020;104(5):600–3. 351
352
19. Karahan E, Zengin MO, Tuncer I. Correlation of choroidal thickness with outer and inner retinal layers. *Ophthalmic Surg Lasers Imaging*. 2013;44(6):544–8. 353
354
20. Lin TN, Yang Y, Lin JH, Zhang JH, Wen Q, He XL, et al. A comparative study of macular and choroidal thickness and blood-flow parameters in patients with intermediate and simple juvenile moderate myopia. *Int J Gen Med*. 2021;14:1343–8. 355
356
357
358