

BAB VII

KESIMPULAN

- Kombinasi PRP dan SVFs adalah terapi yang efektif untuk penyembuhan luka bakar deep dermal dibandingkan dengan terapi konvensional menggunakan vaselin. Hal tersebut disebabkan oleh kemampuan agen terapi meningkatkan kadar growth factor dan sifat steemcell yang ada pada SVF sangat berpengaruh terhadap proses Re-epitelisasi, selain itu jalur administrasi regiment turut berpengaruh terhadap efektifitas terapi.
- Kadar FGF2 tertinggi didapatkan pada kelompok dengan jalur administrasi injeksi, memberikan pengaruh sebesar 90,3% atau sebesar 1.836x dibandingkan dengan jalur topikal yang memberikan pengaruh sebesar 76,6% (1.179x)
- Kadar EGF tertinggi didapatkan pada kelopok perlakuan dengan jalur administrasi injeksi yang memberikan pengaruh sebesar 99,8% atau sebesar 6.696x dibandingkan dengan pemberian secara topikal yang memberikan yang memberikan pengaruh sebesar 99,6% (5.314x).
- Proses reepitelisasi didapatkan ke 4 sample di kelompok perlakuan injeksi stemcell kombinasi prp + SVFs dan pada kelompok perlakuan topikal stemcell hanya 2 sampel yang berhasil sampai pada reepitelisasi namun pada sample perlakuan kontrol dengan vaselin tidak ada sampel yang berhasil mencapai proses Re-epitelisasi.
- Penelitian lebih lanjut, khususnya pada manusia dapat dilakukan agar dapat menjadi terapi pilihan yang dapat digunakan dalam praktik klinis sehari-hari.

Referensi

- Abdel-Sayed, P., Michetti, M., Scaletta, C., Flahaut, M., Hirt-Burri, N., de Buys Roessingh, A., Raffoul, W., Applegate, L.A., 2019. Cell therapies for skin regeneration: an overview of 40 years of experience in burn units. *Swiss Med. Wkly.* 149, w20079. <https://doi.org/10.4414/smw.2019.20079>
- Abraham, J.A., Klagsbrun, M., 1988. Modulation of Wound Repair by Members of the Fibroblast Growth Factor Family, in: *The Molecular and Cellular Biology of Wound Repair*. Springer US, Boston, MA, pp. 195–248. https://doi.org/10.1007/978-1-4899-0185-9_6
- Abu-Humaidan, A.H.A., Ananthoju, N., Mohanty, T., Sonesson, A., Alberius, P., Schmidtchen, A., Garred, P., Sørensen, O.E., 2014. The Epidermal Growth Factor Receptor Is a Regulator of Epidermal Complement Component Expression and Complement Activation. *J. Immunol.* 192, 3355 LP – 3364. <https://doi.org/10.4049/jimmunol.1302305>
- Akhundov, K., Pietramaggiori, G., Waselle, L., Darwiche, S., Guerid, S., Scaletta, C., Hirt-Burri, N., Applegate, L.A., Raffoul, W. V, 2012. Development of a cost-effective method for platelet-rich plasma (PRP) preparation for topical wound healing. *Ann. Burns Fire Disasters* 25, 207–213.
- Aleckovic, M., Simon, C., 2008. Is teratoma formation in stem cell research a characterization tool or a window to developmental biology? *Reprod. Biomed. Online* 17, 270–280. [https://doi.org/10.1016/s1472-6483\(10\)60206-4](https://doi.org/10.1016/s1472-6483(10)60206-4)
- ANZBA, 2016. Emergency Management of Severe Burns (EMSB), 18th ed. THE EDUCATION COMMITTEE OF AUSTRALIA AND NEW ZEALAND BURN ASSOCIATION LTD, Albany Creek.
- Baglioni, S., Francalanci, M., Squecco, R., Lombardi, A., Cantini, G., Angeli, R., Gelmini, S., Guasti, D., Benvenuti, S., Annunziato, F., Bani, D., Liotta, F., Francini, F., Perigli, G., Serio, M., Luconi, M., 2009. Characterization of human adult stem-cell populations isolated from visceral and subcutaneous adipose tissue. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 23, 3494–3505. <https://doi.org/10.1096/fj.08-126946>
- Baird, A., 1997. The Regulation of Basic Fibroblast Growth Factor (FGF2-2) Through Limited Bioavailability, in: Ziegler, T.R., Pierce, G.F., Herndon, D.N. (Eds.), *Growth Factors and Wound Healing*. Springer New York, New York, NY, pp. 27–36.
- Bakacak, M., Bostancı, M.S., Inanc, F., Yaylali, A., Serin, S., Attar, R., Yildirim, G., Yildirim, O.K., 2016. Protective Effect of Platelet Rich Plasma on Experimental Ischemia/Reperfusion Injury in Rat Ovary. *Gynecol. Obstet. Invest.* 81, 225–231.

- <https://doi.org/10.1159/000440617>
- Barret-Nerin, J., 2004. Principles and Practice of Burn Surgery. CRC Press.
<https://doi.org/10.1201/b21634>
- Benson, A., Dickson, W.A., Boyce, D.E., 2006. ABC of wound healing: Burns. BMJ 333, 609324. <https://doi.org/10.1136/sbmj.0609324>
- Borrione, P., Gianfrancesco, A. Di, Pereira, M.T., Pigozzi, F., 2010. Platelet-rich plasma in muscle healing. Am. J. Phys. Med. Rehabil. 89, 854–861.
<https://doi.org/10.1097/PHM.0b013e3181f1c1c7>
- Bourin, P., Bunnell, B.A., Casteilla, L., Dominici, M., Katz, A.J., March, K.L., Redl, H., Rubin, J.P., Yoshimura, K., Gimble, J.M., 2013. Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International So. Cytotherapy 15, 641–648. <https://doi.org/10.1016/j.jcyt.2013.02.006>
- Cerqueira, M.T., Pirraco, R.P., Marques, A.P., 2016. Stem Cells in Skin Wound Healing: Are We There Yet? Adv. Wound Care 5, 164–175.
<https://doi.org/10.1089/wound.2014.0607>
- Cervelli, V., De Angelis, B., Lucarini, L., Spallone, D., Balzani, A., Palla, L., Gentile, P., Cerulli, P., 2010. Tissue regeneration in loss of substance on the lower limbs through use of platelet-rich plasma, stem cells from adipose tissue, and hyaluronic acid. Adv. Skin Wound Care 23, 262–272.
<https://doi.org/10.1097/01.ASW.0000363551.82058.36>
- Chieregato, K., Castegnaro, S., Madeo, D., Astori, G., Pegoraro, M., Rodeghiero, F., 2011. Epidermal growth factor, basic fibroblast growth factor and platelet-derived growth factor-bb can substitute for fetal bovine serum and compete with human platelet-rich plasma in the ex vivo expansion of mesenchymal stromal cells derived from adipose tiss. Cytotherapy 13, 933–943.
<https://doi.org/10.3109/14653249.2011.583232>
- Choi, J., Minn, K.W., Chang, H., 2012. The efficacy and safety of platelet-rich plasma and adipose-derived stem cells: an update. Arch. Plast. Surg. 39, 585–592.
<https://doi.org/10.5999/aps.2012.39.6.585>
- Comella, K., Silbert, R., Parlo, M., 2017. Effects of the intradiscal implantation of stromal vascular fraction plus platelet rich plasma in patients with degenerative disc disease. J. Transl. Med. 15, 12. <https://doi.org/10.1186/s12967-016-1109-0>
- Dailey, L., Ambrosetti, D., Mansukhani, A., Basilico, C., 2005. Mechanisms underlying differential responses to FGF2 signaling. Cytokine Growth Factor Rev. 16, 233–247. <https://doi.org/10.1016/j.cy togfr.2005.01.007>
- Darinskas, A., Paskevicius, M., Apanavicius, G., Vilkevicius, G., Labanauskas, L.,

- Ichim, T.E., Rimdeika, R., 2017. Stromal vascular fraction cells for the treatment of critical limb ischemia: a pilot study. *J. Transl. Med.* 15, 143. <https://doi.org/10.1186/s12967-017-1243-3>
- Deng, H., Coumans, J.V., Anderson, R., Houle, T.T., Peterfreund, R.A., 2019. Spinal anesthesia for lumbar spine surgery correlates with fewer total medications and less frequent use of vasoactive agents: A single center experience. *PLoS One* 14, 1–19. <https://doi.org/10.1371/journal.pone.0217939>
- Desai, T.J., Cardoso, W. V, 2002. Growth factors in lung development and disease: friends or foe? *Respir. Res.* 3, 2. <https://doi.org/10.1186/rr169>
- El-Sharkawy, H., Kantarci, A., Deady, J., Hasturk, H., Liu, H., Alshahat, M., Van Dyke, T.E., 2007. Platelet-rich plasma: growth factors and pro- and anti-inflammatory properties. *J. Periodontol.* 78, 661–669. <https://doi.org/10.1902/jop.2007.060302>
- Eppley, B.L., Pietrzak, W.S., Blanton, M., 2006. Platelet-rich plasma: a review of biology and applications in plastic surgery. *Plast. Reconstr. Surg.* 118, 147e-159e. <https://doi.org/10.1097/01.prs.0000239606.92676.cf>
- Esquirol Caussa, J., Herrero Vila, E., 2015. Epidermal growth factor, innovation and safety. *Med. Clínica (English Ed.)* 145, 305–312. <https://doi.org/10.1016/j.medcle.2016.02.028>
- Eswarakumar, V.P., Lax, I., Schlessinger, J., 2005. Cellular signaling by fibroblast growth factor receptors. *Cytokine Growth Factor Rev.* 16, 139–149. <https://doi.org/10.1016/j.cytogfr.2005.01.001>
- Eyuboglu, A.A., Uysal, C.A., Ozgun, G., Coskun, E., Markal Ertas, N., Haberal, M., 2018. The effect of adipose derived stromal vascular fraction on stasis zone in an experimental burn model. *Burns* 44, 386–396. <https://doi.org/10.1016/j.burns.2017.08.016>
- Falanga, V., Eaglstein, W.H., Bucalo, B., Katz, M.H., Harris, B., Carson, P., 1992. Topical use of human recombinant epidermal growth factor (h-EGF) in venous ulcers. *J. Dermatol. Surg. Oncol.* 18, 604–606. <https://doi.org/10.1111/j.1524-4725.1992.tb03514.x>
- Ferraro, G.A., Mizuno, H., Pallua, N., 2016. Adipose Stem Cells: From Bench to Bedside. *Stem Cells Int.* 2016, 6484038. <https://doi.org/10.1155/2016/6484038>
- Finch, P.W., Rubin, J.S., Miki, T., Ron, D., Aaronson, S.A., 1989. Human KGF is FGF2-Related with Properties of a Paracrine Effector of Epithelial Cell Growth. *Science (80-.).* 245, 752–755.
- Fotouhi, A., Maleki, A., Dolati, S., Aghebati-Maleki, A., Aghebati-Maleki, L., 2018. Platelet rich plasma, stromal vascular fraction and autologous conditioned serum in treatment of knee osteoarthritis. *Biomed. Pharmacother.* 104, 652–660. <https://doi.org/10.1016/j.biopharm.2018.05.019>

- Foubert, P., Gonzalez, A.D., Teodosescu, S., Berard, F., Doyle-Eisele, M., Yekkala, K., Tenenhaus, M., Fraser, J.K., 2016. Adipose-derived regenerative cell therapy for burn wound healing: A comparison of two delivery methods. *Adv. Wound Care* 5, 288–298. <https://doi.org/10.1089/wound.2015.0672>
- Fu, Y.-S., Cheng, Y.-C., Lin, M.-Y.A., Cheng, H., Chu, P.-M., Chou, S.-C., Shih, Y.-H., Ko, M.-H., Sung, M.-S., 2006. Conversion of human umbilical cord mesenchymal stem cells in Wharton's jelly to dopaminergic neurons in vitro: potential therapeutic application for Parkinsonism. *Stem Cells* 24, 115–124. <https://doi.org/10.1634/stemcells.2005-0053>
- Gentile, P., Scioli, M.G., Bielli, A., Orlandi, A., Cervelli, V., 2017. Concise Review: The Use of Adipose-Derived Stromal Vascular Fraction Cells and Platelet Rich Plasma in Regenerative Plastic Surgery. *Stem Cells* 35, 117–134. <https://doi.org/10.1002/stem.2498>
- Ghadially, R., Halkier-Sorensen, L., Elias, P.M., 1992. Effects of petrolatum on stratum corneum structure and function. *J. Am. Acad. Dermatol.* 26, 387–396. [https://doi.org/https://doi.org/10.1016/0190-9622\(92\)70060-S](https://doi.org/https://doi.org/10.1016/0190-9622(92)70060-S)
- Ghieh, F., Jurjus, R., Ibrahim, A., Geagea, A.G., Daouk, H., El Baba, B., Chams, S., Matar, M., Zein, W., Jurjus, A., 2015. The Use of Stem Cells in Burn Wound Healing: A Review. *Biomed Res. Int.* 2015, 684084. <https://doi.org/10.1155/2015/684084>
- Gillenwater, J., Garner, W.L., 2020. Thermal, Chemical, and Electrical Injuries, in: Kevin C. Chung (Ed.), *Grabb and Smith's Plastic Surgery*. Wolters Kluwer Health, Philadelphia.
- Gimble, J., Guilak, F., 2003. Adipose-derived adult stem cells: isolation, characterization, and differentiation potential. *Cyotherapy* 5, 362–369. <https://doi.org/10.1080/14653240310003026>
- Gimble, J.M., Katz, A.J., Bunnell, B.A., 2007. Adipose-derived stem cells for regenerative medicine. *Circ. Res.* 100, 1249–1260. <https://doi.org/10.1161/01.RES.0000265074.83288.09>
- Goldfarb, M., 2005. Fibroblast growth factor homologous factors: evolution, structure, and function. *Cytokine Growth Factor Rev.* 16, 215–220. <https://doi.org/10.1016/j.cytogfr.2005.02.002>
- Grigore, T.V., Cozma, C., 2018. Platelet-rich plasma as a site-targeted approach in wound healing: a molecular perspective. *Discoveries* 6, 1–8. <https://doi.org/10.15190/d.2018.8>
- Guo, H.-F., Ali, R.M., Hamid, R.A., Zaini, A.A., Khaza'ai, H., 2017. A new model for studying deep partial-thickness burns in rats. *Int. J. Burns Trauma* 7, 107–114.
- Han, J., Koh, Y.J., Moon, H.R., Ryoo, H.G., Cho, C.-H., Kim, I., Koh, G.Y., 2010.

- Adipose tissue is an extramedullary reservoir for functional hematopoietic stem and progenitor cells. *Blood* 115, 957–964. <https://doi.org/10.1182/blood-2009-05-219923>
- Hardwicke, J., Schmaljohann, D., Boyce, D., Thomas, D., 2008. Epidermal growth factor therapy and wound healing--past, present and future perspectives. *Surgeon* 6, 172–177. [https://doi.org/10.1016/s1479-666x\(08\)80114-x](https://doi.org/10.1016/s1479-666x(08)80114-x)
- Harfouche, G., Vaigot, P., Rachidi, W., Rigaud, O., Moratille, S., Marie, M., Lemaitre, G., Fortunel, N.O., Martin, M.T., 2010. Fibroblast Growth Factor Type 2 Signaling Is Critical for DNA Repair in Human Keratinocyte Stem Cells. *Stem Cells* 28, 1639–1648. <https://doi.org/10.1002/stem.485>
- Harris, R.C., Chung, E., Coffey, R.J., 2003. EGF receptor ligands. *Exp. Cell Res.* 284, 2–13. [https://doi.org/10.1016/s0014-4827\(02\)00105-2](https://doi.org/10.1016/s0014-4827(02)00105-2)
- Harrison, D.A., 2012. The Jak/STAT pathway. *Cold Spring Harb. Perspect. Biol.* 4. <https://doi.org/10.1101/csdperspect.a011205>
- Hayes, M., Curley, G., Ansari, B., Laffey, J.G., 2012. Clinical review: Stem cell therapies for acute lung injury/acute respiratory distress syndrome - hope or hype? *Crit. Care* 16, 205. <https://doi.org/10.1186/cc10570>
- Herndon, D.N., Pierre, E.J., Rose, J.K., Stokes, K.N., Barrow, R.E., 1997. Growth Hormone Therapy in Human Burn Injury, in: Ziegler, T.R., Pierce, G.F., Herndon, D.N. (Eds.), *Growth Factors and Wound Healing*. Springer New York, New York, NY, pp. 231–244.
- Hirase, T., Ruff, E., Surani, S., Ratnani, I., 2018. Topical application of platelet-rich plasma for diabetic foot ulcers: A systematic review. *World J. Diabetes* 9, 172–179. <https://doi.org/10.4239/wjd.v9.i10.172>
- Holland, E.C., Varmus, H.E., 1998. Basic fibroblast growth factor induces cell migration and proliferation after glia-specific gene transfer in mice. *Proc. Natl. Acad. Sci. U. S. A.* 95, 1218–1223. <https://doi.org/10.1073/pnas.95.3.1218>
- Hombach-Klonisch, S., Panigrahi, S., Rashedi, I., Seifert, A., Alberti, E., Pocar, P., Kurpisz, M., Schulze-Osthoff, K., Mackiewicz, A., Los, M., 2008. Adult stem cells and their trans-differentiation potential--perspectives and therapeutic applications. *J. Mol. Med. (Berl.)* 86, 1301–1314. <https://doi.org/10.1007/s00109-008-0383-6>
- Hosni Ahmed, H., Rashed, L.A., Mahfouz, S., Elsayed Hussein, R., Alkaffas, M., Mostafa, S., Abusree, A., 2017. Can mesenchymal stem cells pretreated with platelet-rich plasma modulate tissue remodeling in a rat with burned skin? *Biochem. Cell Biol.* 95, 537–548. <https://doi.org/10.1139/bcb-2016-0224>
- IMAI, K., HIRAMATSU, A., FUKUSHIMA, D., PIERSCHBACHER, M.D., OKADA, Y., 1997. Degradation of decorin by matrix metalloproteinases: identification of the cleavage sites, kinetic analyses and transforming growth factor- β 1 release.

- Biochem. J. 322, 809–814. <https://doi.org/10.1042/bj3220809>
- Itoh, N., Ornitz, D.M., 2008. Functional evolutionary history of the mouse FGF2 gene family. Dev. Dyn. 237, 18–27. <https://doi.org/10.1002/dvdy.21388>
- Itoh, N., Ornitz, D.M., 2004. Evolution of the FGF2 and FGF2r gene families. Trends Genet. 20, 563–569. <https://doi.org/10.1016/j.tig.2004.08.007>
- Josh, F., Kobe, K., Tobita, M., Tanaka, R., Suzuki, K., Ono, K., Hyakusoku, H., Mizuno, H., 2012. Accelerated and safe proliferation of human adipose-derived stem cells in medium supplemented with human serum. J. Nippon Med. Sch. 79, 444–452. <https://doi.org/10.1272/jnms.79.444>
- Ju, W.D., Schiller, J.T., Kazempour, M.K., Lowy, D.R., 1993. TGF alpha enhances locomotion of cultured human keratinocytes. J. Invest. Dermatol. 100, 628–632.
- Karina, Samudra, M.F., Rosadi, I., Afini, I., Widystuti, T., Sobariah, S., Remelia, M., Puspitasari, R.L., Rosliana, I., Tunggadewi, T.I., 2019. Combination of the stromal vascular fraction and platelet-rich plasma accelerates the wound healing process: pre-clinical study in a Sprague-Dawley rat model. Stem cell Investig. 6, 18. <https://doi.org/10.21037/sci.2019.06.08>
- Kazakos, K., Lyras, D.N., Verettas, D., Tilkeridis, K., Tryfonidis, M., 2009. The use of autologous PRP gel as an aid in the management of acute trauma wounds. Injury 40, 801–805. <https://doi.org/10.1016/j.injury.2008.05.002>
- Kim, D.-Y., Ji, Y.-H., Kim, D.-W., Dhong, E.-S., Yoon, E.-S., 2014. Effects of platelet-rich plasma, adipose-derived stem cells, and stromal vascular fraction on the survival of human transplanted adipose tissue. J. Korean Med. Sci. 29 Suppl 3, S193–S200. <https://doi.org/10.3346/jkms.2014.29.S3.S193>
- Kirschstein, R., 2001. Stem cells: scientific progress and future directions. Natl. Institutes Heal. 1–222.
- Kolkova, K., Novitskaya, V., Pedersen, N., Berezin, V., Bock, E., 2000. Neural cell adhesion molecule-stimulated neurite outgrowth depends on activation of protein kinase C and the Ras-mitogen-activated protein kinase pathway. J. Neurosci. 20, 2238–2246. <https://doi.org/10.1523/JNEUROSCI.20-06-02238.2000>
- Komi-Kuramochi, A., Kawano, M., Oda, Y., Asada, M., Suzuki, M., Oki, J., Imamura, T., 2005. Expression of fibroblast growth factors and their receptors during full-thickness skin wound healing in young and aged mice. J. Endocrinol. 186, 273–289. <https://doi.org/10.1677/joe.1.06055>
- Kotani, T., Masutani, R., Suzuka, T., Oda, K., Makino, S., Ii, M., 2017. Anti-inflammatory and anti-fibrotic effects of intravenous adipose-derived stem cell transplantation in a mouse model of bleomycin-induced interstitial pneumonia. Sci. Rep. 7, 1–10. <https://doi.org/10.1038/s41598-017-15022-3>

- Lawrence, W.T., 1998. Physiology of the acute wound. *Clin. Plast. Surg.* 25, 321–340.
- Le, T.M., Morimoto, N., Mitsui, T., Notodihardjo, S.C., Munisso, M.C., Kakudo, N., Kusumoto, K., 2019. The sustained release of basic fibroblast growth factor accelerates angiogenesis and the engraftment of the inactivated dermis by high hydrostatic pressure. *PLoS One* 14, e0208658. <https://doi.org/10.1371/journal.pone.0208658>
- Li, H., Liu, D., Yu, Y., Wu, T., 2009. [Experimental research of the promotion effect of autogeneic PRP on osteogenic differentiation of human adipose-derived stem cells in vitro]. *Zhongguo xi fu chong jian wai ke za zhi = Zhongguo xiufu chongjian waike zazhi = Chinese J. reparative Reconstr. Surg.* 23, 732–736.
- Liang, Z.-J., Lu, X., Li, D.-Q., Liang, Y.-D., Zhu, D.-D., Wu, F.-X., Yi, X.-L., He, N., Huang, Y.-Q., Tang, C., Li, H., 2018. Precise Intradermal Injection of Nanofat-Derived Stromal Cells Combined with Platelet-Rich Fibrin Improves the Efficacy of Facial Skin Rejuvenation. *Cell. Physiol. Biochem.* 47, 316–329. <https://doi.org/10.1159/000489809>
- Lo, B., Parham, L., 2009. Ethical issues in stem cell research. *Endocr. Rev.* 30, 204–213. <https://doi.org/10.1210/er.2008-0031>
- MacLeod, A.S., Mansbridge, J.N., 2016. The Innate Immune System in Acute and Chronic Wounds. *Adv. wound care* 5, 65–78. <https://doi.org/10.1089/wound.2014.0608>
- Maddaluno, L., Urwyler, C., Werner, S., 2017. Fibroblast growth factors: key players in regeneration and tissue repair. *Development* 144, 4047–4060. <https://doi.org/10.1242/dev.152587>
- Mansoub, N.H., Gürdal, M., Karadadas, E., Kabadayi, H., Vatansever, S., Ercan, G., 2018. The role of PRP and adipose tissue-derived keratinocytes on burn wound healing in diabetic rats. *BioImpacts* 8, 5–12. <https://doi.org/10.15171/bi.2018.02>
- Marck, R., Middelkoop, M., Breederveld, R., 2018. Considerations on the use of platelet rich plasma, specifically for burn treatment: *Journal of burn care & research* May-Jun 2014;35(3):219-227, in: *On PLATELETS and Burns*. pp. 20–34.
- Marikovsky, M., Vogt, P., Eriksson, E., Rubin, J.S., Taylor, W.G., Sasse, J., Klagsbrun, M., 1996. Wound Fluid-Derived Heparin-Binding EGF-Like Growth Factor (HB-EGF) Is Synergistic with Insulin-Like Growth Factor-I For Balb/MK Keratinocyte Proliferation. *J. Invest. Dermatol.* 106, 616–621. <https://doi.org/10.1111/1523-1747.ep12345413>
- McGee, G.S., Davidson, J.M., Buckley, A., Sommer, A., Woodward, S.C., Aquino, A.M., Barbour, R., Demetriou, A.A., 1988. Recombinant basic fibroblast growth factor accelerates wound healing. *J. Surg. Res.* 45, 145–153. [https://doi.org/10.1016/0022-4804\(88\)90034-0](https://doi.org/10.1016/0022-4804(88)90034-0)

- Miranti, C.K., Brugge, J.S., 2002. Sensing the environment: a historical perspective on integrin signal transduction. *Nat. Cell Biol.* 4, E83–E90.
<https://doi.org/10.1038/ncb0402-e83>
- Moenadjat, Y., Merlina, M., Surjadi, C.F., Sardjono, C.T., Kusnadi, Y., Sandra, F., 2013. The application of human umbilical cord blood mononuclear cells in the management of deep partial thickness burn. *Med. J. Indones.* 22.
<https://doi.org/10.13181/mji.v22i2.534>
- Mohammed, M.O., Ali, O.J., Muhamad, S.A., Ibrahim, S.H., Raouf, G.M., Salih, N.A., 2019. Role of Bovine Fetal Platelet-rich Plasma (PRP) on Skin Wound Healing in Mice. *Int. J. Pharmacol.* 16, 18–26. <https://doi.org/10.3923/ijp.2020.18.26>
- Motamed, S., Taghiabadi, E., Molaei, H., Sodeifi, N., Hassanpour, S.E., Shafieyan, S., Azargashb, E., Farajzadeh-Vajari, F., Aghdami, N., Bajouri, A., 2017. Cell-based skin substitutes accelerate regeneration of extensive burn wounds in rats. *Am. J. Surg.* 214, 762–769. <https://doi.org/10.1016/j.amjsurg.2017.04.010>
- Nauta, A., Seidel, C., Deveza, L., Montoro, D., Grova, M., Ko, S.H., Hyun, J., Gurtner, G.C., Longaker, M.T., Yang, F., 2013. Adipose-derived stromal cells overexpressing vascular endothelial growth factor accelerate mouse excisional wound healing. *Mol. Ther.* 21, 445–455. <https://doi.org/10.1038/mt.2012.234>
- Nazzal, M., Osman, M.F., Albeshri, H., Abbas, D.B., Angel, C.A., 2019. Wound Healing, in: F. Charles Brunicardi, Anderse, D.K., Billiar, T.R., Dunn, D.L., Kao, L.S., Hunter, J.G., Matthews, J.B., Pollock, R.E. (Eds.), Schwartz's Principles of Surgery. McGraw-Hill, New York, pp. 271–304.
- Nexø, E., Jørgensen, P.E., Hansen, M.R., 1992. Human epidermal growth factor—on molecular forms present in urine and blood. *Regul. Pept.* 42, 75–84.
[https://doi.org/https://doi.org/10.1016/0167-0115\(92\)90025-P](https://doi.org/https://doi.org/10.1016/0167-0115(92)90025-P)
- Nikolidakis, D., Jansen, J.A., 2008. The biology of platelet-rich plasma and its application in oral surgery: literature review. *Tissue Eng. Part B. Rev.* 14, 249–258. <https://doi.org/10.1089/ten.teb.2008.0062>
- Oryan, A., Alemzadeh, E., Moshiri, A., 2017. Burn wound healing: Present concepts, treatment strategies and future directions. *J. Wound Care* 26, 5–19.
<https://doi.org/10.12968/jowc.2017.26.1.5>
- Park, J.W., Hwang, S.R., Yoon, I.-S., 2017. Advanced Growth Factor Delivery Systems in Wound Management and Skin Regeneration. *Molecules* 22, 1259.
<https://doi.org/10.3390/molecules22081259>
- Pastar, I., Stojadinovic, O., Yin, N.C., Ramirez, H., Nusbaum, A.G., Sawaya, A., Patel, S.B., Khalid, L., Isseroff, R.R., Tomic-Canic, M., 2014. Epithelialization in Wound Healing: A Comprehensive Review. *Adv. Wound Care* 3, 445–464.
<https://doi.org/10.1089/wound.2013.0473>

- Patel, D.P., Redshaw, J.D., Breyer, B.N., Smith, T.G., Erickson, B.A., Majercik, S.D., Gaither, T.W., Craig, J.R., Gardner, S., Presson, A.P., Zhang, C., Hotaling, J.M., Brant, W.O., Myers, J.B., 2015. High-grade renal injuries are often isolated in sports-related trauma. *Injury* 46, 1245–1249.
<https://doi.org/10.1016/j.injury.2015.02.008>
- Patel, P., Duttaroy, D., Kacheriwala, S., 2014. Management of renal injuries in blunt abdominal trauma. *J. Res. Med. Dent. Sci.* 2, 38.
<https://doi.org/10.5455/jrmlds.2014229>
- Pearson, G., Robinson, F., Beers Gibson, T., Xu, B.E., Karandikar, M., Berman, K., Cobb, M.H., 2001. Mitogen-activated protein (MAP) kinase pathways: regulation and physiological functions. *Endocr. Rev.* 22, 153–183.
<https://doi.org/10.1210/edrv.22.2.0428>
- Petry, T., Bury, D., Fautz, R., Hauser, M., Huber, B., Markowitz, A., Mishra, S., Rettinger, K., Schuh, W., Teichert, T., 2017. Review of data on the dermal penetration of mineral oils and waxes used in cosmetic applications. *Toxicol. Lett.* 280, 70–78. [https://doi.org/https://doi.org/10.1016/j.toxlet.2017.07.899](https://doi.org/10.1016/j.toxlet.2017.07.899)
- Plichta, J.K., Radek, K.A., 2012. Sugar-Coating Wound Repair: A Review of FGF2-10 and Dermatan Sulfate in Wound Healing and Their Potential Application in Burn Wounds. *J. Burn Care Res.* 33, 299–310.
<https://doi.org/10.1097/BCR.0b013e318240540a>
- Powers, C.J., McLeskey, S.W., Wellstein, A., 2000. Fibroblast growth factors, their receptors and signaling. *Endocr. Relat. Cancer* 7, 165–197.
<https://doi.org/10.1677/erc.0.0070165>
- Raposio, E., Bertozzi, N., Bonomini, S., Bernuzzi, G., Formentini, A., Grignaffini, E., Pio Grieco, M., 2016. Adipose-derived Stem Cells Added to Platelet-rich Plasma for Chronic Skin Ulcer Therapy. *Wounds a Compend. Clin. Res. Pract.* 28, 126–131.
- Rigotti, G., Marchi, A., Sbarbati, A., 2009. Adipose-derived mesenchymal stem cells: past, present, and future. *Aesthetic Plast. Surg.* <https://doi.org/10.1007/s00266-009-9339-7>
- Rodrigues, M., Kosaric, N., Bonham, C.A., Gurtner, G.C., 2019. Wound Healing: A Cellular Perspective. *Physiol. Rev.* 99, 665–706.
<https://doi.org/10.1152/physrev.00067.2017>
- Rohovsky, S., D'Amore, P.A., 1997. Growth Factors and Angiogenesis in Wound Healing, in: Thomas R. Ziegler, Pierce, G.F., Herndon, D.N. (Eds.), *Growth Factors and Wound Healing*. Springer New York, New York, NY, pp. 8–26.
https://doi.org/10.1007/978-1-4612-1876-0_2
- Rose, L.F., Chan, R.K., 2016. The Burn Wound Microenvironment. *Adv. Wound Care*

- 5, 106–118. <https://doi.org/10.1089/wound.2014.0536>
- Rosenstrauch, D., Poglajen, G., Zidar, N., Gregoric, I.D., 2005. Stem celltherapy for ischemic heart failure. *Texas Hear. Inst. J.* 32, 339–347.
- Rumalla, V.K., Borah, G.L., 2001. Cytokines, growth factors, and plastic surgery. *Plast. Reconstr. Surg.* 108, 719–733. <https://doi.org/10.1097/00006534-200109010-00019>
- Said, A., Wahid, F., Bashir, K., Rasheed, H.M., Khan, T., Hussain, Z., Siraj, S., 2019. Sauromatum guttatum extract promotes wound healing and tissue regeneration in a burn mouse model via up-regulation of growth factors. *Pharm. Biol.* 57, 736–743. <https://doi.org/10.1080/13880209.2019.1676266>
- Schöler, H.R., 2016. The Potential of Stem Cells: An Inventory, in: Schipanski, D., Knoepffle, N., S L Sorgner (Eds.), *Humanbiotechnology as Social Challenge An Interdisciplinary Introduction to Bioethics*. Taylor & Francis, London, pp. 1–28.
- Sethi, A., Kaur, T., Malhotra, S.K., Gambhir, M.L., 2016. Moisturizers: The Slippery Road. *Indian J. Dermatol.* 61, 279–287. <https://doi.org/10.4103/0019-5154.182427>
- Seyhan, N., Alhan, D., Ural, A.U., Gunal, A., Avunduk, M.C., Savaci, N., 2015. The Effect of Combined Use of Platelet-Rich Plasma and Adipose-Derived Stem Cells on Fat Graft Survival. *Ann. Plast. Surg.* 74, 615–620. <https://doi.org/10.1097/SAP.0000000000000480>
- Shpichka, A., Butnaru, D., Bezrukov, E.A., Sukhanov, R.B., Atala, A., Burdukovskii, V., Zhang, Y., Timashev, P., 2019. Skin tissue regeneration for burn injury. *Stem Cell Res. Ther.* 10, 1–16. <https://doi.org/10.1186/s13287-019-1203-3>
- Shukla, S.K., Sharma, A.K., Bharti, R., Kulshrestha, V., Kalonia, A., Shaw, P., 2020. Can miRNAs serve as potential markers in thermal burn injury: An in silico approach. *J. Burn Care Res.* 41, 57–64. <https://doi.org/10.1093/jbcr/irz183>
- Singh, V.K., Saini, A., Kalsan, M., Kumar, N., Chandra, R., 2016. Describing the Stem Cell Potency: The Various Methods of Functional Assessment and In silico Diagnostics. *Front. cell Dev. Biol.* 4, 134. <https://doi.org/10.3389/fcell.2016.00134>
- Subramaniam, N., Petrik, J.J., Vickaryous, M.K., 2018. VEGF, FGF2-2 and TGF β expression in the normal and regenerating epidermis of geckos: implications for epidermal homeostasis and wound healing in reptiles. *J. Anat.* 232, 768–782. <https://doi.org/10.1111/joa.12784>
- Sun, M., He, Y., Zhou, T., Zhang, P., Gao, J., Lu, F., 2017. Adipose Extracellular Matrix/Stromal Vascular Fraction Gel Secretes Angiogenic Factors and Enhances Skin Wound Healing in a Murine Model. *Biomed Res. Int.* 2017, 1–11. <https://doi.org/10.1155/2017/3105780>
- Tajima, S., Tobita, M., Orbay, H., Hyakusoku, H., Mizuno, H., 2014. Direct and

- Indirect Effects of a Combination of Adipose-Derived Stem Cells and Platelet-Rich Plasma on Bone Regeneration. *Tissue Eng. Part A* 21. <https://doi.org/10.1089/ten.TEA.2014.0336>
- Tan, S.T., Dosan, R., 2019. Lessons From Epithelialization: The Reason Behind Moist Wound Environment. *Open Dermatol. J.* 13, 34–40. <https://doi.org/10.2174/1874372201913010034>
- Tarnuzzer, R.W., Macauley, S.P., Mast, B.A., Gibson, J.S., Stacey, M.C., Trengrove, N., Moldawer, L.L., Burslem, F., Schultz, G.S., 1997. Epidermal Growth Factor in Wound Healing: A Model for the Molecular Pathogenesis of Chronic Wounds., in: Ziegler, T.R., Pierce, G.F., Herndon, D.N. (Eds.), *Growth Factors and Wound Healing*. Springer New York, New York, NY, pp. 206–228.
- Tavares Pereira, D. dos S., Lima-Ribeiro, M.H.M., de Pontes-Filho, N.T., Carneiro-Leão, A.M. dos A., Correia, M.T. dos S., 2012. Development of Animal Model for Studying Deep Second-Degree Thermal Burns. *J. Biomed. Biotechnol.* 2012, 460841. <https://doi.org/10.1155/2012/460841>
- Thisse, B., Thisse, C., 2005. Functions and regulations of fibroblast growth factor signaling during embryonic development. *Dev. Biol.* 287, 390–402. <https://doi.org/10.1016/j.ydbio.2005.09.011>
- Tohidnezhad, M., Varoga, D., Wruck, C.J., Brandenburg, L.O., Seekamp, A., Shakibaei, M., Sönmez, T.T., Pufe, T., Lippross, S., 2011. Platelet-released growth factors can accelerate tenocyte proliferation and activate the anti-oxidant response element. *Histochem. Cell Biol.* 135, 453–460. <https://doi.org/10.1007/s00418-011-0808-0>
- Toussaint, J., Singer, A.J., 2014. The evaluation and management of thermal injuries: 2014 update. *Clin. Exp. Emerg. Med.* 1, 8–18. <https://doi.org/10.15441/ceem.14.029>
- Tsien, L., 2006. Stem Cell Basics. *Postgrad. Obstet. Gynecol.* 26, 1–6. <https://doi.org/10.1097/00256406-200612310-00001>
- Turner, N., Grose, R., 2010. Fibroblast growth factor signalling: from development to cancer. *Nat. Rev. Cancer* 10, 116–129. <https://doi.org/10.1038/nrc2780>
- Vaghardoost, R., Mousavi Majd, S.G., Tebyanian, H., Babavalian, H., Malaei, L., Niazi, M., Javdani, A., 2018. The Healing Effect of Sesame Oil, Camphor and Honey on Second Degree Burn Wounds in Rat. *World J. Plast. Surg.* 7, 67–71.
- Van Pham, P., Bui, K.H.-T., Ngo, D.Q., Vu, N.B., Truong, N.H., Phan, N.L.-C., Le, D.M., Duong, T.D., Nguyen, T.D., Le, V.T., Phan, N.K., 2013. Activated platelet-rich plasma improves adipose-derived stem cell transplantation efficiency in injured articular cartilage. *Stem Cell Res. Ther.* 4, 91. <https://doi.org/10.1186/scrt277>

- Velnar, T., Bailey, T., Smrkolj, V., 2009. The Wound Healing Process: An Overview of the Cellular and Molecular Mechanisms. *J. Int. Med. Res.* 37, 1528–1542.
<https://doi.org/10.1177/147323000903700531>
- Venturi, S., Venturi, M., 2009. Iodine in Evolution of Salivary Glands and in Oral Health. *Nutr. Health* 20, 119–134. <https://doi.org/10.1177/026010600902000204>
- Wang, Y., Beekman, J., Hew, J., Jackson, S., Issler-Fisher, A.C., Parungao, R., Lajevardi, S.S., Li, Z., Maitz, P.K.M., 2018. Burn injury: Challenges and advances in burn wound healing, infection, pain and scarring, *Advanced Drug Delivery Reviews*. Elsevier B.V. <https://doi.org/10.1016/j.addr.2017.09.018>
- Werner, S., Grose, R., 2003. Regulation of wound healing by growth factors and cytokines. *Physiol. Rev.* 83, 835–870.
<https://doi.org/10.1152/physrev.2003.83.3.835>
- Widowati, W., Widyanto, R.M., 2013. Sel Punca sebagai Transformasi Alternatif Terapi. *Zenit* 2, 1–5.
- Witte, M.B., Barbul, A., 1997. General principles of wound healing. *Surg. Clin. North Am.* 77, 509–528. [https://doi.org/10.1016/s0039-6109\(05\)70566-1](https://doi.org/10.1016/s0039-6109(05)70566-1)
- Wobus, A.M., Boheler, K.R., 2005. Embryonic Stem Cells: Prospects for Developmental Biology and Cell Therapy. *Physiol. Rev.* 85, 635–678.
<https://doi.org/10.1152/physrev.00054.2003>
- Wong, A., Lamothe, B., Lee, A., Schlessinger, J., Lax, I., 2002. FRS2 alpha attenuates FGF2 receptor signaling by Grb2-mediated recruitment of the ubiquitin ligase Cbl. *Proc. Natl. Acad. Sci. U. S. A.* 99, 6684–6689.
<https://doi.org/10.1073/pnas.052138899>
- Yun, Y.-R., Won, J.E., Jeon, E., Lee, S., Kang, W., Jo, H., Jang, J.-H., Shin, U.S., Kim, H.-W., 2010. Fibroblast growth factors: biology, function, and application for tissue regeneration. *J. Tissue Eng.* 2010, 218142.
<https://doi.org/10.4061/2010/218142>
- Zakrzewski, W., Dobrzański, M., Szymonowicz, M., Rybak, Z., 2019. Stem cells: past, present, and future. *Stem Cell Res. Ther.* 10, 68. <https://doi.org/10.1186/s13287-019-1165-5>
- Zhang, Y., He, J., Xiao, G., Li, Q., 2011. [Effect of platelet-rich plasma on the proliferation and adipogenic differentiation of human adipose-derived stem cells in vitro]. *Nan Fang Yi Ke Da Xue Xue Bao* 31, 525–528.
- Zuk, P.A., Zhu, M., Ashjian, P., De Ugarte, D.A., Huang, J.I., Mizuno, H., Alfonso, Z.C., Fraser, J.K., Benhaim, P., Hedrick, M.H., 2002. Human adipose tissue is a source of multipotent stem cells. *Mol. Biol. Cell* 13, 4279–4295.
<https://doi.org/10.1091/mbc.e02-02-0105>

Lampiran : Dokumentasi Penelitian

Sampel tikus Wistar



Menimbang BB tikus



Proses pengambilan Donor PRP dan SVFs
(Insert: Gambar 1. Pembiusan dengan Eter fisiologis. 2. Pengambilan Donor PRP.
3 dan 4. Pengambilan Donor SVFS)



Proses preparasi Donor SVFs


(Insert: Gambar 1. Penimbangan Lemak Donor. 2. Pencacahan lemak Donor.

3. Penambahan cairan kolagenase 0.15% & Sentrifugasi. 4. Inkubasi.


5. Proses penghitungan Jumlah Stem cell 6. Hasil Stem cell)



Cara Pemodelan Luka Bakar
(Tavacres Pereira et al., 2012; Guo et al., 2017).



Perlakuan Injeksi PRP dan SVFs.
(Insert: Gambar 1. Pencukuran Bulu Wistar. 2. Injeksi Stem cell. -
3. Lima lokasi injeksi Stem cell)



Bahan dan Proses Pemeriksaan EGF.

(Insert: Gambar 1. KIT FGF2. 2. Serum Hewan coba. 3. Well Elisa Kit. 4. Inkubasi 37° selama 90 menit. 5. Penambahan Wash buffer 1x sebanyak 3x & 100 µL HRP conjugate. 6. Baca dan Ukur memakai alat microplate reader).

