

## DAFTAR PUSTAKA

- Ahn, H. S., Lee, J. B., Seo, M., Park, S. H., & Choi, B. I. (2018). Distinguishing benign from malignant thyroid nodules using thyroid ultrasonography: utility of adding superb microvascular imaging and elastography. *Radiologia Medica*, 123(4), 260–270. <https://doi.org/10.1007/s11547-017-0839-2>
- Algin, O., Algin, E., Gokalp, G., Ocakoğlu, G., Cüneyt, E., Saraydaroglu, O., & Tuncel, E. (2010). Role of duplex power Doppler ultrasound in differentiation between malignant and benign thyroid nodules. *Korean Journal of Radiology*, 11(6), 594–602. <https://doi.org/10.3348/kjr.2010.11.6.594>
- Aslan, A., Sancak, S., Aslan, M., Cimsit, N. C., Güllüoğlu, B. M., Ahiskali, R. A., Akalin, N. S., & Aribal, M. E. (2013). Ultrasonography and duplex doppler ultrasonography based indices in nodular thyroid disease. *Acta Endocrinologica*, 9(4), 575–588. <https://doi.org/10.4183/aeb.2013.575>
- Aslan, Ahmet, Sancak, S., Aslan, M., Ayaz, E., Inan, I., Ozkanli, S. S., Alimoğlu, O., & Yıkılmaz, A. (2018). Diagnostic value of duplex doppler ultrasound parameters in papillary thyroid carcinoma. *Acta Endocrinologica*, 14(1), 43–48. <https://doi.org/10.4183/aeb.2018.43>
- Azizi, G., & Malchoff, C. D. (2018). Ultrasound Elastography of Thyroid Nodules. *Thyroid and Parathyroid Ultrasound and Ultrasound-Guided FNA*, 489–516. [https://doi.org/10.1007/978-3-319-67238-0\\_16](https://doi.org/10.1007/978-3-319-67238-0_16)
- Baig, F. N., Van Lunenburg, J. T. J., Liu, S. Y. W., Yip, S. P., Law, H. K. W., & Ying, M. (2017). Computer-aided assessment of regional vascularity of thyroid nodules for prediction of malignancy. *Scientific Reports*, 7(1), 1–9. <https://doi.org/10.1038/s41598-017-14432-7>
- Bertazza, L., Mocellin, S., Marchet, A., Pilati, P., Gabrieli, J., Scalerta, R., & Nitti, D. (2009). Survivin gene levels in the peripheral blood of patients with gastric cancer independently predict survival. *Journal of Translational Medicine*, 7, 1–8. <https://doi.org/10.1186/1479-5876-7-111>
- Cantisani, V., Lodise, P., Grazhdani, H., Mancuso, E., Maggini, E., Di Rocco, G., D'Ambrosio, F., Calliada, F., Redler, A., Ricci, P., & Catalano, C. (2014). Ultrasound elastography in the evaluation of thyroid pathology. Current status. *European Journal of Radiology*, 83(3), 420–428. <https://doi.org/10.1016/j.ejrad.2013.05.008>
- Chen, S., Niu, C., Peng, Q., & Tang, K. (2021). Sonographic Characteristics of Papillary Thyroid Carcinoma With Coexistent

- Hashimoto's Thyroiditis in the Preoperative Prediction of Central Lymph Node Metastasis. *Frontiers in Endocrinology*, 12(March), 1–9. <https://doi.org/10.3389/fendo.2021.556851>
- Choi, S. H., Kim, E. K., Kim, S. J., & Kwak, J. Y. (2014). Thyroid ultrasonography: Pitfalls and techniques. *Korean Journal of Radiology*, 15(2), 267–276. <https://doi.org/10.3348/kjr.2014.15.2.267>
- Cosgrove, D., Barr, R., Bojunga, J., Cantisani, V., Chammas, M. C., Dighe, M., Vinayak, S., Xu, J. M., & Dietrich, C. F. (2017). WFUMB Guidelines and Recommendations on the Clinical Use of Ultrasound Elastography: Part 4. Thyroid. *Ultrasound in Medicine and Biology*, 43(1), 4–26. <https://doi.org/10.1016/j.ultrasmedbio.2016.06.022>
- Di, Z., Li, Z., Tian, J., Wang, D., Liu, L., & Liu, C. (2019). The Value of Elasticity Contrast Index in the Differential Diagnosis of Thyroid Solid Nodules. *Ultrasound Quarterly*, 35(3), 259–263. <https://doi.org/10.1097/RUQ.0000000000000457>
- Dietrich, C. F., Barr, R. G., Farrokh, A., Dighe, M., Hocke, M., Jenssen, C., Dong, Y., Saftoiu, A., & Havre, R. F. (2016). 1 Introduction to Elastography. *Elastography*. <https://doi.org/10.1055/b-0036-140635>
- Durante, C., Grani, G., Lamartina, L., Filetti, S., Mandel, S. J., & Cooper, D. S. (2018). The diagnosis and management of thyroid nodules a review. In *JAMA - Journal of the American Medical Association* (Vol. 319, Issue 9). <https://doi.org/10.1001/jama.2018.0898>
- Franklin NTesser, William D Middleton, E. G. G. (2018). thyroid imaging Reporting and data system (TI-RADS): A User's Guide 1 HOW I DO IT: Thyroid Imaging Reporting and Data System Tessler et al. *Radiology*, 287, 29–36. <https://doi.org/10.1148/radiol.2017171240>
- Ferreira de Souza, M., Kuasne, H., de Camargo Barros-Filho, M., Lizotti Cilião, H., Albuquerque Marchi, F., Emilio Fuganti, P., Rossi Paschoal, A., Regina Rogatto, S., & Mara de Syllis Có Ius, I. (2017). Circulating mRNAs and miRNAs as candidate markers for the diagnosis and prognosis of prostate cancer. *PLoS ONE*, 12(9), e0184094.
- Gao, L., Xi, X., Jiang, Y., Yang, X., Wang, Y., Zhu, S., Lai, X., Zhang, X., Zhao, R., & Zhang, B. (2019). Comparison among TIRADS (ACR TI-RADS and KWAK- TI-RADS) and 2015 ATA Guidelines in the diagnostic efficiency of thyroid nodules. *Endocrine*, 64(1), 90–96. <https://doi.org/10.1007/s12020-019-01843-x>
- Gede, I. W., Eka, A., Epid, M., Luh, N., Suariyani, P., Septarini, N. W. (2016). UJI DIAGNOSTIK DAN SKRINING.
- Grant, E. G., Tessler, F. N., Hoang, J. K., Langer, J. E., Beland, M. D., Berland, L. L., Cronan, J. J., Desser, T. S., Frates, M. C., Hamper, U. M., Middleton, W. D., Reading, C. C., Scoult, L. M., Stavros, A. T., &

- Teefey, S. A. (2015). *Thyroid Ultrasound Reporting Lexicon: White Paper of the ACR Thyroid Imaging, Reporting and Data System (TIRADS) Committee*. <https://doi.org/10.1016/j.jacr.2015.07.011>
- Habib, L. A. M., Abd Rabou, A. M., Geneidi, E. A. S., & Sultan, Y. M. (2016). Role of ultrasound elastography in assessment of indeterminate thyroid nodules. *The Egyptian Journal of Radiology and Nuclear Medicine*, 47(1), 141–147. <https://doi.org/10.1016/j.ejrm.2015.11.002>
- Hairu, L., Yulan, P., Yan, W., Hong, A., Xiaodong, Z., Lichun, Y., Kun, Y., Ying, X., Lisha, L., Baoming, L., Qiang, Y., Shuzhen, C., Shuangquan, J., Xin, F., Buyun, M., Yi, L., Xixi, Z., Xue, G., Haitao, C., ... Jiawei, T. (2020). Elastography for the diagnosis of high-suspicion thyroid nodules based on the 2015 American Thyroid Association guidelines: A multicenter study. *BMC Endocrine Disorders*, 20(1), 1–10. <https://doi.org/10.1186/s12902-020-0520-y>
- Hatta M. 2018. Pelatihan Aplikasi Teknik Biologi Molekuler dan Imunologi dalam Penelitian bidang Kesehatan. Laboratorium Biologi Molekuler dan Imunologi, Fakultas Kedokteran, Universitas Hasanuddin, Makassar
- Hee Shin, J., Hwan Baek, J., Chung, J., Ju Ha, E., Kim, J., Hen Lee, Y., Kyung Lim, H., Moon, W.-J., Gyu Na, D., Seon Park, J., Jung Choi, Y., Yeon Hahn, S., Jeong Jeon, S., Lyung Jung, S., Wook Kim, D., Kim, E.-K., Young Kwak, J., Yoon Lee, C., Joong Lee, H., ... Young Sung, J. (2016). Ultrasonography Diagnosis and Imaging-Based Management of Thyroid Nodules: Revised Korean Society of Thyroid Radiology Consensus Statement and Recommendations. <https://doi.org/10.3348/kjr.2016.17.3.370>
- Horvath, E., Majlis, S., Rossi, R., Franco, C., Niedmann, J. P., Castro, A., Dominguez, M., & Board, T. (2009). An Ultrasonogram Reporting System for Thyroid Nodules Stratifying Cancer Risk for Clinical Management. *J Clin Endocrinol Metab*, 90, 1748–1751. <https://doi.org/10.1210/jc.2008-1724>
- Jankovic, B., Le, K. T., & Hershman, J. M. (2013). Hashimoto's thyroiditis and papillary thyroid carcinoma: Is there a correlation? *Journal of Clinical Endocrinology and Metabolism*, 98(2), 474–482. <https://doi.org/10.1210/jc.2012-2978>
- Jayaraj, R., Muhyambere, B., & Suja, S. (2018). *Cytokeratin 19 (CK19) as a marker for Epithelial Differentiation and Malignant Transformation: Its Clinical relevance in Diagnosis, Prognosis and Treatment response monitoring*.
- Kalantari, S. (2018). The diagnostic value of color doppler ultrasonography in predicting thyroid nodules malignancy. *International Tinnitus Journal*, 22(1), 35–39. <https://doi.org/10.5935/0946-5448.20180006>
- Khayyata, S., Barroeta, J. E., LiVolsi, V. A., & Baloch, Z. W. (2008).

- Papillary hyperplastic nodule: Pitfall in the cytopathologic diagnosis of papillary thyroid carcinoma. *Endocrine Practice*, 14(7), 863–868. <https://doi.org/10.4158/EP.14.7.863>
- Kwak, J. Y., & Kim, E. K. (2014). Ultrasound elastography for thyroid nodules: Recent advances. *Ultrasonography*, 33(2), 75–82. <https://doi.org/10.14366/usg.13025>
- Liao, L., Chen, H., Hsu, W., & Chen, Y. (2019). Comparison of Strain Elastography , Shear Wave Elastography , and Conventional Ultrasound in Diagnosing Thyroid Nodules. 26–32. <https://doi.org/10.4103/JMU.JMU>
- Levine, R. A. (2018). *Doppler ultrasound*. 43–70. <https://doi.org/10.3109/9781439814697-51>
- Mahmoudian-Sani, M.-R., Alghasi, A., Saeedi-Boroujeni, A., Jalali, A., Jamshidi, M., & Khodadadi, A. (2019). *Survivin as a diagnostic and therapeutic marker for thyroid cancer*. <https://doi.org/10.1016/j.prp.2019.01.025>
- Marcy, P. Y., Russ, G., Saba, L., Sanglier, J., Ghanassia, E., Sharara, H., Thariat, J., Morvan, J. B., & Bizeau, A. (2022). Opinion: leading position of ultrasound in decision algorithm for small papillary thyroid carcinoma. *Insights into Imaging*, 13(1), 5–11. <https://doi.org/10.1186/s13244-022-01240-5>
- Mulyadi, R., Hatta, M., Islam, A. A., Murtala, B., Tammase, J., Firdaus, M., Susanto, E., & Prihartono, J. (2021). Intratumoral and Peritumoral Apparent Diffusion Coefficient and MGMT mRNA Expression in Different Meningioma Histopathological Grade. *Indonesian Biomedical Journal*, 13(1), 97–105. <https://doi.org/10.18585/inabj.v13i1.1338>
- Sastroasmoro S, Ismail S. (2014) Dasar-dasar Metodologi Penelitian Klinis. edisi ke-5. Sagung Seto.
- Okasha, H. H., Mansor, M., Sheriba, N., Assem, M., Ashoush, O. A., Rakha, M., Abdelfatah, D., El-sawy, S. S., Elshenoufy, M., Mohsen, A. A., Sedrak, H. K., & Abdellatif, A. A. (2020). and TIRADS score in predicting malignant thyroid nodule. <https://doi.org/10.20945/2359-3997000000283>
- Qin, L., Liu, Q., Sun, L., & Wang, H. (2022). Accuracy of ultrasound in distinguishing pathology of malignant thyroid diseases: A protocol for systematic review and meta-analysis. *Medicine (United States)*, 101(2), E28556. <https://doi.org/10.1097/MD.00000000000028556>
- Rosario, P. W., da Silva, A. L., Borges, M. A. R., & Calsolari, M. R. (2015). Is Doppler ultrasound of additional value to gray-scale ultrasound in differentiating malignant and benign thyroid nodules? *Archives of Endocrinology and Metabolism*, 59(1), 79–83. <https://doi.org/10.1590/2359-3997000000014>

- Selemetjev, S. A., Savin, S. B., Paunovic, I. R., Tatic, S. B., & Cvejic, D. (2015). Änderungen im Expressionsmuster von apoptotischen Molekülen (Galectin-3, Bcl-2, Bax, Survivin) im Rahmen der Progression der Malignität von Schilddrüsentumoren und deren klinische Bedeutung. *Wiener Klinische Wochenschrift*, 127(9–10), 337–344. <https://doi.org/10.1007/s00508-014-0674-6>
- Shin, J. H. (2017). Ultrasonographic imaging of papillary thyroid carcinoma variants. *Ultrasonography*, 36(2), 103–110. <https://doi.org/10.14366/usg.16048>
- Tessler, F. N., Middleton, W. D., Grant, E. G., Hoang, J. K., Berland, L. L., Teeffey, S. A., Cronan, J. J., Beland, M. D., Desser, T. S., Frates, M. C., Hammers, L. W., Hamper, U. M., Langer, J. E., Reading, C. C., Scoutt, L. M., & Stavros, A. T. (2017). ACR Thyroid Imaging, Reporting and Data System (TI-RADS): White Paper of the ACR TI-RADS Committee. *Journal of the American College of Radiology*, 14(5), 587–595. <https://doi.org/10.1016/j.jacr.2017.01.046>
- Thomas, S., & Shah, G. (2005). Calcitonin induces apoptosis resistance in prostate cancer cell lines against cytotoxic drugs via the Akt/survivin pathway. *Cancer Biology and Therapy*, 4(11), 1226–1233. <https://doi.org/10.4161/cbt.4.11.2093>
- Yu, J., Wang, Z., Zhang, H., Wang, Y., & Li, D. Q. (2021). Survivin-positive circulating tumor cells as a marker for metastasis of hepatocellular carcinoma. *World Journal of Gastroenterology*, 27(43), 7546–7562. <https://doi.org/10.3748/wjg.v27.i43.7546>
- Xie, C., Cox, P., Taylor, N., & LaPorte, S. (2016). Ultrasonography of thyroid nodules: a pictorial review. *Insights into Imaging*, 7(1), 77–86. <https://doi.org/10.1007/s13244-015-0446-5>
- Yuen, H. Y., Wong, K. T., & Ahuja, A. T. (2016). Sonography of diffuse thyroid disease. *Australasian Journal of Ultrasound in Medicine*, 19(1), 13–29. <https://doi.org/10.1002/ajum.12001>

## LAMPIRAN




Lampiran 1. Primer RTPCR Gen Human Survivin / Housekeeping Gen GAPDH



Lampiran 2. Primer RTPCR Gen Human Cytokeratin-19 / Housekeeping Gen GAPDH



Lampiran 3. PCR Master Mix SYBR Green



Lampiran 4. Pembuatan PCR Mix



Lampiran 5. Pengukuran template pada Nanodrop



Lampiran 6. Memasukkan template ke dalam PCR Mix



Lampiran 7. Mesin PCR



Lampiran 8. Analisa hasil PCR

## LAMPIRAN 9. ANALISA STATISTIK

### KARAKTERISTIK SUBJEK

**JENIS KELAMIN**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Laki-laki	4	9.5	9.5	9.5
	Perempuan	38	90.5	90.5	100.0
	Total	42	100.0	100.0	

**UMUR**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	≤40 Tahun	27	64.3	64.3	64.3
	41-50 Tahun	5	11.9	11.9	76.2
	51-60 Tahun	7	16.7	16.7	92.9
	>60 Tahun	3	7.1	7.1	100.0
	Total	42	100.0	100.0	

**TIRADS**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	2.0	5	11.9	11.9	11.9
	3.0	17	40.5	40.5	52.4
	4.0	15	35.7	35.7	88.1
	5.0	5	11.9	11.9	100.0
	Total	42	100.0	100.0	

**HISTOLOGI**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	GANAS	20	47.6	47.6	47.6
	JINAK	22	52.4	52.4	100.0
	Total	42	100.0	100.0	

**PATOLOGI ANATOMI**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	ADENOMATOUS GOITER	16	38.1	38.1	38.1
	HASHIMOTO THYROIDITIS	2	4.8	4.8	42.9
	PAPILLARY THYROID				
	CARCINOMA	20	47.6	47.6	90.5
	SOLITARY PAPILLARY				
	HYPERPLASTIC	1	2.4	2.4	92.9
	WELL DIFFERENTIATED				
	NEOPLASMA	3	7.1	7.1	100.0
	Total	42	100.0	100.0	

**LOKASI NODUL**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	BILATERAL	4	9.5	9.5	9.5
	KANAN	24	57.1	57.1	66.7
	KIRI	14	33.3	33.3	100.0
	Total	42	100.0	100.0	

**UKURAN NODUL**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	≤1 cm	7	16.7	16.7	16.7
	1.1 - 2 cm	19	45.2	45.2	61.9
	2.1 - 4 cm	14	33.3	33.3	95.2
	>4 cm	2	4.8	4.8	100.0
	Total	42	100.0	100.0	

**UKURAN NODUL**

N	Valid	42
	Missing	0
Mean		2.2619
Std. Error of Mean		.12316

## KOMBINASI SERIAL TIRADS +RI

. diagt patologianatomi serialtiradsri

PATOLOGI ANATOMI	serialtiradsri		Total
	Pos.	Neg.	
Abnormal	13	7	20
Normal	1	21	22
Total	14	28	42

True abnormal diagnosis defined as patologianatomi = 1

[95% Confidence Interval]				
Prevalence	Pr(A)	48%	32%	63.6%
Sensitivity	Pr(+ A)	65%	40.8%	84.6%
Specificity	Pr(- N)	95.5%	77.2%	99.9%
ROC area	(Sens. + Spec.)/2	.802	.686	.918
Likelihood ratio (+)	Pr(+ A)/Pr(+ N)	14.3	2.05	99.7
Likelihood ratio (-)	Pr(- A)/Pr(- N)	.367	.2	.671
Odds ratio	LR(+) / LR(-)	39	5.24	.
Positive predictive value	Pr(A +)	92.9%	66.1%	99.8%
Negative predictive value	Pr(N -)	75%	55.1%	89.3%

## KOMBINASI SERIAL TIRADS +PI

. diagt patologianatomi serialtiradspi

PATOLOGI ANATOMI	serialtiradspi		Total
	Pos.	Neg.	
Abnormal	12	8	20
Normal	1	21	22
Total	13	29	42

True abnormal diagnosis defined as patologianatomi = 1

[95% Confidence Interval]				
Prevalence	Pr(A)	48%	32%	63.6%
Sensitivity	Pr(+ A)	60%	36.1%	80.9%
Specificity	Pr(- N)	95.5%	77.2%	99.9%
ROC area	(Sens. + Spec.)/2	.777	.658	.896
Likelihood ratio (+)	Pr(+ A)/Pr(+ N)	13.2	1.88	92.6
Likelihood ratio (-)	Pr(- A)/Pr(- N)	.419	.243	.722
Odds ratio	LR(+) / LR(-)	31.5	4.29	.
Positive predictive value	Pr(A +)	92.3%	64%	99.8%
Negative predictive value	Pr(N -)	72.4%	52.8%	87.3%

## KOMBINASI SERIAL TIRADS + ELASTOGRAPHY

. diagt patologianatomi serialtiradselasto

PATOLOGI ANATOMI	serialtiradselasto		Total
	Pos.	Neg.	
Abnormal	13	7	20
Normal	1	21	22
Total	14	28	42

True abnormal diagnosis defined as patologianatomi = 1

[95% Confidence Interval]

Prevalence	Pr(A)	48%	32%	63.6%
Sensitivity	Pr(+ A)	65%	40.8%	84.6%
Specificity	Pr(- N)	95.5%	77.2%	99.9%
ROC area	(Sens. + Spec.)/2	.802	.686	.918
Likelihood ratio (+)	Pr(+ A)/Pr(+ N)	14.3	2.05	99.7
Likelihood ratio (-)	Pr(- A)/Pr(- N)	.367	.2	.671
Odds ratio	LR(+) / LR(-)	39	5.24	.
Positive predictive value	Pr(A +)	92.9%	66.1%	99.8%
Negative predictive value	Pr(N -)	75%	55.1%	89.3%

## KOMBINASI SERIAL TIRADS + SURVIVIN

. diagt patologianatomi serialTIRADSSURVIVIN

PATOLOGI ANATOMI	serialTIRADSSURVIVIN		Total
	Pos.	Neg.	
Abnormal	17	3	20
Normal	0	22	22
Total	17	25	42

True abnormal diagnosis defined as patologianatomi = 1

[95% Confidence Interval]

Prevalence	Pr(A)	48%	32%	63.6%
Sensitivity	Pr(+ A)	85%	62.1%	96.8%
Specificity	Pr(- N)	100%	84.6%	100%
ROC area	(Sens. + Spec.)/2	.925	.845	1
Likelihood ratio (+)	Pr(+ A)/Pr(+ N)	.	.	.
Likelihood ratio (-)	Pr(- A)/Pr(- N)	.15	.0528	.426
Odds ratio	LR(+) / LR(-)	.	24.4	.
Positive predictive value	Pr(A +)	100%	80.5%	100%
Negative predictive value	Pr(N -)	88%	68.8%	97.5%

## KOMBINASI SERIAL TIRADS + CYTOKERATIN-19

. diagt patologianatomi serialtiradscytokeratin19

PATOLOGI ANATOMI	serialtiradscytokerat in19		
	Pos.	Neg.	Total
Abnormal	16	4	20
Normal	0	22	22
Total	16	26	42

True abnormal diagnosis defined as patologianatomi = 1

[95% Confidence Interval]

Prevalence	Pr (A)	48%	32%	63.6%
Sensitivity	Pr (+ A)	80%	56.3%	94.3%
Specificity	Pr (- N)	100%	84.6%	100%
ROC area	(Sens. + Spec.) / 2	.9	.81	.99
Likelihood ratio (+)	Pr (+ A) / Pr (+ N)	.	.	.
Likelihood ratio (-)	Pr (- A) / Pr (- N)	.2	.0832	.481
Odds ratio	LR (+) / LR (-)	.	18.3	.
Positive predictive value	Pr (A +)	100%	79.4%	100%
Negative predictive value	Pr (N -)	84.6%	65.1%	95.6%

## KOMBINASI PARAREL TIRADS + RI

. diagt patologianatomi parareltiradsri

PATOLOGI ANATOMI	parareltiradsri		
	Pos.	Neg.	Total
Abnormal	19	1	20
Normal	5	17	22
Total	24	18	42

True abnormal diagnosis defined as patologianatomi = 1

[95% Confidence Interval]

Prevalence	Pr (A)	48%	32%	63.6%
Sensitivity	Pr (+ A)	95%	75.1%	99.9%
Specificity	Pr (- N)	77.3%	54.6%	92.2%
ROC area	(Sens. + Spec.) / 2	.861	.759	.964
Likelihood ratio (+)	Pr (+ A) / Pr (+ N)	4.18	1.92	9.09
Likelihood ratio (-)	Pr (- A) / Pr (- N)	.0647	.0095	.443
Odds ratio	LR (+) / LR (-)	64.6	8.24	.
Positive predictive value	Pr (A +)	79.2%	57.8%	92.9%
Negative predictive value	Pr (N -)	94.4%	72.7%	99.9%

## KOMBINASI PARAREL TIRADS + PI

. diagt patologianatomi parareltiradspi

PATOLOGI ANATOMI	parareltiradspi		Total
	Pos.	Neg.	
Abnormal	19	1	20
Normal	11	11	22
Total	30	12	42

True abnormal diagnosis defined as patologianatomi = 1

[95% Confidence Interval]

Prevalence	Pr(A)	48%	32%	63.6%
Sensitivity	Pr(+ A)	95%	75.1%	99.9%
Specificity	Pr(- N)	50%	28.2%	71.8%
ROC area	(Sens. + Spec.)/2	.725	.607	.843
Likelihood ratio (+)	Pr(+ A)/Pr(+ N)	1.9	1.24	2.92
Likelihood ratio (-)	Pr(- A)/Pr(- N)	.1	.0141	.707
Odds ratio	LR(+) / LR(-)	19	2.66	.
Positive predictive value	Pr(A +)	63.3%	43.9%	80.1%
Negative predictive value	Pr(N -)	91.7%	61.5%	99.8%

## KOMBINASI PARAREL TIRADS + ELASTOGRAPHY

. diagt patologianatomi parareltiradselasto

PATOLOGI ANATOMI	parareltiradselasto		Total
	Pos.	Neg.	
Abnormal	19	1	20
Normal	6	16	22
Total	25	17	42

True abnormal diagnosis defined as patologianatomi = 1

[95% Confidence Interval]

Prevalence	Pr(A)	48%	32%	63.6%
Sensitivity	Pr(+ A)	95%	75.1%	99.9%
Specificity	Pr(- N)	72.7%	49.8%	89.3%
ROC area	(Sens. + Spec.)/2	.839	.732	.946
Likelihood ratio (+)	Pr(+ A)/Pr(+ N)	3.48	1.75	6.94
Likelihood ratio (-)	Pr(- A)/Pr(- N)	.0688	.01	.472
Odds ratio	LR(+) / LR(-)	50.7	6.68	.
Positive predictive value	Pr(A +)	76%	54.9%	90.6%
Negative predictive value	Pr(N -)	94.1%	71.3%	99.9%

## KOMBINASI PARAREL TIRADS + SURVIVIN

. diagt patologianatomi parareltIRADSSURVIVIN

PATOLOGI ANATOMI	parareltIRADSSURVIVIN			Total
	Pos.	Neg.		
Abnormal	19	1		20
Normal	4	18		22
Total	23	19		42

True abnormal diagnosis defined as patologianatomi = 1

[95% Confidence Interval]

Prevalence	Pr(A)	48%	32%	63.6%
Sensitivity	Pr(+ A)	95%	75.1%	99.9%
Specificity	Pr(- N)	81.8%	59.7%	94.8%
ROC area	(Sens. + Spec.)/2	.884	.788	.98
Likelihood ratio (+)	Pr(+ A)/Pr(+ N)	5.22	2.14	12.8
Likelihood ratio (-)	Pr(- A)/Pr(- N)	.0611	.009	.417
Odds ratio	LR(+)/LR(-)	85.5	10.4	.
Positive predictive value	Pr(A +)	82.6%	61.2%	95%
Negative predictive value	Pr(N -)	94.7%	74%	99.9%

## KOMBINASI PARAREL TIRADS + CYTOKERATIN-19

. diagt patologianatomi parareltiradscytokeratin19

PATOLOGI ANATOMI	parareltiradscytoker atin19			Total
	Pos.	Neg.		
Abnormal	19	1		20
Normal	4	18		22
Total	23	19		42

True abnormal diagnosis defined as patologianatomi = 1

[95% Confidence Interval]

Prevalence	Pr(A)	48%	32%	63.6%
Sensitivity	Pr(+ A)	95%	75.1%	99.9%
Specificity	Pr(- N)	81.8%	59.7%	94.8%
ROC area	(Sens. + Spec.)/2	.884	.788	.98
Likelihood ratio (+)	Pr(+ A)/Pr(+ N)	5.22	2.14	12.8
Likelihood ratio (-)	Pr(- A)/Pr(- N)	.0611	.009	.417
Odds ratio	LR(+)/LR(-)	85.5	10.4	.
Positive predictive value	Pr(A +)	82.6%	61.2%	95%
Negative predictive value	Pr(N -)	94.7%	74%	99.9%