

DAFTAR PUSTAKA

1. Mosayyebi A, Manes C, Carugo D, Somani BK. Advances in Ureteral Stent Design and Materials. *Curr Urol Rep.* 2018;19(5).
2. Aggarwal SP, Priyadarshi S, Tomar V, Yadav SS, Gangkak G, Vyas N, et al. A randomized controlled trial to compare the safety and efficacy of Tadalafil and Tamsulosin in relieving double J stent related symptoms. *Adv Urol.* 2015;2015.
3. Lu J, Lu Y, Xun Y, Chen F, Wang S, Cao S. Impact of Endourological procedures with or without double-J stent on sexual function: A systematic review and meta-analysis. *BMC Urol.* 2020;20(1):1–8.
4. Vogt B, Chokri I. Histological Inflammation in Human Ureter either Healthy or Fitted with Double-Pigtail Stent or a Thin 0.3 F Suture Thread: A Preliminary Study. *Adv Urol.* 2020;2020.
5. Lange D, Bidnur S, Hoag N, Chew BH. Ureteral stent-associated complications—where we are and where we are going. *Nat Rev Urol [Internet].* 2015;12(1):17–25. Available from: <http://dx.doi.org/10.1038/nrurol.2014.340>
6. Furuta A. Comparison of inflammatory urine markers in patients with interstitial cystitis and overactive bladder. 2018;3–8.
7. Tanidir Y, Mangir N, Sahan A, Sulukaya M. Turkish version of the Ureteral Stent Symptoms Questionnaire: linguistic and psychometric validation. *World J Urol.* 2017;35(7):1149–54.
8. Petes C, Mariani MK, Yang Y, Grandvaux N, Gee K. Interleukin (IL)-6 inhibits IL-27- and IL-30-mediated inflammatory responses in human monocytes. *Front Immunol.* 2018;9(FEB):1–16.
9. Al-eisa A. Serum and Urine Interleukin-6 and Interleukin-8 Levels Do Not Differentiate Acute Pyelonephritis from Lower Urinary Tract Infections in Children. 2020;
10. Lotz M, Villiger P, Li THUG, Koziol J, W BLZ. Interleukin-6 and interstitial cystitis. 1994;152(September):869–73.
11. Tae BS, Cho S, Jeon BJ, Choi H, Park JY, Cho SY, et al. Does mirabegron relieve ureteric stent-related discomfort? A prospective, randomized, multicentre study. *BJU Int.* 2018;122(5):866–72.
12. Otsuki H, Yamasaki T, Hori S, Aoki H, Kosaka T, Uehara S, et al. The efficacy of mirabegron for the relief of ureteral stent-related symptoms. *Acta Med Okayama.* 2020;74(2):145–50.
13. Lena M Di, Tolls V, Kelly K, Nickel JC. Mirabegron as adjuvant treatment for patients with interstitial cystitis / bladder pain syndrome. 2018;12(3).
14. Navanimitkul N, Lojanapiwat B. Efficacy of tamsulosin 0.4 mg/day in relieving double-J stent-related symptoms: A randomized controlled study. *J Int Med Res.* 2010;38(4):1436–41.
15. Beysens M, Tailly TO. Ureteral stents in urolithiasis. *Asian J Urol [Internet].* 2018;5(4):274–86. Available from: <https://doi.org/10.1016/j.ajur.2018.07.002>
16. Gibson W, MacDiarmid S, Huang M, Siddiqui E, Stölzel M, Choudhury N, et

- al. Treating Overactive Bladder in Older Patients with a Combination of Mirabegron and Solifenacin: A Prespecified Analysis from the BESIDE Study. *Eur Urol Focus*. 2017;3(6):629–38.
17. Kelleher C, Hakimi Z, Zur R, Siddiqui E, Maman K, Aballéa S, et al. Efficacy and Tolerability of Mirabegron Compared with Antimuscarinic Monotherapy or Combination Therapies for Overactive Bladder: A Systematic Review and Network Meta-analysis [Figure presented]. *Eur Urol*. 2018;74(3):324–33.
 18. Camtosun A, Bicer S. The impact of Double J stent on the quality of sexual life and job performance. *Clin Exp Obstet Gynecol*. 2020;47(2):199–201.
 19. Zumstein V, Betschart P, Albrich WC, Buhmann MT, Ren Q, Schmid H-P, et al. Biofilm formation on ureteral stents - Incidence, clinical impact, and prevention. *Swiss Med Wkly*. 2017 Feb;
 20. Sighinolfi MC, Sighinolfi GP, Galli E, Micali S, Ferrari N, Mofferdin A, et al. Chemical and Mineralogical Analysis of Ureteral Stent Encrustation and Associated Risk Factors. *Urology*. 2015;86(4):703–6.
 21. Fischer KM, Louie M, Mucksavage P. Ureteral Stent Discomfort and Its Management. *Curr Urol Rep*. 2018;19(8):5–8.
 22. Gonzalez EJ, Arms L, Vizzard MA. The Role (s) of Cytokines / Chemokines in Urinary Bladder Inflammation and Dysfunction. 2014;2014.
 23. Rodhe N, Löfgren S, Strindhall J, Matussek A. Cytokines in urine in elderly subjects with acute cystitis and asymptomatic bacteriuria. 2009;3432.
 24. Akdis M, Burgler S, Crameri R, Eiwegger T, Fujita H, Gomez E, et al. Interleukins, from 1 to 37, and interferon- γ : Receptors, functions, and roles in diseases. *J Allergy Clin Immunol*. 2011;127(3):701-721.e70.
 25. Choy E. Interleukin-6 as a multifunctional regulator: inflammation, immune response, and fibrosis. 2017;2(Suppl 2):1–5.
 26. White WB, Siddiqui E, Tat T, Franks B, Schermer CR. Cardiovascular safety of mirabegron: analysis of an integrated clinical trial database of patients with overactive bladder syndrome. *J Am Soc Hypertens* [Internet]. 2018;12(11):768-778.e1. Available from: <https://doi.org/10.1016/j.jash.2018.08.001>
 27. Nachawati D, Patel J. Alpha Blockers. *Encycl Toxicol* Third Ed. 2021;
 28. Robinson D, Cardozo L. Solifenacin : pharmacology and clinical efficacy. *Expert Rev Clin Pharmacol*. 2009;2(3):239–53.
 29. Chapple CR, Siddiqui E. Mirabegron for the treatment of overactive bladder: a review of efficacy, safety and tolerability with a focus on male, elderly and antimuscarinic poor-responder populations, and patients with OAB in Asia. *Expert Rev Clin Pharmacol*. 2017;10(2):131–51.
 30. Zhou L, Cai X, Li H, Wang K. Effects of α -Blockers, Antimuscarinics, or Combination Therapy in Relieving Ureteral Stent-Related Symptoms: A Meta-Analysis. *J Endourol*. 2015 Jun;29(6):650–6.
 31. Liu Q, Luo D, Yang T, Liao B, Li H, Wang KJ. Protective Effects of Antimuscarinics on the Bladder Remodeling after Bladder Outlet Obstruction. *Cell Physiol Biochem*. 2017;44(3):907–19.
 32. Wood MW, Breitschwerdt EB, Gookin JL. Autocrine effects of interleukin-6 mediate acute-phase proinflammatory and tissue-reparative transcriptional

- responses of canine bladder mucosa. *Infect Immun.* 2011;79(2):708–15.
33. Segarra-Medrano A, Carnicer-Caceres C, Valtierra-Carmeno N, Agraz-Pamplona I, Ramos-Terrades N, Jatem Escalante E, et al. Valor de los niveles urinarios de interleucina 6, factor de crecimiento epidérmico, proteína quimioattractante de monocitos de tipo 1 y factor de crecimiento transformante β 1 para la predicción de la extensión de las lesiones de fibrosis en biopsias de enf. *Nefrología* [Internet]. 2017;37(5):531–8. Available from: <http://dx.doi.org/10.1016/j.nefro.2016.11.017>
34. Grover S, Srivastava A, Lee R, Tewari AK, te AE. Role of inflammation in bladder function and interstitial cystitis. *Ther Adv Urol.* 2011;3(1):19–33.
35. Alexander K, Palinrungi A, Palinrungi M, Kholis K, Syahrir S, Syarif S, Faruk M (2020) Effectiveness of Tamsulosin compared with Mirabegron in treatment of Double-J Stent-Related Lower Urinary Tract Symptoms. *Int J Med Rev Case Reports* 4:8–13
36. Yavuz A, Kilinc MF, Aydin M, Ofluoglu Y, Bayar G (2020) Does tamsulosin or mirabegron improve ureteral stent-related symptoms? A prospective placebo-controlled study. *Low Urin Tract Symptoms* 1–5
37. Hasbi BE, Palinrungi MA, Kholis K, Bakri S, Syahrir S, Kasim F (2021) The Effectiveness Between Tamsulosin and Solifenacin Combined with Tamsulosin on Treatment of Ureteral Stent-Related Symptoms. *Nusant Med Sci J* 6:1–17
38. Ilyas MRF, Kholis K, Palinrungi MA, Palinrungi AM, Seweng A, Syahrir S, Syarif, Faruk M (2021) Comparative effectiveness of Tamsulosin and Tadalafil therapy in ureteral stents-related symptoms (uSRS). *Med Clin Pract* 4:1–5
39. Singh R, Bhangu GS, Bansal D, Rattan A (2021) Effect of tamsulosin on ureteric stent related morbidity: a double-blind randomized placebo-controlled study. *Int Surg J* 8:294–299
40. Archer M, Dogra N, Dovey Z, et al. (2021) Role of α - and β -adrenergic signaling in phenotypic targeting: significance in benign and malignant urologic disease. *Cell Commun Signal* 19:1–21
41. Todorova VB, Baxan N, Delahaye M, Harding SE, Rankin SM (2023) Drug-based mobilisation of mesenchymal stem/stromal cells improves cardiac function post myocardial infarction. *Dis Model Mech* 16:1–10
42. Kovács ZZA, Szűcs G, Freiwan M, et al. (2021) Comparison of the antiremodeling effects of losartan and mirabegron in a rat model of uremic cardiomyopathy. *Sci Rep* 11:1–18
43. da Silva CPV, Calmasini F, Alexandre EC, Raposo HF, Delbin MA, Monica FZ, Zanesco A (2021) The effects of mirabegron on obesity-induced inflammation and insulin resistance are associated with brown adipose tissue activation but not beigeing in the subcutaneous white adipose tissue. *Clin Exp Pharmacol Physiol* 48:1477–1487
44. Ateyah MA, Abdulridha MK, Alkabee MJ (2021) Effects of Saw Palmetto Therapy on some Inflammatory Biomarkers in a Sample of Iraqi Male with Symptomatic Benign Prostatic Hyperplasia. *Al Mustansiriyah J Pharm Sci* 21:1–9
45. Sun L, Sun C, Zhou S, Zhang L, Hu W (2021) Tamsulosin attenuates high

- glucose- induced injury in glomerular endothelial cells. *Bioengineered* 12:5184–5194
- 46. Ye Y, Xu Q, Wan M (2021) Tamsulosin combined with levofloxacin can effectively improve the clinical symptoms and reduce inflammation in CP/CPPS patients. *Int J Clin Exp Med* 14:326–332
 - 47. Bantis A, Tsakaldimis G, Zissimopoulos A, Giannakopoulos S, Kalaitzis C, Pitiakoudis M, Polichronidis A, Toulopoulos S (2014) Can Tumor Necrosis Factor- α and Interleukin-6 Be Used as Prognostic Markers of Infection following Ureteroscopic Lithotripsy? *ISRN Urol* 2014:1–5
 - 48. Cinar O, Tanidir Y, Ozer S, et al. (2020) Effects of mirabegron on JJ stent-related symptoms: A multicentric study. *Int J Clin Pract* 00:1–5
 - 49. Wang W, Jian B, Qiao S, Ji J, Wang L (2018) Effect of Tamsulosin on the Patients with Lower Ureteral Stones after Extracorporeal Shock Wave Lithotripsy and Serum Levels of IL-6, IL-10 and CRP. *Prog Mod Biomed* 18:4099–4102
 - 50. Niemczyk G, Fus L, Czarzasta K, Jesion A, Radziszewski P, Gornicka B, Cudnoch-Jedrzejewska A (2020) Expression of Toll-Like Receptors in the Animal Model of Bladder Outlet Obstruction. *Biomed Res Int.* <https://doi.org/10.1155/2020/6632359>
 - 51. Zhang K, Yan W, Li H, Chen J, Wang Q, Chai Y, Yuan L, Zhu G (2022) Comparison of Mirabegron plus Tamsulosin and Tamsulosin Monotherapy for the Treatment of Ureteral Stent-Related Symptoms: A Prospective Randomized Study. *Urol Int* 106:1226–1232

Lampiran Tabel 1

Statistics

	Jenis Kelamin	IMT	Suku	PKJ kode
N	Valid	20	20	20
	Missing	0	0	0

Jenis Kelamin

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Perempuan	10	50,0	50,0
	Laki-laki	10	50,0	100,0
	Total	20	100,0	100,0

IMT

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Kurang (<18,5)	2	10,0	10,0
	Baik (18,5-24,9)	11	55,0	65,0
	Obesitas (>25)	7	35,0	100,0
	Total	20	100,0	100,0

Suku

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Bugis	16	80,0	80,0
	Banjar	2	10,0	90,0
	Toraja	2	10,0	100,0
	Total	20	100,0	100,0

PKJ kode

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	IRT	9	45,0	45,0
	Wiraswasta	3	15,0	60,0
	Karyawan	3	15,0	75,0
	PNS	1	5,0	80,0
	Pensiunan	2	10,0	90,0
	Petani	1	5,0	95,0
	Ojek	1	5,0	100,0
	Total	20	100,0	100,0

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Umur	,210	20	,021	,899	20	,040

a. Lilliefors Significance Correction

Descriptives

		Statistic	Std. Error
Umur	Mean	43,60	2,981
	95% Confidence Interval for	Lower Bound	37,36
	Mean	Upper Bound	49,84
	5% Trimmed Mean		43,78
	Median		42,00
	Variance		177,726
	Std. Deviation		13,331
	Minimum		23
	Maximum		61
	Range		38
	Interquartile Range		21
	Skewness		-,185 ,512
	Kurtosis		-1,448 ,992

Lampiran Tabel 2

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Umur	,210	20	,021	,899	20	,040

a. Lilliefors Significance Correction

Descriptives

		Statistic	Std. Error
Umur	Mean	43,60	2,981
	95% Confidence Interval for Mean	Lower Bound	37,36
	Mean	Upper Bound	49,84
	5% Trimmed Mean		43,78
	Median		42,00
	Variance		177,726
	Std. Deviation		13,331
	Minimum		23
	Maximum		61
	Range		38
	Interquartile Range		21
	Skewness	-,185	,512
	Kurtosis	-1,448	,992

Jenis Kelamin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Perempuan	10	50,0	50,0	50,0
	Laki-laki	10	50,0	50,0	100,0
	Total	20	100,0	100,0	

IMT

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Kurang (<18,5)	2	10,0	10,0	10,0
	Baik (18,5-24,9)	11	55,0	55,0	65,0
	Obesitas (>25)	7	35,0	35,0	100,0
	Total	20	100,0	100,0	

Suku

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Bugis	16	80,0	80,0	80,0
	Banjar	2	10,0	10,0	90,0
	Toraja	2	10,0	10,0	100,0
	Total	20	100,0	100,0	

PKJ kode

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	IRT	9	45,0	45,0	45,0
	Wiraswasta	3	15,0	15,0	60,0
	Karyawan	3	15,0	15,0	75,0
	PNS	1	5,0	5,0	80,0
	Pensiunan	2	10,0	10,0	90,0
	Petani	1	5,0	5,0	95,0
	Ojek	1	5,0	5,0	100,0
	Total	20	100,0	100,0	

Crosstab

Jenis Kelamin	Perempuan	Obat			Total
		Mirabegron	Tamsulosin		
		Count	% within Obat		
Laki-laki	Perempuan	6	60,0%	4	10
		4	40,0%	6	10
	Laki-laki	4	60,0%	6	10
Total		10	100,0%	10	20
		10	100,0%	10	20

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,800 ^a	1	,371		
Continuity Correction ^b	,200	1	,655		
Likelihood Ratio	,805	1	,369		
Fisher's Exact Test				,656	,328
Linear-by-Linear Association	,760	1	,383		
N of Valid Cases	20				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 5,00.

b. Computed only for a 2x2 table

Tests of Normality

	Obat	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Umur	Mirabegron	,240	10	,108	,902	10	,229
	Tamsulosin	,189	10	,200*	,875	10	,113

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Group Statistics

	Obat	N	Mean	Std. Deviation	Std. Error Mean
Umur	Mirabegron	10	45,00	12,454	3,938
	Tamsulosin	10	42,20	14,688	4,645

Independent Samples Test

		Levene's Test for Equality of Variances						t-test for Equality of Means			
								Mean Difference		Std. Error Difference	95% Confidence Interval of the Difference
		F	Sig.	t	df	Sig. (2-tailed)	e	ce	Lower	Upper	
Umur	Equal variances assumed	,277	,605	,460	18	,651	2,800	6,090	-9,994	15,594	
	Equal variances not assumed			,460	17,5	,651	2,800	6,090	-10,019	15,619	32

Crosstab

			Obat		Total
			Mirabegron	Tamsulosin	
IMT	Kurang (<18,5)	Count	2	0	2
		% within Obat	20,0%	0,0%	10,0%
	Baik (18,5-24,9)	Count	5	6	11
		% within Obat	50,0%	60,0%	55,0%
	Obesitas (>25)	Count	3	4	7
		% within Obat	30,0%	40,0%	35,0%
	Total	Count	10	10	20
		% within Obat	100,0%	100,0%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	2,234 ^a	2	,327
Likelihood Ratio	3,007	2	,222
Linear-by-Linear Association	1,103	1	,294
N of Valid Cases	20		

a. 4 cells (66,7%) have expected count less than 5. The minimum expected count is 1,00.

Crosstab

			Obat		Total
			Mirabegron	Tamsulosin	
PKJ kode	IRT	Count	5	4	9
		% within Obat	50,0%	40,0%	45,0%
	Wiraswasta	Count	3	0	3
		% within Obat	30,0%	0,0%	15,0%
	Karyawan	Count	0	3	3
		% within Obat	0,0%	30,0%	15,0%
	PNS	Count	0	1	1
		% within Obat	0,0%	10,0%	5,0%
	Pensiunan	Count	1	1	2
		% within Obat	10,0%	10,0%	10,0%
	Petani	Count	0	1	1
		% within Obat	0,0%	10,0%	5,0%
	Ojek	Count	1	0	1

	% within Obat	10,0%	0,0%	5,0%
Total	Count	10	10	20
	% within Obat	100,0%	100,0%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	9,111 ^a	6	,167
Likelihood Ratio	12,588	6	,050
Linear-by-Linear Association	,344	1	,557
N of Valid Cases	20		

a. 14 cells (100,0%) have expected count less than 5. The minimum expected count is ,50.

Crosstab

Suku	Bugis	Obat		Total
		Mirabegron	Tamsulosin	
Suku	Bugis	Count	8	16
		% within Obat	80,0%	80,0%
Banjar		Count	2	2
		% within Obat	20,0%	0,0%
Toraja		Count	0	2
		% within Obat	0,0%	20,0%
Total		Count	10	20
		% within Obat	100,0%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	4,000 ^a	2	,135
Likelihood Ratio	5,545	2	,063
Linear-by-Linear Association	,463	1	,496
N of Valid Cases	20		

a. 4 cells (66,7%) have expected count less than 5. The minimum expected count is 1,00.

Lampiran Tabel 3

Descriptives

IL-6 serum	Mirabegron	Mean	2,104370470933934	,393375490166099
pre stent		95% Confidence Interval for Mean	Lower Bound	1,214493288185438
ureter			Upper Bound	2,994247653682430
			Bound	
		5% Trimmed Mean		2,074559860981391
		Median		1,839399660056303
		Variance		1,547
		Std. Deviation		1,243962524610041
		Minimum		,622208798013647
		Maximum		4,123123123000000
		Range		3,500914324986353
		Interquartile Range		2,078555548919548
		Skewness		,299 ,687
		Kurtosis		-1,626 1,334
IL-6 serum	Mirabegron	Mean	,980998876575459	,194522761159853
post stent		95% Confidence Interval for Mean	Lower Bound	,540957819090413
ureter			Upper Bound	1,421039934060504
			Bound	
		5% Trimmed Mean		,959135056028287
		Median		,944420008067538
		Variance		,378
		Std. Deviation		,615134982010072
		Minimum		,232423423000000
		Maximum		2,123123100000000
		Range		1,890699677000000
		Interquartile Range		1,061587932226972
		Skewness		,537 ,687
		Kurtosis		-,642 1,334

Test Statistics^a

Obat		
	IL-6 serum Operasi I –	IL-6 serum Operasi II
Mirabegron	Z	-2,803 ^b
	Asymp. Sig. (2-tailed)	,005

a. Wilcoxon Signed Ranks Test

b. Based on positive ranks.

Lampiran Tabel 4

IL-6 serum pre stent ureter	Tamsulosin	Mean	2,032270874947530	,530637904781076
		95% Confidence Interval for Mean	Lower Bound	,831884537794787
			Upper Bound	3,232657212100273
		5% Trimmed Mean	1,873301728823726	
		Median	1,406434576956139	
		Variance	2,816	
		Std. Deviation	1,678024391927752	
		Minimum	,502072890608204	
		Maximum	6,423913489515340	
		Range	5,921840598907136	
		Interquartile Range	1,319565010635070	
		Skewness	2,322	,687
		Kurtosis	6,117	1,334
IL-6 serum post stent ureter	Tamsulosin	Mean	4,378156240816292	,839191185835194
		95% Confidence Interval for Mean	Lower Bound	2,479773888822089
			Upper Bound	6,276538592810495
		Mean	Bound	
		5% Trimmed Mean	4,366993777275975	
		Median	5,238298288699999	
		Variance	7,042	
		Std. Deviation	2,653755539576844	

Minimum	,723002485358285
Maximum	8,234234340000000
Range	7,511231854641715
Interquartile Range	4,767222215918944
Skewness	-,092
Kurtosis	-1,568 1,334

Test Statistics^a

Obat	IL-6 serum Operasi I –	
	IL-6 serum Operasi II	
Tamsulosin	Z	-2,701 ^c
	Asymp. Sig. (2-tailed)	,007

a. Wilcoxon Signed Ranks Test

b. Based on positive ranks.

Lampiran Tabel 5

IL-6	Mirabegron	Mean	28,233028454870272	20,559186619605804
Urine		95% Confidence Interval for Mean	Lower Bound	-18,275082817976020
pre stent			Upper Bound	74,741139727716560
ureter				
		5% Trimmed Mean	19,932194059640604	
		Median	2,079358376013585	
		Variance	4226,802	
		Std. Deviation	65,013856558412100	
		Minimum	,754478804583476	
		Maximum	205,126597219291000	
		Range	204,372118414707530	
		Interquartile Range	17,903616973984718	
		Skewness	2,743	,687
		Kurtosis	7,687	1,334
IL-6	Mirabegron	Mean	53,814844795432620	19,391123443028754
Urine		95% Confidence Interval for Mean	Lower Bound	9,949076004080930
post				

stent	Upper	97,680613586784320	
ureter	Bound		
	5% Trimmed Mean	51,435046620765400	
	Median	17,928483321104054	
	Variance	3760,157	
	Std. Deviation	61,320116469457176	
	Minimum	,449063248680019	
	Maximum	150,016993486195360	
	Range	149,567930237515360	
	Interquartile Range	114,545058504486320	
	Skewness	,636	,687
	Kurtosis	-1,620	1,334

Test Statistics^a

Obat	IL-6 Urine Operasi I –	
	IL-6 Urine Operasi II	
Mirabegron	Z	-1,580 ^c
	Asymp. Sig. (2-tailed)	,114

a. Wilcoxon Signed Ranks Test

b. Based on positive ranks.

Lampiran Tabel 6

IL-6	Tamsulosin	Mean	4,403801446096930	,807768601608361
Urine		95% Confidence	Lower	2,576501918085086
pre stent		Interval for Mean	Bound	
ureter			Upper	6,231100974108774
			Bound	
		5% Trimmed Mean		4,416691271160063
		Median		4,538066535601001
		Variance		6,525
		Std. Deviation		2,554388603451572
		Minimum		,624259448347163
		Maximum		7,951326592710300
		Range		7,327067144363137
		Interquartile Range		4,666589195150616
		Skewness		-,087
		Kurtosis		-1,596
				1,334
IL-6	Tamsulosin	Mean	30,503758128737520	18,076594571322154
Urine		95%	Lower	-10,388339759778045
post		Confidence	Bound	
stent		Interval for	Upper	71,395856017253080
ureter		Mean	Bound	
		5% Trimmed Mean		25,281657117574657
		Median		4,878844400670000
		Variance		3267,633
		Std. Deviation		57,163211184813050
		Minimum		,778388969095970
		Maximum		154,226945489310570
		Range		153,448556520214600
		Interquartile Range		33,107837931228070
		Skewness		1,854
		Kurtosis		1,947
				1,334

Test Statistics^a

Obat	IL-6 Urine Operasi I –	
	Z	IL-6 Urine Operasi II
Tamsulosin		-,051 ^c
	Asymp. Sig. (2-tailed)	,959

- a. Wilcoxon Signed Ranks Test
b. Based on positive ranks.

Lampiran Tabel 7

Test Statistics^a

	IL-6 serum pre stent ureter	IL-6 serum post stent ureter
Mann-Whitney U	48,000	11,000
Wilcoxon W	103,000	66,000
Z	-,151	-2,948
Asymp. Sig. (2-tailed)	,880	,003
Exact Sig. [2*(1-tailed Sig.)]	,912 ^b	,002 ^b

- a. Grouping Variable: Obat
b. Not corrected for ties.

Lampiran Tabel 8

Test Statistics^a

	IL-6 Urine Operasi I	IL-6 Urine Operasi II
Mann-Whitney U	35,000	38,000
Wilcoxon W	90,000	93,000
Z	-1,134	-,907
Asymp. Sig. (2-tailed)	,257	,364
Exact Sig. [2*(1-tailed Sig.)]	,280 ^b	,393 ^b

- a. Grouping Variable: Obat
b. Not corrected for ties.

Lampiran Tabel 9

Tests of Normality

	Obat	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Delta serum	Mirabegron	,195	10	,200*	,883	10	,140
	Tamsulosin	,174	10	,200*	,946	10	,622
Delta urine	Mirabegron	,257	10	,060	,906	10	,255
	Tamsulosin	,465	10	,000	,557	10	,000

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Descriptives

	Obat	Statistic	Std. Error
Delta serum	Mirabegron	Mean	-1,123371594400000
		95% Confidence Interval for Mean	-1,63374445884055
		Lower Bound	
		Upper Bound	-,612998742915945
		5% Trimmed Mean	-1,128972225166667
		Median	-,888925218000000
		Variance	,509
		Std. Deviation	,713452050611760
		Minimum	-2,009891928000000
		Maximum	-,136039907000000
		Range	1,873852021000000
		Interquartile Range	1,413128426250000
		Skewness	-,208
		Kurtosis	-1,662
	Tamsulosin	Mean	2,345885365700000
		95% Confidence Interval for Mean	,979637136705366
		Lower Bound	
		Upper Bound	3,712133594694635
		5% Trimmed Mean	2,329381562000000
		Median	2,314175441000000

Variance	3,648	
Std. Deviation	1,909883328994746	
Minimum	-340502727000000	
Maximum	5,329341925000000	
Range	5,669844651999999	
Interquartile Range	3,377123154000000	
Skewness	,104	,687
Kurtosis	-1,410	1,334

Test Statistics^a

Delta serum		
Mann-Whitney U	1,000	
Wilcoxon W	56,000	
Z	-3,705	
Asymp. Sig. (2-tailed)	,000	
Exact Sig. [2*(1-tailed Sig.)]	,000 ^b	

a. Grouping Variable: Obat

b. Not corrected for ties.

Lampiran Tabel 10

Delta urine	Mirabegron	Mean	25,581816344600000	22,927898398143956
		95% Confidence Interval for Mean	Lower Bound	-26,284693244670848
			Upper Bound	77,448325933870850
		5% Trimmed Mean	26,515608649555556	
		Median	3,638025696500000	
		Variance	5256,885	
		Std. Deviation	72,504380899061000	
		Minimum	-114,631429900000000	
		Maximum	148,986801100000000	
		Range	263,61823100000040	
		Interquartile Range	83,157254035750000	
		Skewness	-,079	,687
		Kurtosis	1,017	1,334

Tamsulosin	Mean	26,099956681000002	18,074512136961538
	95% Confidence Interval for Mean	Lower Bound	-14,787430413710638
		Upper Bound	66,987343775710640
	5% Trimmed Mean	20,694351388444446	
	Median	-,576836610000000	
	Variance	3266,880	
	Std. Deviation	57,156625949155710	
	Minimum	-3,063509872000000	
	Maximum	152,564318500000000	
	Range	155,627828372000000	
	Interquartile Range	30,951784962250002	
	Skewness	1,889	,687
	Kurtosis	2,179	1,334

Test Statistics^a

	Delta urine
Mann-Whitney U	34,000
Wilcoxon W	89,000
Z	-1,209
Asymp. Sig. (2-tailed)	,226
Exact Sig. [2*(1-tailed Sig.)]	,247 ^b

a. Grouping Variable: Obat

b. Not corrected for ties.