

DAFTAR PUSTAKA

1. Noorisa R, Apriliwati D, Aziz A, Bayusentono S. The characteristic of patients with femoral fracture in department of orthopaedic and traumatology RSUD dr. soetomo surabaya 2013 - 2016. *J Orthop Traumatol Surabaya*. 2017;6(1):1-11.
2. Barcega B, Minasyan L. Lower extremity trauma. In: Baren JM, Rothrock SG, Brennan JA, Brown L, editor. *Pediatric emergency medicine*. Philadelphia: W.B. Saunders; 2008. p. 182-7.
3. Keehan R, Rees D, Kendrick E, Bradshaw C, Flavell E, Deglurkar M. Enhanced recovery for fractured neck of femur: a report of 3 cases. *Geriatr Orthop Surg Rehabil*. 2014;5(2):37-42.
4. Gupta N, Jain P, Kapadia M, Sanwatsarkar S, Saxena D, Maheshwari M. The effect of 0.25% levobupivacaine and 0.25% ropivacaine in fascia iliaca block in elderly patients with fracture femur given before positioning for subarachnoid block. *Int J Res Med Sci*. 2018;6(1):191-4.
5. Tosounidis TH, Sheikh H, Stone MH, Giannoudis PV. Pain relief management following proximal femoral fractures: options, issues and controversies. *Injury*. 2015;46(5):52-58.
6. Femur Shaft Fractures (Broken Thighbone) - OrthoInfo - AAOS. Tersedia dari: <https://orthoinfo.aaos.org/en/diseases-conditions/femur-shaft-fractures-broken-thighbone/>
7. Trompeter A, Newman K. Femoral shaft fractures in adults. *Orthop Trauma*. 2013;1:27(5):322-31.
8. Salminen S. Femoral shaft fractures in adults: epidemiology, fracture patterns, nonunions, and fatigue fractures. PhD [dissertation]. Helsinki: University of Helsinki; 2005.
9. Kootstra G. Femoral shaft fractures in adults : a study of 329 consecutive cases with a statistical analysis of different methods of treatment [*e-book*]; 1973 [diunduh 16 Juni 2022]. Tersedia dari: <https://research.rug.nl/en/publications/femoral-shaft-fractures-in-adults-a-study-of-329-consecutive-case>.

10. Diwan S. Fascia iliaca block - an anatomical and technical description. *JACCR*. 2015;1(1):27-30.
11. Raja SN, Carr DB, Cohen M, Finnerup NB, Flor H, Gibson S, et al. The revised international association for the study of pain definition of pain: concepts, challenges, and compromises. *Pain*. 2020;1:161(9):1976-82.
12. European Society of Regional Anesthesia and Pain Therapy. Postoperative pain management good clinical practice.pdf [Internet]. Tersedia dari: <https://www.fpmx.com.au/resources/clinicians/postoperative-pain-management-good-clinical-practice.pdf>
13. Corke P. Postoperative pain management. *Aust Prescr*. 2013. Tersedia dari: <https://www.australianprescriber.com>
14. Rao M. Acute post operative pain. *Indian J Anaesth*. 2006;50(5):340.
15. Council (U.S.) NP, Organizations JC on A of H. Pain: current understanding of assessment, management, and treatments. National Pharmaceutical Council, Incorporated; 2001; 4-12.
16. Dureja GP, Iyer RN, Das G, Ahdal J, Narang P. Evidence and consensus recommendations for the pharmacological management of pain in india. *J Pain Res*. 2017;10:709-36.
17. Swieboda P, Filip R, Prystupa A, Drozd M. Assessment of pain: types, mechanism and treatment. *Ann Agric Environ Med AAEM*. 2013;1:2-7.
18. Breivik H, Borchgrevink PC, Allen SM, Rosseland LA, Romundstad L, Hals EKB, et al. Assessment of pain. *Br J Anaesth*. 2008;101(1):17-24.
19. Maryniak K. Pain assessment and management. 2013. Tersedia dari: <https://lms.rn.com/getpdf.php/1918.pdf>
20. Ong KS, Seymour RA. Pain measurement in humans. *Surg J R Coll Surg Edinb Irel*. 2004;2(1):15-27.
21. Innovation A for C. Fascia iliaca block [Internet]. Agency for Clinical Innovation (ACI); 2022. Tersedia dari: <https://aci.health.nsw.gov.au/networks/pain-management/resources/fascia-iliaca-block>

22. O'Reilly N, Desmet M, Kearns R. Fascia iliaca compartment block. *BJA Educ.* 2019;19(6):191-7.
23. Review of current practices of peripheral nerve blocks for hip fracture and surgery. SpringerLink [Internet]. Tersedia dari: <https://link.springer.com/article/10.1007/s40140-020-00393-7>
24. Fascia iliaca compartment block: landmark and ultrasound approach [Internet]. WFSA Resource Library. Tersedia dari: <https://resources.wfsahq.org/atotw/fascia-iliaca-compartment-block-landmark-and-ultrasound-approach/>
25. Tarekegn F, Sisay A. A review of evidence-based clinical practice on fascia iliaca compartment block for lower limb injury. *Ann Med Surg* 2012. 2021;69:102758.
26. Ultrasound-guided fascia iliaca nerve block - NYSORA [internet]. Tersedia dari: <https://www.nysora.com/topics/regional-anesthesia-for-specific-surgical-procedures/lower-extremity-regional-anesthesia-for-specific-surgical-procedures/ultrasound-guided-fascia-iliaca-block/>
27. Arias MG. Levobupivacaine: a long acting local anesthetic, with less cardiac and neurotoxicity. *Update in anesthesia.* 2002;40(4):249-8.
28. Butterworth IV JF, Mackey DC, Wasnick JD. Local anesthetics. In: Morgan & Mikhail's *Clinical Anesthesiology*. 6th ed. New York, NY: McGraw-Hill Education; 2018. p.452-70.
29. Bajwa SJS, Kaur J. Clinical profile of levobupivacaine in regional anesthesia: A systematic review. *J Anaesthesiol Clin Pharmacol.* 2013;29(4):530-9.
30. McLeod GA, Burke D. Levobupivacaine. *Anaesthesia.* 2001;56(4):331-4.
31. Burlacu CL, Buggy DJ. Update on local anesthetics: focus on levobupivacaine. *The Clin Risk Manag.* 2008;4(2):381-92.
32. Li A, Wei Z, Liu Y, Shi J, Ding H, Tang H, et al. Ropivacaine versus levobupivacaine in peripheral nerve block: A PRISMA-compliant meta-analysis of randomized controlled trials. *Medicine.* 2017;96(14).
33. Kummer KK, Zeidler M, Kalpachidou T, Kress M. Role of IL-6 in the regulation of neuronal development, survival and function. *Cytokine.* 2021;144(4):82-89.

34. Sebba A. Pain: a review of interleukin-6 and its roles in the pain of rheumatoid arthritis. *Open Access Rheumatol.* 2021;13:31-43.
35. Krishnamurthy P, MN. Girish, Dileep PK. Ultrasound guided fascia iliaca block versus pericapsular nerve group for postoperative analgesia prior to spinal anesthesia for hip surgeries. *Intern J of Heal Sci.* 2022;6(1):5084-92.
36. Gao Y, He L, Hongfei H, Yu X, Jun Z, Youtan L. Effect of continous fascia iliaca compartment block on early quality of recovery after total hip arthroplasty in elderly patients: a randomized controlled trial. *J of Pain Res.* 2022;15:1837-44.
37. El-Shaarawy, Asfour MS, Rashwan DA, Amer MM, El-Menshawe SF, Elkomy MH. Comparison of three different concentrations of levobupivacaine for epidural labor analgesia: clinical effect and pharmacokinetic profile. *Annest Essays Res.* 2018; 12(1):60-66.
38. Metesky JL, Junping C, Meg R. Enchanced recovery after surgery pathway: the use of fascia iliaca blocks causes delayed ambulation after total hip arthroplasty. *World J Anesthesiol.* 2019; 8(2): 13-18.
39. Casati A, Federico, Gianluca C, Giorgio G., Crispino G., Marta P. Levobupivacaine 0,2% or 0,125% for continous sciatic nerve block: a prospective, randomized, double-blind comparison with 0,2% ropivacaine. *Anesth Analg.* 2004;99:919-23.
40. Mose OK, Udin S, Sitanggang RH, Boom CE. Perbandingan analgesia epidural menggunakan bupivakain 0,125% dengan kombinasi bupivakain 0,0625% dan fentanyl 2µg/mL terhadap nyeri dan blok motorik pada persalinan normal. *JAP.*2013;1(2):94-104.
41. Vadhanan P, Tripaty DK, S Adinarayanan. Physiological and pharmacologic aspects of peripheral nerve blocks. *J Anaesthesiol Clin Pharmacol.* 2015;31(3):384-93.
42. C.P. Kuo, S.W. Jao, K.M. Chen, C.S. Wong, C.C. Yeh, M.J.Sheen, C.T.Wu. Comparison of the effects of thoracic epidural analgesia and i.v. infusion with lidocaine on cytokine response, postoperative pain and bowel function in patients undergoing colonic surgery. *Br J Anaesth.* 2006; 97: 640–6.
43. Ibrahim A.S., Farrag W.S., Aly M.G. Ultrasound guided bilateral rectus sheath block and serum TNF- α and IL-6 after radical prostatectomy: a randomized double blinded study. *Egypt J Anaesth.* 2018; 34: 139-143.

44. Tantri A.R., Raden B.S., Singkat D.A., Christella N. Comparing the effect of classical and modified thoracolumbar interfascial plane block on postoperative pain and IL-6 level in posterior lumbar decompression and stabilization surgery. *Anesth Pain Med.* 2022; 12(2):e122174.



REKOMENDASI PERSETUJUAN ETIK

Nomor : 705/UN4.6.4.5.31/ PP36/ 2022

Tanggal: 11 Nopember 2022

Dengan ini Menyatakan bahwa Protokol dan Dokumen yang Berhubungan Dengan Protokol berikut ini telah mendapatkan Persetujuan Etik :

No Protokol	UH22100596	No Sponsor Protokol	
Peneliti Utama	dr. Eva Satya Nugraha	Sponsor	
Judul Peneliti	PENGARUH PERBEDAAN KONSENTRASI LEVOBUPIVAKAIN ISOBARIK 0,25%, 0,125% DAN 0,0625% PADA BLOK FASCIA ILIACA TERHADAP SKOR NYERI, DURASI ANALGESIA, RESCUE ANALGESIA, DAN KEBUTUHAN OPIOID PASCA OPERASI PADA OPEN REDUCTION INTERNAL FIXATION (ORIF) FEMUR		
No Versi Protokol	2	Tanggal Versi	6 Nopember 2022
No Versi PSP	2	Tanggal Versi	6 Nopember 2022
Tempat Penelitian	RSUP Dr. Wahidin Sudirohusodo Makassar		
Jenis Review	<input type="checkbox"/> Exempted <input type="checkbox"/> Expedited <input checked="" type="checkbox"/> Fullboard Tanggal 3 Nopember 2022	Masa Berlaku 11 Nopember 2022 sampai 11 Nopember 2023	Frekuensi review lanjutan
Ketua KEP Universitas Hasanuddin	Nama Prof.Dr.dr. Suryani As'ad, M.Sc.,Sp.GK (K)	Tanda tangan	
Sekretaris KEP Universitas Hasanuddin	Nama dr. Agussalim Bukhari, M.Med.,Ph.D.,Sp.GK (K)	Tanda tangan	

Kewajiban Peneliti Utama:

- Menyerahkan Amandemen Protokol untuk persetujuan sebelum di implementasikan
- Menyerahkan Laporan SAE ke Komisi Etik dalam 24 Jam dan dilengkapi dalam 7 hari dan Laporan SUSAR dalam 72 Jam setelah Peneliti Utama menerima laporan
- Menyerahkan Laporan Kemajuan (progress report) setiap 6 bulan untuk penelitian resiko tinggi dan setiap setahun untuk penelitian resiko rendah
- Menyerahkan laporan akhir setelah Penelitian berakhir
- Melaporkan penyimpangan dari prokol yang disetujui (protocol deviation / violation)
- Mematuhi semua peraturan yang ditentukan