

## DAFTAR PUSTAKA

- Afzali, T. *et al.* (2018) 'Cost-effectiveness of treatments for non-osteoarthritic knee pain conditions: A systematic review', *PLoS ONE*, 13(12). doi: 10.1371/journal.pone.0209240.
- Alfonsin, M. M. *et al.* (2019) 'Correlations among algometry, the visual analogue scale, and the numeric rating scale to assess chronic pelvic pain in women', *European Journal of Obstetrics and Gynecology and Reproductive Biology*: X, 3. doi: 10.1016/j.eurox.2019.100037.
- Anderson, K. L. *et al.* (2020) 'Risk factors for canine osteoarthritis and its predisposing arthropathies: A systematic review', *Frontiers in Veterinary Science*. doi: 10.3389/fvets.2020.00220.
- Aweid, O. *et al.* (2018) 'Treatment modalities for hip and knee osteoarthritis: A systematic review of safety', *Journal of Orthopaedic Surgery*, 26(3). doi: 10.1177/2309499018808669.
- Bae, G. *et al.* (2021) 'Prolotherapy for the patients with chronic musculoskeletal pain: systematic review and meta-analysis', *Anesthesia and Pain Medicine*, 16(1). doi: 10.17085/apm.20078.
- Bryk, F. F. *et al.* (2016) 'Exercises with partial vascular occlusion in patients with knee osteoarthritis: a randomized clinical trial', *Knee Surgery, Sports Traumatology, Arthroscopy*, 24(5). doi: 10.1007/s00167-016-4064-7.
- Carlson, A. K. *et al.* (2019) 'Characterization of synovial fluid metabolomic phenotypes of cartilage morphological changes associated with osteoarthritis', *Osteoarthritis and Cartilage*, 27(8). doi: 10.1016/j.joca.2019.04.007.
- Chang-Miller, A. M. D. (2018) *Osteoarthritis - Symptoms and causes - Mayo Clinic, Mayo Clinic*.
- Collins, N. J., Hart, H. F. and Mills, K. A. G. (2019) 'Osteoarthritis year in review 2018: rehabilitation and outcomes', *Osteoarthritis and Cartilage*. doi: 10.1016/j.joca.2018.11.010.
- Cross, M. *et al.* (2014) 'The global burden of hip and knee osteoarthritis: Estimates

from the Global Burden of Disease 2010 study', *Annals of the Rheumatic Diseases*, 73(7). doi: 10.1136/annrheumdis-2013-204763.

Décary, S. *et al.* (2018) 'Initial derivation of diagnostic clusters combining history elements and physical examination tests for symptomatic knee osteoarthritis', *Musculoskeletal Care*, 16(3). doi: 10.1002/msc.1245.

Delgado, D. A. *et al.* (2018) 'Validation of Digital Visual Analog Scale Pain Scoring With a Traditional Paper-based Visual Analog Scale in Adults', *Journal of the American Academy of Orthopaedic Surgeons Global Research and Reviews*, 2(3). doi: 10.5435/JAAOSGlobal-D-17-00088.

Eslamian, F. and Amouzandeh, B. (2015) 'Therapeutic effects of prolotherapy with intra-articular dextrose injection in patients with moderate knee osteoarthritis: A single-arm study with 6 months follow up', *Therapeutic Advances in Musculoskeletal Disease*, 7(2). doi: 10.1177/1759720X14566618.

Gupton, M. and Terreberry, R. R. (2020) *Anatomy, Bony Pelvis and Lower Limb, Knee, StatPearls*.

Hulshof, C. T. J. *et al.* (2019) 'WHO/ILO work-related burden of disease and injury: Protocol for systematic reviews of exposure to occupational ergonomic risk factors and of the effect of exposure to occupational ergonomic risk factors on osteoarthritis of hip or knee and selected other musculoskeletal diseases', *Environment International*, 125. doi: 10.1016/j.envint.2018.09.053.

Hyland, S. and Varacallo, M. (2019) 'Anatomy, Bony Pelvis and Lower Limb, Popliteal Region', *StatPearls Publishing*, 01.

Jang, S., Lee, K. and Ju, J. H. (2021) 'Recent updates of diagnosis, pathophysiology, and treatment on osteoarthritis of the knee', *International Journal of Molecular Sciences*. doi: 10.3390/ijms22052619.

Ji, M. and Yu, Q. (2015) 'Primary osteoporosis in postmenopausal women', *Chronic Diseases and Translational Medicine*, 1(1). doi: 10.1016/j.cdtm.2015.02.006.

- Juge, P.-A. *et al.* (2016) 'Are metabolic factors associated with shoulder osteoarthritis? A multicentric study', *Osteoarthritis and Cartilage*, 24. doi: 10.1016/j.joca.2016.01.416.
- Juneja, P., Munjal, A. and Hubbard, J. B. (2022) 'Anatomy , Joints', *StatPearls [Internet]*.
- Khairurizal, K. (2019) 'Perbandingan Pengaruh Kombinasi Latihan Hold Relax Dan Open Kinetic Chain Dengan Latihan Hold Relax Dan Close Kinetic Chain Terhadap Peningkatan Kemampuan Fungsional Pasien Osteoarthritis Knee', *Nusantara Medical Science Journal*, 4(2). doi: 10.20956/nmsj.v4i2.6564.
- Kim, M. J. *et al.* (2020) 'Association of the western ontario and mcmaster universities osteoarthritis index (WOMAC) with muscle strength in community-dwelling elderly with knee osteoarthritis', *International Journal of Environmental Research and Public Health*, 17(7). doi: 10.3390/ijerph17072260.
- Kim, M. K. and Yoo, K. T. (2017) 'The effects of open and closed kinetic chain exercises on the static and dynamic balance of the ankle joints in young healthy women', *Journal of Physical Therapy Science*, 29(5). doi: 10.1589/jpts.29.845.
- Kloppenburg, M. *et al.* (2019) '2018 update of the EULAR recommendations for the management of hand osteoarthritis', *Annals of the Rheumatic Diseases*, 78(1). doi: 10.1136/annrheumdis-2018-213826.
- Kwon, Y. J. *et al.* (2013) 'The effect of open and closed kinetic chain exercises on dynamic balance ability of normal healthy adults', *Journal of Physical Therapy Science*, 25(6). doi: 10.1589/jpts.25.671.
- Long, H. *et al.* (2022) 'Prevalence Trends of Site-Specific Osteoarthritis From 1990 to 2019: Findings From the Global Burden of Disease Study 2019', *Arthritis and Rheumatology*, 74(7). doi: 10.1002/art.42089.
- Manlapaz, D. G. *et al.* (2019) 'Risk Factors for Falls in Adults with Knee Osteoarthritis: A Systematic Review', *PM and R*. doi: 10.1002/pmrj.12066.

- Martel-Pelletier, J. *et al.* (2019) 'A new decision tree for diagnosis of osteoarthritis in primary care: international consensus of experts', *Aging Clinical and Experimental Research*, 31(1). doi: 10.1007/s40520-018-1077-8.
- Michael, J. W.-P., Schlüter-Brust, K. U. and Eysel, P. (2010) 'The Epidemiology, Etiology, Diagnosis, and Treatment of Osteoarthritis of the Knee', *Deutsches Ärzteblatt international*. doi: 10.3238/arztebl.2010.0152.
- Nissen, N. *et al.* (2022) 'Clinicians' beliefs and attitudes to physical activity and exercise therapy as treatment for knee and/or hip osteoarthritis: a scoping review', *Osteoarthritis and Cartilage*. doi: 10.1016/j.joca.2021.11.008.
- Øiestad, B. E. *et al.* (2015) 'Knee extensor muscle weakness is a risk factor for development of knee osteoarthritis. A systematic review and meta-analysis', *Osteoarthritis and Cartilage*. doi: 10.1016/j.joca.2014.10.008.
- Prevention, C. for D. C. and (2018) 'Osteoarthritis (OA) | Basics | Arthritis | CDC', *Osteoarthritis (OA)*.
- Rabago, D. *et al.* (2013) 'Dextrose prolotherapy for knee osteoarthritis: A randomized controlled trial', *Annals of Family Medicine*, 11(3). doi: 10.1370/afm.1504.
- Raj, M. A. and Bubnis, M. A. (2019) *Knee Meniscal Tears, StatPearls*.
- Rannou, F., Pelletier, J. P. and Martel-Pelletier, J. (2016) 'Efficacy and safety of topical NSAIDs in the management of osteoarthritis: Evidence from real-life setting trials and surveys', *Seminars in Arthritis and Rheumatism*, 45(4). doi: 10.1016/j.semarthrit.2015.11.007.
- 'Relieving Pain in America: A Blueprint for Transforming Prevention, Care, Education, and Research' (2016) *Military medicine*, 181(5). doi: 10.7205/MILMED-D-16-00012.
- Riddle, D. L. and Perera, R. A. (2020) 'The WOMAC pain scale and crosstalk from co-occurring pain sites in people with knee pain: A causal modeling study', *Physical Therapy*, 100(10). doi: 10.1093/ptj/pzaa098.
- Sam, N. *et al.* (2022) 'Effectiveness of Prolotherapy Combined with Physical Therapy Versus Physical Therapy Only for Frozen Shoulder: A Case

Report', *American Journal of Case Reports*, 23. doi: 10.12659/AJCR.936995.

Talsania, M. and Scofield, R. H. (2017) 'Menopause and Rheumatic Disease', *Rheumatic Disease Clinics of North America*. doi: 10.1016/j.rdc.2016.12.011.

Tu, C. *et al.* (2019) 'An extensive review regarding the adipokines in the pathogenesis and progression of osteoarthritis', *Cytokine*. doi: 10.1016/j.cyto.2018.06.019.

Wang, H. and Ma, B. (2022) 'Healthcare and Scientific Treatment of Knee Osteoarthritis', *Journal of Healthcare Engineering*. doi: 10.1155/2022/5919686.

**L**

**A**

**M**

**P**

**I**

**R**

**A**

**N**

## Lampiran 1. Biodata Penulis



Nama Mahasiswa:	Annisa Larasati
No. Stambuk:	C011201203
Tempat, Tanggal Lahir :	Makassar, 31 juli 2002
Alamat :	Jl. Bandara Baru
Riwayat Pendidikan :	SD Angkasa II SMP 2 Unggulan Maros SMA 1 Maros
Nama Orang Tua :	Nur Alam (Ayah) Herawati (Ibu)
Nomor HP Mahasiswa :	082293872239
Email Mahasiswa :	Annisalarasati3107@gmail.com
Dosen Pembimbing :	Dr.dr. Nuralam Sam, Sp.KFR.,M.S(K)
Nomor HP Pembimbing :	08990501396

*Quality Assessment Tool for Case Control Studies*

Criteria	Yes	No	Other (CD, NR, NA)*
1. Was the research question or objective in this paper clearly stated and appropriate?			
2. Was the study population clearly specified and defined?			
3. Did the authors include a sample size justification?			
4. Were controls selected or recruited from the same or similar population that gave rise to the cases (including the same timeframe)?			
5. Were the definitions, inclusion and exclusion criteria, algorithms or processes used to identify or select cases and controls valid, reliable, and implemented consistently across all study participants?			
6. Were the cases clearly defined and differentiated from controls?			
7. If less than 100 percent of eligible cases and/or controls were selected for the study, were the cases and/or controls randomly selected from those eligible?			
8. Was there use of concurrent controls?			
9. Were the investigators able to confirm that the exposure/risk occurred prior to the development of the condition or event that defined a participant as a case?			
10. Were the measures of exposure/risk clearly defined, valid, reliable, and implemented consistently (including the same time period) across all study participants?			
11. Were the assessors of exposure/risk blinded to the case or control status of participants?			
12. Were key potential confounding variables measured and adjusted statistically in the analyses? If matching was used, did the investigators account for matching during study analysis?			



*Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies*

Criteria	Yes	No	Other (CD, NR, NA)*
1. Was the research question or objective in this paper clearly stated and appropriate?			
2. Was the study population clearly specified and defined?			
3. Did the authors include a sample size justification?			
4. Were controls selected or recruited from the same or similar population that gave rise to the cases (including the same timeframe)?			
5. Were the definitions, inclusion and exclusion criteria, algorithms or processes used to identify or select cases and controls valid, reliable, and implemented consistently across all study participants?			
6. Were the cases clearly defined and differentiated from controls?			
7. If less than 100 percent of eligible cases and/or controls were selected for the study, were the cases and/or controls randomly selected from those eligible?			
8. Was there use of concurrent controls?			
9. Were the investigators able to confirm that the exposure/risk occurred prior to the development of the condition or event that defined a participant as a case?			
10. Were the measures of exposure/risk clearly defined, valid, reliable, and implemented consistently (including the same time period) across all study participants?			
11. Were the assessors of exposure/risk blinded to the case or control status of participants?			
12. Were key potential confounding variables measured and adjusted statistically in the analyses? If matching was used, did the investigators account for matching during study analysis?			

*Quality Assessment Tool for Before-After (Pre-Post) Studies With No Control Group*

Criteria	Yes	No	Other (CD, NR, NA)*
1. Was the research question or objective in this paper clearly stated and appropriate?			
2. Was the study population clearly specified and defined?			
3. Did the authors include a sample size justification?			
4. Were controls selected or recruited from the same or similar population that gave rise to the cases (including the same timeframe)?			
5. Were the definitions, inclusion and exclusion criteria, algorithms or processes used to identify or select cases and controls valid, reliable, and implemented consistently across all study participants?			
6. Were the cases clearly defined and differentiated from controls?			
7. If less than 100 percent of eligible cases and/or controls were selected for the study, were the cases and/or controls randomly selected from those eligible?			
8. Was there use of concurrent controls?			
9. Were the investigators able to confirm that the exposure/risk occurred prior to the development of the condition or event that defined a participant as a case?			
10. Were the measures of exposure/risk clearly defined, valid, reliable, and implemented consistently (including the same time period) across all study participants?			
11. Were the assessors of exposure/risk blinded to the case or control status of participants?			
12. Were key potential confounding variables measured and adjusted statistically in the analyses? If matching was used, did the investigators account for matching during study analysis?			

*Quality Assessment Tool for Case-Series Studies*

Criteria	Yes	No	Other (CD, NR, NA)*
1. Was the study question or objective clearly stated?			
2. Was the study population clearly and fully described, including a case definition?			
3. Were the cases consecutive?			
4. Were the subjects comparable?			
5. Was the intervention clearly described?			
6. Were the outcome measures clearly defined, valid, reliable, and implemented consistently across all study participants?			
7. Was the length of follow-up adequate?			
8. Were the statistical methods well-described?			
9. Were the results well-described?			