

DAFTAR PUSTAKA

1. Morgan GE, Michail MS, Murray MJ (2006). Local Anesthesia. Clinical Anesthesiology. 4thed. New York: Mc Graw-Hill; 263-75.
2. Kaye KW. Changing trends in urology practice: increasing outpatient surgery. Austr New Zealand J Surg 2015;65(1):31
3. Glass PS, Hardman D, Kamiyama Y, Quill TJ, Marton G, Donn KH, et al. Preliminary pharmacokinetics and pharmacodynamics of an ultra-short-acting opioid: L Remifentanil (GI87084B). Anesth Analg 2013;77(5):1031-40.
4. Douma MR, Verway RA, Kam-Endtz CE, van der Linden PD, Stienstra R. Obstetric analgesia: a comparison of patient-controlled meperidine, Remifentanil, and fentanyl in labour. Br J Anaesth 2010; 104: 209-15.
5. Stoelting RK, Hillier SC (2006). Local Anesthetics. *Handbook of Pharmacology and Physiology in Anesthetic Practice* 2nd ed. Philadelphia: Lippincott Williams And Wilkins; 7: 180-209.
6. Balcio glu O, Akin S, Demir S, Aribogan A. Patient-controlled intravenous analgesia with Remifentanil in nulliparous subjects in labor. Expert Opin Pharmacother 2007; 8: 3089-96.
7. I Constant., et al. "Reflex pupillary dilatation in response to skin incision and alfentanil in children anaesthetized with sevoflurane: a more sensitive measure of noxious stimulation than the commonly used variables". British Journal of Anaesthesia 96.5 (2006): 614-619.
8. Park HP, Hwang JW, Park SH, Jeon YT, et.al (2007. The Effects of Glossopharyngeal Nerve Block on Postoperative Pain Relief After Tonsillectomy, The Importance of the Extent of Obtunded Gag Reflex as a Clinical Indicator. *Anesth Analg J. Med*; 105 (1): 267-71
9. Rudra A (2005). Airway Management in Obstetric. *Indian J. Anesth*; 49(4): 328-35.



Mari P, Gupta D, Kumar A, Singh (2005). Opioid Sparing Duringotrakheal Intubation Using Mccoy Laryngoscope In Neurosurgical

- Patient: The Comparison Haemodynamic Change With Macintosh Blade. *J. Postgrad Med*; 51(1): 260-5.
11. Hagberg C, Georgi R, Krier C (2005). Complication of Managing the Airway Responses to Intubation. *Best Practice And Research Clinical Anaesthesiology*. USA: Elsevier; 7: 641-69.
 12. Finucane BT, Tsui BCH, Santora AH (2011). Anatomy of the Airway. *Principles of Airway Management 4th ed.* New York: Springer; 1: 10-7.
 13. Orebaugh, Steven L (2007). Atlas of Airway Management: Technique and Tools. 1st ed. Lippincott Williams and Wilkins; 1-6.
 14. Calder I, Pearce A (2005). Tracheal intubation of the Adult Patient. In
 15. Talmage,et al. (2001) Remifentanil Pharmacokinetics and Pharmacodynamics. University of Utah school of medicine.
 16. Ben et al. (2016) Remifentanil use in Anaesthesia. and Critical Care. Anasthesia tutuorial of the week.
 17. Minto F Charles. (2007). Pharmacokinetics and Pharmacodynamics of Remifentanil. American Society of anesthesiologist
 18. Melljeans et al. (2003). Remifentanil versus fentanyl for analgesia based sedation to provide patient comfort in the intensive care unit: a randomized, double-blind controlled trial.
 19. Somers TJ, Kurakula PC, Criscione-Schreiber L, Keefe FJ, Clowse MEB. Self-efficacy and pain catastrophizing in systemic lupus erythematosus: relationship to pain, stiffness, fatigue, and psychological distress. *Arthritis Care Res.* 2012;64:1334-40
 20. M Larson., et al. "Pupillary response to noxious stimulation during isoflurane and propofol anesthesia". *Anesthesia and Analgesia* 76.5 (1993): 1072-1078.
 21. S Isnardon., et al. "Pupilometry to detect pain response during general anaesthesia following unilateral popliteal sciatic nerve block". *European Journal of Anaesthesiology* 30.7



ez Pournajafi, et al. Endotracheal Intubation in Preeclamptic Parturient Candidate for Cesarean Delivery. *Anesth Pain.* 2012;2(2):90-93. DOI: 10.1128/aapm.688

23. Bernd Muellejans et al. Remifentanil versus fentanyl for analgesia based sedation to provide patient comfort in the intensive care unit: a randomized,double-blind controlled trial [ISRCTN43755713]. *Critical Care* 2004, 8:R1-R11 (DOI 10.1186/cc2398)
24. Cevik,M. Celik, P. M. Clark, and C.Macit. Sedation and Analgesia in Intensive Care: A Comparison of Fentanyl and Remifentanil. Hindawi Publishing Corporation Pain Research and Treatment 2011. doi:10.1155/2011/650320
25. Karabinis A, Hantson P, Speelberg B, Stergiopoulos S, Illievich UM, Maas A, Upadhyaya BK: A Remifentanil-based techniquefor analgesia and sedation in ICU patients with neurotrauma: preliminary data [abstract A549]. *Intensive Care Med* 2001, Suppl 2:S275.
26. Jhon Guy MD, et al. Comparison of Remifentanil and fentanyl in patients undergoing craniotomy for supratentorial space occupying lesion. 2007. American association of anesthesiologist.
27. Q.-Y. Yang, F.-S. Xue, X. Liao et al., “Comparison of bolusre mifentanil versus bolus fentanyl for blunting cardiovascular intubation responses in children: a randomized, double-blind study,” *Chinese Medical Journal*, vol. 122, no. 1, pp. 44–50,2009.
28. Mangano DT, Siliciano D, Hollenberg M, et al. Postoperative myocardial ischemia. Therapeutic trials using intensive analgesia following surgery. The Study of Perioperative Ischemia (SPI) Research Group. *Anesthesiology* 2012; 76: 342±53.

