

## DAFTAR PUSTAKA

- American Diabetes Association (2013) ‘Diagnosis and classification of diabetes mellitus.’, *Diabetes care*, 36 Suppl 1(Suppl 1), pp. S67-74. doi: 10.2337/dc13-S067.
- Anderson, J. L. and Morrow, D. A. (2017) ‘Acute Myocardial Infarction.’, *The New England journal of medicine*, 376(21), pp. 2053–2064. doi: 10.1056/NEJMra1606915.
- Andrade, G. N. et al. (2021) ‘Prolonged heart rate recovery time after 6-minute walk test is an independent risk factor for cardiac events in heart failure: a prospective cohort study’, *Physiotherapy*. doi: 10.1016/j.physio.2021.03.012.
- Asue, B. H. Y., Izuno, Y. M. and Arada, E. H. (2019) ‘Review Coronary artery spasm — Clinical features , pathogenesis and treatment —’, 95(2), pp. 53–66.
- Beatty, A. L. and , Schiller, Nelson B. , and Whooley, M. A. (2013) ‘Six-minute walk test as a prognostic tool in stable coronary heart disease: data from the Heart and Soul Study’, 172(14), pp. 1096–1102. doi: 10.1001/archinternmed.2012.2198.Six-minute.
- Boccara, F. et al. (2020) ‘HIV Infection and Long-Term Residual Cardiovascular Risk After Acute Coronary Syndrome.’, *Journal of the American Heart Association*, 9(17), p. e017578. doi: 10.1161/JAHA.119.017578.
- Cahalin, L. P. et al. (2013) ‘Heart rate recovery after the 6 min walk test rather than distance ambulated is a powerful prognostic indicator in heart failure with reduced and preserved ejection fraction: a comparison with cardiopulmonary exercise testing.’, *European journal of heart failure*, 15(5), pp. 519–527. doi: 10.1093/eurjhf/hfs216.
- Choudhury, T., West, N. E. and El-Omar, M. (2016) ‘ST elevation myocardial infarction.’, *Clinical medicine (London, England)*, 16(3), pp. 277–282. doi: 10.7861/clinmedicine.16-3-277.
- Cole, C. R., Blackstone, E. H., Pashkow, F. J., Snader, C. E., & Lauer, M. S. (2000). Heart rate recovery immediately after exercise as a predictor of mortality. *Journal of Cardiopulmonary Rehabilitation and*

*Prevention*, 20(2), 131-132.

- Dar, M. I. *et al.* (2020) ‘Prevalence of type 2 diabetes mellitus and association of HbA1c with severity of coronary artery disease in patients presenting as non-diabetic acute coronary syndrome.’, *The Egyptian heart journal : (EHJ) : official bulletin of the Egyptian Society of Cardiology*, 72(1), p. 66. doi: 10.1186/s43044-020-00101-0.
- Dewi, R. *et al.* (2022) ‘IJNHS Predictor of Major Adverse Cardiac Event ( MACE ) in Acute Coronary Syndrome ( ACS ) patients : A Scoping Review’, 5(4). doi: 10.35654/ijnhs.v5i4.634.
- Dimopoulos, S., Manetos, C. and Panagopoulou, N. (2015) ‘The Prognostic Role of Heart Rate Recovery after Exercise in Health and Disease’, 2(2).
- Diniz, L. S. *et al.* (2017) ‘Safety of early performance of the six-minute walk test following acute myocardial infarction: a cross-sectional study’, *Brazilian Journal of Physical Therapy*, 21(3), pp. 167–174. doi: 10.1016/j.bjpt.2017.03.013.
- Elshazly, A. *et al.* (2018) ‘Effect of exercise training on heart rate recovery in patients post anterior myocardial infarction’, *The Egyptian Heart Journal*, pp. 4–6. doi: 10.1016/j.ehj.2018.04.007.
- Emilia, E. *et al.* (2022) ‘Acute coronary syndromes in diabetic patients , outcome , revascularization , and antithrombotic therapy’, *Biomedicine & Pharmacotherapy*, 148(January), p. 112772. doi: 10.1016/j.bioph.2022.112772.
- Emren, S. V., Gediz, R. B., Senöz, O., Karagöz, U., Şimşek, E. Ç., Levent, F., ... & Nazli, C. (2019). Decreased heart rate recovery may predict a high SYNTAX score in patients with stable coronary artery disease. *Bosnian Journal of Basic Medical Sciences*, 19(1), 109.
- Ferreira, P. A. *et al.* (2015) ‘Safety of the Six-Minute Walk Test in Hospitalized Cardiac Patients’, 28(1), pp. 70–77.
- Goda, A., Koike, A., Hoshimoto-Iwamoto, M., Nagayama, O., Yamaguchi, K., Tajima, A., ... & Aizawa, T. (2009). Prognostic value of heart rate profiles during cardiopulmonary exercise testing in patients with cardiac disease. *International Heart Journal*, 50(1), 59-71.

- Hegazy, M. A. *et al.* (2021) ‘Myocardial Infarction : Risk Factors , Pathophysiology , Classification , Assessment and Management’, pp. 1–11. doi: 10.31579/2692-9759/056.
- Huang, Z., Chan, T. and Dong, W. (2017) ‘MACE prediction of acute coronary syndrome via boosted resampling classification using electronic medical records’, *Journal of Biomedical Informatics*, 66, pp. 161–170. doi: 10.1016/j.jbi.2017.01.001.
- Jan, S. *et al.* (2016) ‘Bulletin of the World Health Organization Catastrophic health expenditure on acute coronary events in Asia : a prospective study’, *Bulletin of the World Health Organization*, 94(November 2015), pp. 1–12. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26966330%5Cnhttp://www.ncbi.nlm.nih.gov/articlerender.fcgi?artid=PMC4773930>.
- Karjalainen, J. J., Kiviniemi, A. M., Hautala, A. J., Piira, O. P., Lepojärvi, E. S., Peltola, M. A., ... & Tulppo, M. P. (2014). Determinants and prognostic value of cardiovascular autonomic function in coronary artery disease patients with and without type 2 diabetes. *Diabetes care*, 37(1), 286–294.
- Kementerian Kesehatan Republik Indonesia (2014) *Lingkungan Sehat, Jantung Sehat*. Available at: [www.depkes.go.id](http://www.depkes.go.id) (Accessed: 8 May 2021).
- Klabunde, R. E. (2012) *Cardiovascular Physiology Concepts*. 2nd edn. Philadelphia: Lippincott Williams & Wilkins.
- Kolettis, T. M. (2018) ‘Autonomic function and ventricular tachyarrhythmias during acute myocardial infarction’, 8(1). doi: 10.5493/wjem.v8.i1.8.
- Kontos, M. C. *et al.* (2015) ‘Cardiac arrest and clinical characteristics, treatments and outcomes among patients hospitalized with ST-elevation myocardial infarction in contemporary practice: A report from the National Cardiovascular Data Registry.’, *American heart journal*, 169(4), pp. 515–22.e1. doi: 10.1016/j.ahj.2015.01.010.
- Lanza, G. A., Careri, G. and Crea, F. (2011) ‘Contemporary Reviews in Cardiovascular Medicine Mechanisms of Coronary Artery Spasm’, pp. 1774–1782. doi: 10.1161/CIRCULATIONAHA.111.037283.

- Lipinski, M. J., Vetrovec, G. W., & Froelicher, V. F. (2004). Importance of the first two minutes of heart rate recovery after exercise treadmill testing in predicting mortality and the presence of coronary artery disease in men. *The American journal of cardiology*, 93(4), 445-449.
- Malik, M. A., Bansal, R., Seth, S., Parakh, N., Roy, A., & Bahl, V. K. (2022). Prognostic Value of Resting Heart Rate and Heart Rate Recovery in Acute Decompensated Heart Failure: A Prospective Cohort Study. *Journal of the Practice of Cardiovascular Sciences*, 8(3), 161-167.
- Martalena, D. *et al.* (2012) ‘Pengaruh Hiperglikemia Admisi terhadap Major Adverse Cardiac Events Selama Perawatan pada Pasien Sindrom Koroner Akut The Effect of Hyperglycemia on Admission on Survival of Major Adverse Cardiac Events during Hospitalization in Patients with Acute Corona’.
- Mechanic OJ, Gavin M, G. S. (2022) ‘Acute Myocardial Infarction - StatPearls - NCBI Bookshelf’.
- Messinger-Rapport, B., Snader, C. E. P., Blackstone, E. H., Yu, D., & Lauer, M. S. (2003). Value of exercise capacity and heart rate recovery in older people. *Journal of the American Geriatrics Society*, 51(1), 63-68.
- Mihardja, L. *et al.* (2013) *Penyakit Tidak Menular*. Jakarta: Kementerian Kesehatan Republik Indonesia.
- Montalescot, G. (2007) ‘On behalf of the OPERA investigators. STEMI and NSTEMI are two distinct pathophysiological entities: reply’, *Eur Heart J*, 21, pp. 2685–6.
- Nishime, E. O., Cole, C. R., Blackstone, E. H., Pashkow, F. J., & Lauer, M. S. (2000). Heart rate recovery and treadmill exercise score as predictors of mortality in patients referred for exercise ECG. *Jama*, 284(11), 1392-1398.
- Pierpont, G. L. *et al.* (2013) ‘Pathophysiology of Exercise Heart Rate Recovery : A Comprehensive Analysis’, pp. 107–117. doi: 10.1111/anec.12061.
- Qiu, S. *et al.* (2017) ‘Heart Rate Recovery and Risk of Cardiovascular Events and All-Cause Mortality: A Meta-Analysis of Prospective Cohort Studies.’,

- Journal of the American Heart Association*, 6(5). doi: 10.1161/JAHA.117.005505.
- Qiu, S., Cai, X., Sun, Z., Li, L., Zuegel, M., Steinacker, J. M., & Schumann, U. (2017). Heart rate recovery and risk of cardiovascular events and all-cause mortality: a meta-analysis of prospective cohort studies. *Journal of the American Heart Association*, 6(5), e005505.
- Ramy, M. et al. (2021) ‘Prediabetes as a risk factor for major adverse cardiovascular events’, *Annals of Medicine*, 53(1), pp. 2090–2098. doi: 10.1080/07853890.2021.2000633.
- Rilantono and Rahajoe (2014) *Penyakit Kardiovaskular pada Perempuan : Tantangan Abad Ke-21*. Jakarta: Badan Penerbit Fakultas Kedokteran Universitas Indonesia.
- Schwartz, G. G. et al. (2021) ‘Relation of insulin treatment for type 2 diabetes to the risk of major adverse cardiovascular events after acute coronary syndrome : an analysis of the BETOnMACE randomized clinical trial’, *Cardiovascular Diabetology*, pp. 1–8. doi: 10.1186/s12933-021-01311-9.
- Seravalle, G., Mancia, G. and Grassi, G. (2014) ‘Role of the sympathetic nervous system in hypertension and hypertension-related cardiovascular disease.’, *High blood pressure & cardiovascular prevention : the official journal of the Italian Society of Hypertension*, 21(2), pp. 89–105. doi: 10.1007/s40292-014-0056-1.
- Shetler, K., Marcus, R., Froelicher, V. F., Vora, S., Kalisetti, D., Prakash, M., ... & Myers, J. (2001). Heart rate recovery: validation and methodologic issues. *Journal of the American College of Cardiology*, 38(7), 1980–1987.
- Smith, S. W. et al. (2012) ‘Diagnosis of ST-elevation myocardial infarction in the presence of left bundle branch block with the ST-elevation to S-wave ratio in a modified Sgarbossa rule.’, *Annals of emergency medicine*, 60(6), pp. 766–776. doi: 10.1016/j.annemergmed.2012.07.119.
- Thygesen, K. et al. (2018) ‘Fourth Universal Definition of Myocardial Infarction (2018).’, *Journal of the American College of Cardiology*, 72(18), pp.

2231–2264. doi: 10.1016/j.jacc.2018.08.1038.

Tiksnadi, B. B., Ambari, A. M. and Adriana, M. (2019) ‘Uji Jalan 6 Menit (UJ6M) pada Pasien Pasca Sindrom Koroner Akut’, *Indonesian Journal of Cardiology*, 40(1), pp. 222–231. doi: 10.30701/ijc.v40i1.913.

Vivekananthan, D. P., Blackstone, E. H., Pothier, C. E., & Lauer, M. S. (2003). Heart rate recovery after exercise is a predictor of mortality, independent of the angiographic severity of coronary disease. *Journal of the American College of Cardiology*, 42(5), 831-838.

World Health Organization (2017) *World Heart Day 2017*. Available at: [http://www.who.int/cardiovascular\\_diseases/world-heart-day-2017](http://www.who.int/cardiovascular_diseases/world-heart-day-2017) (Accessed: 8 May 2021).

Yanagisawa, S., Miki, K., Yasuda, N., Hirai, T., Suzuki, N., & Tanaka, T. (2011). The prognostic value of treadmill exercise testing in very elderly patients: heart rate recovery as a predictor of mortality in octogenarians. *Europace*, 13(1), 114-120.