

DAFTAR PUSTAKA

1. Li X, Zhou M, Shi X, Yang H, Li Y, Li J, Yang M, Yuan H. Local anaesthetic wound infiltration used for Sesarean section pain relief: a meta-analysis. *Int J Clin Exp Med.* 2015;8(6):10213-10224.
2. Borges NC, de Deus JM, Guimarães RA, Conde DM, Bachion MM, de Moura LA, Pereira LV. The incidence of chronic pain following Cesarean section and associated risk factors: a cohort of women followed up for three months. *PLoS One.* 2020;15(9):1-14.
3. Demelash G, Berhe YW, Gebregzi AH, Chekol WB. Prevalence and factors associated with postoperative pain after Cesarean section at a comprehensive specialized hospital in Northwest Ethiopia: prospective observational study. *Open Access Surg.* 2022;15(3):1-8.
4. Yam MF, Loh YC, Tan CS, Adam SK, Manan NA, Basir R. General pathways of pain sensation and the major neurotransmitters involved in pain regulation. *Int J Mol Sci.* 2018;19(8):1-10.
5. Joshi GP, Janis JE, Haas EM, Ramshaw BJ, Nihira MA, Dunkin BJ. Surgical site infiltration for abdominal surgery: a novel neuroanatomical-based approach. *Plast Reconstr Surg - Glob Open.* 2016;4(12):1-5.
6. Reddi D, Curran N. Chronic pain after surgery: pathophysiology, risk factors and prevention. *Postgrad Med J.* 2014;90(1062):222-227.
7. Gan TJ. Poorly controlled postoperative pain : prevalence, consequences, and prevention. *J Pain Res.* 2017;10(2):2287-2298.
8. Pogatzki-Zahn EM, Segelcke D, Schug SA. Postoperative pain from mechanisms to treatment. *Pain Reports.* 2017;2(2):1-10.
9. Roofthooft E, Joshi GP, Rawal N, Van de Velde M, PROSPECT guideline for elective Sesarean section: updated systematic review and procedure-specific postoperative pain management recommendations. *Anaesthesia.* 2021;76(5):665-680.
10. Jolly C, Jathières F, Keita H, Jaouen E, Guyot B, Torre A. Cesarean analgesia using levobupivacaine continuous wound infiltration: a randomized trial.

Eur J Obstet Gynecol Reprod Biol. 2015;194:125-130.

11. Chou R, Gordon DB, De Leon-Casasola OA, Rosenberg JM, Bickler S, Brennan T, Carter T, Cassidy CL, et al. Management of postoperative pain: a clinical practice guideline from the American pain society, the American society of regional anesthesia and pain medicine, and the American society of anesthesiologists' committee on regional anesthesia, executive committee, and administrative. *J Pain*. 2016;17(2):131-157.
12. Adesope O, Ituk U, Habib AS. Local anaesthetic wound infiltration for post Sesarean section analgesia: a systematic review and meta-analysis. *Eur J Anaesthesiol*. 2016;33(10):731-142.
13. Telnes A, Skogvoll E, Lonnée H. Transversus abdominis plane block vs. wound infiltration in Sesarean section: a randomised controlled trial. *Acta Anaesthesiol Scand*. 2015;59(4):496-504.
14. Thomas D, Kundra SPP, Rudingwa P, Sivakumar RK, Dorairajan G. Continuous wound infiltration of bupivacaine at two different anatomical planes for Sesarean analgesia: a randomised clinical trial. *Indian J Anaesth*. 2019;49(4):257-262.
15. Sebba A. Pain: a review of interleukin-6 and its roles in the pain of rheumatoid arthritis. *Open access rheumatol res rev*. 2021;13:31-43.
16. Kuchálik J, Magnuson A, Tina E, Gupta A. Does local infiltration analgesia reduce peri-operative inflammation following total hip arthroplasty? a randomized, double-blind study. *BMC Anesthesiol*. 2017;17(1):1-9.
17. Carvalho B, Clark DJ, Angst MS. Local and systemic release of cytokines, nerve growth factor, prostaglandin E2, and substance P in incisional wounds and serum following Cesarean delivery. *J Pain*. 2008;9(7):650-657.
18. Raja SN, Carr DB, Cohen M, Finnerup NB, Flor H, Gibson S, Keefe FJ, Mogil JS, et al. The revised international association for the study of pain definition of pain: concepts, challenges, and compromises. *pain*. 2020;161(9):1976-1782.
19. Eisenach JC, Brennan TJ. Pain after surgery. *Pain*. 2018;159(2):321–328.
20. Bonezzi C, Fornasari D, Cricelli C, Magni A, Ventriglia G. Not all pain is

- created equal: basic definitions and diagnostic work-up. *Pain Ther.* 2020;9(s1):1-15.
21. Gupta A, Kaur K, Sharma S, Goyal S, Arora S, Murthy RSR. Clinical aspects of acute post-operative pain management & its assessment. *J Adv Pharm Technol Res.* 2010;1(2):97-108.
 22. Raney EB, Thankam FG, Dilisio MF, Agrawal DK. Pain and pathogenesis of bicep tendinopathy. *Am J Transl Res.* 2017;9(6):2668-83.
 23. Doody O, Bailey ME. Understanding pain physiology and its application to person with intellectual disability. *J Intellect Disabil.* 2019;23(1):5–18.
 24. Mazda Y, Jadin S, Khan JS. Postoperative pain management. *Can J Gen Intern Med.* 2021;16(1):1011-1018.
 25. Mohamed RE, Amin MA, Omar HM. Computed tomography-guided celiac plexus neurolysis for intractable pain of unresectable pancreatic cancer. *Egypt J Radiol Nucl Med.* 2017;48(3):627-637.
 26. Sung S, Mahdy H. Sung dan Mahdy. Treasure Island (FL): Stat Pearls Publishing; 2022.
 27. Harini R, Juwitasari J, Setyowati L, Oktavia RD. Post Sesarean section pain and quality of sleep among mothers who delivered by Sesarean section under spinal anesthesia. *Malahayati Int J Nurs Heal Sci.* 2021;3(2):110-116.
 28. Borges NC, e Silva BC, Pedroso CF, Silva TC, Tatagiba BSF, Pereira LV. Postoperative pain in women undergoing Sesarean section. *Enfermeria Global.* 2017;16(4):364-373.
 29. Kintu A, Abdulla S, Lubikire A, Nabukenya MT, Igaga E, Bulamba F, Semakula D, Olufolabi AJ, et al. Postoperative pain after Cesarean section: Assessment and management in a tertiary hospital in a low-income country. *BMC Health Serv Res.* 2019;19(1):1-6.
 30. Ahmad MR, Taufik RH. Manajemen nyeri terkini pada pasien pasca seksio sesarea. *J Anestesi Obstet Indones.* 2021;4(1):63-78.
 31. Abrão J, Antunes M, Vicente Garcia L. Local anesthetics infiltration and wound healing process. *Top Local Anesth.* 2020;1-16.
 32. Safari S, Rahimzadeh P, Haghghi M. Local infiltration anesthesia: Does it

- really work? Ann Transl Med. 2015;3(18):12-24.
33. Kastelik J, Fuchs M, Krämer M, Trauzeddel RF, Ertmer M, Von Roth P, Perka C, Kirschbaum SM, et al. Local infiltration anaesthesia versus sciatic nerve and adductor canal block for fast-track knee arthroplasty: a randomised controlled clinical trial. Eur J Anaesthesiol. 2019;36(4):255-263.
 34. Joshi D, Shristi S. To determine functional visual improvement after cataract surgery in eyes with age related macular degeneration. Indian J Clin Exp Ophthalmol. 2022;7(4):655-658.
 35. Stamenkovic DM, Bezmarevic M, Bojic S, Unicstojanovic D, Stojkovic D, Slavkovic DZ, Bancevic V, Maric N, et al. Updates on wound infiltration use for postoperative pain management: a narrative review. J Clin Med. 2021;10(20):1-36.
 36. Mcgee D. Local and topical anesthesia anesthetic and analgesic techniques. In: Chanmugam A, Chudnofskey C, DeBlieux P, editors. Roberts and Hedges' clinical procedures in emergency medicine and acute care. 7th Ed. Elsevier; 2019. p. 533.
 37. Latham JL, Martin SN. Infiltrative anesthesia in office practice. Am Fam Physician. 2014;89(12):956-962.
 38. Bagatin T, Bagatin D, Šakić L, Šakić K. Impact of local infiltration anesthesia on postoperative pain management after rhinoplasty in day care surgery. Acta Clin Croat. 2019;58:62-6.
 39. Miller RD, Cohen NH, Ericson LI, Fleisher LA, Wiener-Kronish JP, Young WL. Miller's Anesthesia. Canada: Elsevier; 2015. p. 1042.
 40. Ghenaee MM, Rahmani S, Jafarabadi M. Local lidocaine 2% in postoperative pain management in Cesarean delivery. J Fam Reprod Heal. 2015;9(1):19-21.
 41. Corsini T, Cuvillon P, Forgeot A, Chapelle C, Seffert P, Chauleur C. Single-dose intraincisional levobupivacaine infiltration in sesarean postoperative analgesia: a placebo-controlled double-blind randomized trial. Ann Fr Anesth Reanim. 2013;32(1):25-30.

42. Demiraran Y, Albayrak M, Yorulmaz IS, Ozdemir I. Tramadol and levobupivacaine wound infiltration at Cesarean delivery for postoperative analgesia. *J Anesth*. 2013;27(2):175-179.
43. Wagner-Kovacec J, Povalej-Brzan P, Mekis D. Efficacy of continuous in-wound infusion of levobupivacaine and ketorolac for post-Sesarean section analgesia: a prospective, randomised, double-blind, placebo-controlled trial. *BMC Anesthesiol*. 2018;18(1):1-9.
44. Lalmand M, Wilwerth M, Fils JF, Van Der Linden P. Continuous ropivacaine subfascial wound infusion compared with intrathecal morphine for post Cesarean analgesia: a prospective, randomized controlled, double-blind study. *Anesth Analg*. 2017;125(3):907-912.
45. Larsen KR, Kristensen BB, Rasmussen MA, Rasmussen YH, Weber T, Kristensen B, Kehlet, H. Effect of high-volume systematic local infiltration analgesia in Sesarean section: a randomised, placebo - controlled trial. *Acta Anaesthesiol Scand*. 2015;59(5):632-639.
46. Nasir F, Sohail I, Sadiq H, Habib M. Local wound infiltration with ropivacaine for postoperative pain control in Sesarean section. *Cureus*. 2019;11(9):5-11.
47. Ducarme G, Sillou S, Wernet A, Davitian C, Poujade O, Ceccaldi PF, Bougeois B, Luton D. Single-shot ropivacaine wound infiltration during cesarean section for postoperative pain relief. *Gynecol Obstet Fertil*. 2012;40(1):10-13.
48. Reinikainen M, Syväoja S, Hara K. Continuous wound infiltration with ropivacaine for analgesia after Sesarean section: a randomised, placebo-controlled trial. *Acta Anaesthesiol Scand*. 2014;58(8):973-979.
49. Wang J, Zhao G, Song G, Liu J. The efficacy and safety of local anesthetic techniques for postoperative analgesia after Cesarean section: a bayesian network meta-analysis of randomized controlled trials. *J Pain Res*. 2021;14(4l):1559–1572.
50. Ryu C, Choi GJ, Jung YH, Baek CW, Cho CK, Kang H. Postoperative analgesic effectiveness of peripheral nerve blocks in Cesarean delivery: a

- systematic review and network meta-analysis. *J Pers Med.* 2022;12(4):1-10.
51. Kummer KK, Zeidler M, Kalpachidou T, Kress M. Role of IL-6 in the regulation of neuronal development, survival and function. *Cytokine.* 2021;144(4):82-89.
 52. Zhou YQ, Liu Z, Liu ZH, Chen SP, Li M, Shahveranov A, Ye DW, Tian YK. Interleukin-6: an emerging regulator of pathological pain. *J Neuroinflammation.* 2016;13(1):1-9.
 53. Cook AD, Christensen AD, Tewari D, McMahon SB, Hamilton JA. Immune cytokines and their receptors in inflammatory pain. *Trends Immunol.* 2018;39(3):240-255.
 54. Cruz FF, Rocco PRM, Pelosi P. Anti-inflammatory properties of anesthetic agents. *Crit Care.* 2017;21(1):1-10.
 55. Ikeuchi M, Kamimoto Y, Izumi M, Fukunaga K, Aso K, Sugimura N, Yokoyama M, Tani T. Effects of dexamethasone on local infiltration analgesia in total knee arthroplasty: a randomized controlled trial. *Knee Surgery, Sport Traumatol Arthrosc.* 2014;22(7):1638-1643.
 56. Sahmeddini MA, Azemati S, Motlagh EM. Local infiltration of tramadol versus bupivacaine for post cesarean section pain control: A double-blind randomized study. *Iran J Med Sci.* 2017;42(3):235-41.
 57. Bagul A, Taha R, Metcalfe MS, Brook NR, Nicholson ML. Pre-incision infiltration of local anesthetic reduces postoperative pain with no effects on bruising and wound cosmesis after thyroid surgery. *Thyroid.* 2005;15(11):1245-8.
 58. Santosh UP, Sanjay VC. Comparative study of 0.5% bupivacaine infiltration versus sterile water infiltration in tonsillar fossa during tonsillectomy performed under general anaesthesia. *Int J Otorhinolaryngol Head Neck Surg.* 2020;6(5):964.
 59. Sharifian HA, Fathololoomi MR, Fattahi Bafghi A, Safavi Naini SA. Effect of local bupivacaine infiltration on post-tonsillectomy pain. *Tanaffos.* 2006;5(1):45-9.

60. Steagall P V., Robertson S, Simon B, Warne LN, Shilo-Benjamini Y, Taylor S. Consensus guidelines on the management of acute pain in cats. *J Feline Med Surg.* 2022;24(1):4–30.
61. Euliano T, Gravenstein J, Gravenstein N, Gravenstein D. Essential Anesthesia. United States of America: Cambridge University Press; 2011.
62. Yari M, Rooshani B, Golfam P, Nazari N. Intraperitoneal bupivacaine effect on postoperative nausea and vomiting following laparoscopic cholecystectomy. *Anesthesiol Pain Med.* 2014;4(3).
63. Sherif FA El, Tohamy M, A. M. Mostafa M, Adlan S de la, Mohamed SA-B, Mansour S, et al. Effect of local wound infiltration with ketamine versus dexmedetomidine added to bupivacaine on inflammatory cytokines, a randomized clinical trial. *Open J Anesthesiol.* 2022;12(08):261–77.
64. Chang L, Ye F, Luo Q, Wang Z, Wang Y, Xia Z, Shu H. Effects of three forms of local anesthesia on perioperative fentanyl-induced hyperalgesia. *Biosci Trends.* 2018;12(2):177–84.
65. Patel L, Shah K, Padhiyar V, Patel K, Patel B. Pre-emptive Incision Infiltration Versus Post-Operative Wound Infiltration with 0.5% ropivacaine in Patients Undergoing Lumbar Laminectomy. *Indian J Clin Anaesth.* 2015;2(4):217.
66. Carvalho AL De, Bonini F, Emanuel B, Gatto O, Muraro SK, Schiavuzzo FA, Ashmawi HA, Vieira JE, et al. Preemptive activity of incision infiltration with 0.75 % ropivacaine in patients submitted to inguinal hernia repair. *Rev Dor Sao Paulo.* 2011;12(4):321–6.
67. Cantore F, Boni L, Di Giuseppe M, Giavarini L, Rovera F, Dionigi G. Pre-incision local infiltration with levobupivacaine reduces pain and analgesic consumption after laparoscopic cholecystectomy: A new device for day-case procedure. *Int J Surg.* 2008;6(1):89–92.
68. Jebeles JA, Reilly JS, Gutierrez JF, Bradley EL, Kissin I. The effect of pre-incisional infiltration of tonsils with bupivacaine on the pain following tonsillectomy under general anesthesia. *Pain.* 1991;47(3):305–8.

LAMPIRAN

Lampiran 1. Data Induk penelitian

Kelompok kontrol

No	Umur	IMT	Paritas	Leukosit	NRS					Rescue Fentanyl					Efek Samping			IL6		
					2	4	8	16	24	2	4	8	16	24	Mual	Muntah	Edema	Pre	4 Jam	8 Jam
1	39	24.07	2	9.7	1	0	4	2	2	0	0	30	30	0	Ya	Tidak	Tidak	54	60	44
2	34	27.92	3	8.5	0	0	2	2	0	0	0	0	30	0	Ya	Tidak	Tidak	46	44	22
3	19	27.27	0	6.6	0	0	1	2	2	0	0	0	30	0	Ya	Ya	Tidak	39	67	47
4	19	27.60	1	9.2	1	4	0	4	1	0	0	0	0	0	Ya	Ya	Tidak	29	90	60
5	34	28.00	1	8.8	0	0	4	4	0	0	0	0	35	0	Tidak	Tidak	Tidak	43	56	28
6	36	22.69	4	8.1	0	2	4	5	2	0	0	30	30	0	Ya	Ya	Tidak	76	98	28
7	40	26.00	5	9.6	1	0	1	0	0	0	0	0	0	0	Ya	Tidak	Tidak	78	56	31
8	41	25.19	1	9	1	0	1	0	0	0	0	0	0	0	Ya	Ya	Tidak	52	69	60

Kelompok sebelum insisi

No	Umur	IMT	Paritas	Leukosit	NRS					Rescue Fentanyl					Efek Samping			IL6		
					2	4	8	16	24	2	4	8	16	24	Mual	Muntah	Edema	Pre	4 Jam	8 Jam
1	18	22.41	0	7.5	0	0	0	1	0	0	0	0	0	0	Tidak	Tidak	Tidak	32	11	8
2	38	20.94	3	9.8	0	0	0	0	0	0	0	0	0	0	Tidak	Tidak	Tidak	16	18	17
3	17	25.00	0	8	0	0	0	1	0	0	0	0	0	0	Tidak	Ya	Tidak	17	12	4
4	39	26.25	2	7	1	0	2	2	1	0	30	0	0	30	Tidak	Tidak	Tidak	40	19	14

5	19	26.15	2	9.4	0	1	0	0	0	0	0	0	0	0	Tidak	Tidak	Tidak	30	19	17
6	28	27.08	0	8.6	0	1	1	5	2	0	0	0	30	0	Ya	Tidak	Tidak	12	18	15
7	18	26.82	0	7.7	0	0	0	2	0	0	0	0	0	0	Tidak	Tidak	Tidak	39	17	17
8	23	27.83	0	6.9	1	0	0	1	0	0	0	0	0	0	Tidak	Tidak	Tidak	39	22	20

Kelompok setelah insisi

No	Umur	IMT	Paritas	Leukosit	NRS					Rescue Fentanyl					Efek Samping			IL6		
					2	4	8	16	24	2	4	8	16	24	Mual	Muntah	Edema	Pre	4 Jam	8 Jam
1	44	25.22	0	10	0	0	1	0	0	0	0	35	0	0	Tidak	Tidak	Tidak	69	60	22
2	20	26.40	1	8.8	0	2	0	2	0	0	0	0	0	0	Tidak	Tidak	Tidak	34	37	32
3	20	27.27	0	9	0	0	0	0	1	0	0	0	0	0	Ya	Ya	Tidak	45	33	32
4	20	28.00	0	7.9	0	1	0	0	1	0	0	0	0	0	Tidak	Tidak	Tidak	50	52	29
5	20	24.07	0	8.3	1	0	0	1	0	0	0	0	0	0	Tidak	Tidak	Tidak	57	53	25
6	20	27.60	0	10	0	0	2	1	0	0	0	0	0	0	Tidak	Ya	Tidak	41	53	17
7	36	26.67	5	9.5	0	0	0	3	0	0	0	30	0	0	Ya	Tidak	Tidak	26	42	14
8	29	28.00	1	9.8	0	0	0	5	2	0	0	0	0	0	Tidak	Tidak	Tidak	53	24	28

Lampiran 2. Hasil olah data SPSS

KARAKTERISTIK SAMPEL

Tests of Normality

	Kelompok	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Umur	Tanpa	.306	8	.026	.795	8	.025
	Sebelum	.246	8	.169	.811	8	.038
	Sesudah	.369	8	.002	.730	8	.005
Paritas	Tanpa	.243	8	.184	.919	8	.425
	Sebelum	.384	8	.001	.720	8	.004
	Sesudah	.346	8	.005	.592	8	.000
BB	Tanpa	.213	8	.200*	.903	8	.308
	Sebelum	.140	8	.200*	.973	8	.924
	Sesudah	.173	8	.200*	.911	8	.364
TB	Tanpa	.226	8	.200*	.907	8	.335
	Sebelum	.217	8	.200*	.943	8	.641
	Sesudah	.259	8	.123	.918	8	.410
IMT	Tanpa	.226	8	.200*	.894	8	.254
	Sebelum	.261	8	.115	.870	8	.151
	Sesudah	.178	8	.200*	.895	8	.261

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Descriptives

		N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
						Lower Bound	Upper Bound		
Umur	Tanpa	8	32.7500	8.87613	3.13819	25.3294	40.1706	19.00	41.00
	Sebelum	8	25.0000	9.07115	3.20713	17.4163	32.5837	17.00	39.00
	Sesudah	8	26.1250	9.35701	3.30820	18.3023	33.9477	20.00	44.00
	Total	24	27.9583	9.37378	1.91341	24.0001	31.9165	17.00	44.00
Paritas	Tanpa	8	2.1250	1.72689	.61055	.6813	3.5687	.00	5.00
	Sebelum	8	.8750	1.24642	.44068	-.1670	1.9170	.00	3.00
	Sesudah	8	.8750	1.72689	.61055	-.5687	2.3187	.00	5.00
	Total	24	1.2917	1.62799	.33231	.6042	1.9791	.00	5.00
BB	Tanpa	8	65.3750	4.03334	1.42600	62.0030	68.7470	59.00	70.00
	Sebelum	8	65.1250	3.35676	1.18679	62.3187	67.9313	59.00	70.00
	Sesudah	8	65.2500	4.49603	1.58959	61.4912	69.0088	58.00	70.00
	Total	24	65.2500	3.81359	.77845	63.6397	66.8603	58.00	70.00
TB	Tanpa	8	158.4330	5.23709	1.85159	154.0546	162.8113	148.32	164.32
	Sebelum	8	160.9471	10.47657	3.70403	152.1884	169.7057	148.32	178.89
	Sesudah	8	156.4591	4.84477	1.71289	152.4088	160.5095	148.32	164.32
	Total	24	158.6131	7.23986	1.47783	155.5559	161.6702	148.32	178.89
IMT	Tanpa	8	26.0926	1.96755	.69563	24.4477	27.7375	22.69	28.00
	Sebelum	8	25.3103	2.41863	.85512	23.2883	27.3324	20.94	27.83
	Sesudah	8	26.6539	1.39825	.49436	25.4849	27.8228	24.07	28.00
	Total	24	26.0189	1.96732	.40158	25.1882	26.8497	20.94	28.00

Test of Homogeneity of Variances

	Levene Statistic	df1	df2	Sig.
Umur	.065	2	21	.937
Paritas	.310	2	21	.737
BB	.464	2	21	.635
TB	3.929	2	21	.036
IMT	1.271	2	21	.301

ANOVA

		Sum of Squares	df	Mean Square	F	Sig.
Umur	Between Groups	280.583	2	140.292	1.693	.208
	Within Groups	1740.375	21	82.875		
	Total	2020.958	23			
Paritas	Between Groups	8.333	2	4.167	1.663	.214
	Within Groups	52.625	21	2.506		
	Total	60.958	23			
BB	Between Groups	.250	2	.125	.008	.992
	Within Groups	334.250	21	15.917		
	Total	334.500	23			
TB	Between Groups	80.956	2	40.478	.756	.482
	Within Groups	1124.602	21	53.552		
	Total	1205.557	23			
IMT	Between Groups	7.285	2	3.643	.936	.408
	Within Groups	81.733	21	3.892		
	Total	89.018	23			

Test Statistics^{a,b}

	Umur	Paritas
Chi-Square	2.977	4.335
df	2	2
Asymp. Sig.	.226	.114

a. Kruskal Wallis Test

b. Grouping Variable: Kelompok

HASIL OLAH DATA NRS

HASIL OLAH DATA BERDASARKAN KELOMPOK INTERVENSI ANESTESI
UJI NORMALITAS

Tests of Normality

	Kelompok	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
NRS_2jam	Tanpa	.391	8	.001	.641	8	.000
	Sebelum	.455	8	.000	.566	8	.000
	Sesudah	.513	8	.000	.418	8	.000
NRS_4jam	Tanpa	.455	8	.000	.566	8	.000
	Sebelum	.443	8	.000	.601	8	.000
	Sesudah	.443	8	.000	.601	8	.000

NRS_8jam	Tanpa	.253	8	.140	.827	8	.055
	Sebelum	.513	8	.000	.418	8	.000
	Sesudah	.300	8	.033	.789	8	.022
NRS_16jam	Tanpa	.224	8	.200*	.866	8	.137
	Sebelum	.310	8	.023	.753	8	.009
	Sesudah	.455	8	.000	.566	8	.000
NRS_24jam	Tanpa	.377	8	.001	.693	8	.002
	Sebelum	.443	8	.000	.601	8	.000
	Sesudah	.391	8	.001	.641	8	.000

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Descriptives

		N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
						Lower Bound	Upper Bound		
NRS_2jam	Tanpa	8	.6250	.51755	.18298	.1923	1.0577	.00	1.00
	Sebelum	8	.2500	.46291	.16366	-.1370	.6370	.00	1.00
	Sesudah	8	.1250	.35355	.12500	-.1706	.4206	.00	1.00
	Total	24	.3333	.48154	.09829	.1300	.5367	.00	1.00
NRS_4jam	Tanpa	8	.2500	.46291	.16366	-.1370	.6370	.00	1.00
	Sebelum	8	.7500	1.48805	.52610	-.4940	1.9940	.00	4.00
	Sesudah	8	.3750	.74402	.26305	-.2470	.9970	.00	2.00
	Total	24	.4583	.97709	.19945	.0457	.8709	.00	4.00
NRS_8jam	Tanpa	8	2.1250	1.64208	.58056	.7522	3.4978	.00	4.00
	Sebelum	8	.1250	.35355	.12500	-.1706	.4206	.00	1.00
	Sesudah	8	1.8750	2.23207	.78916	.0089	3.7411	.00	5.00
	Total	24	1.3750	1.78916	.36521	.6195	2.1305	.00	5.00
NRS_16jam	Tanpa	8	1.8750	1.88509	.66648	.2990	3.4510	.00	5.00
	Sebelum	8	1.2500	1.66905	.59010	-.1454	2.6454	.00	5.00
	Sesudah	8	.2500	.46291	.16366	-.1370	.6370	.00	1.00
	Total	24	1.1250	1.56906	.32028	.4624	1.7876	.00	5.00
NRS_24jam	Tanpa	8	.6250	.91613	.32390	-.1409	1.3909	.00	2.00
	Sebelum	8	.7500	1.48805	.52610	-.4940	1.9940	.00	4.00
	Sesudah	8	.7500	1.03510	.36596	-.1154	1.6154	.00	2.00
	Total	24	.7083	1.12208	.22904	.2345	1.1821	.00	4.00

Test of Homogeneity of Variances

	Levene Statistic	df1	df2	Sig.
NRS_2jam	2.722	2	21	.089
NRS_4jam	3.608	2	21	.045
NRS_8jam	11.892	2	21	.000
NRS_16jam	2.539	2	21	.103
NRS_24jam	.728	2	21	.495

ANOVA

		Sum of Squares	df	Mean Square	F	Sig.
NRS_2jam	Between Groups	1.083	2	.542	2.676	.092
	Within Groups	4.250	21	.202		
	Total	5.333	23			
NRS_4jam	Between Groups	1.083	2	.542	.545	.588
	Within Groups	20.875	21	.994		
	Total	21.958	23			
NRS_8jam	Between Groups	19.000	2	9.500	3.652	.044
	Within Groups	54.625	21	2.601		
	Total	73.625	23			
NRS_16jam	Between Groups	10.750	2	5.375	2.460	.110
	Within Groups	45.875	21	2.185		
	Total	56.625	23			
NRS_24jam	Between Groups	.083	2	.042	.030	.970
	Within Groups	28.875	21	1.375		
	Total	28.958	23			

Test Statistics^{a,b}

	NRS_2jam	NRS_4jam	NRS_8jam	NRS_16jam	NRS_24jam
Chi-Square	4.672	.093	7.541	4.551	.132
df	2	2	2	2	2
Asymp. Sig.	.097	.954	.023	.103	.936

a. Kruskal Wallis Test

b. Grouping Variable: Kelompok

HASIL OLAH DATA BERDASARKAN KELOMPOK WAKTU ANESTESI

Test of Homogeneity of Variances

	Levene Statistic	df1	df2	Sig.
NRS_tanpa	5.085	4	35	.002
NRS_Sesudah	12.985	4	35	.000
NRS_Sebelum	2.785	4	35	.041

ANOVA

		Sum of Squares	df	Mean Square	F	Sig.
NRS_tanpa	Between Groups	22.600	4	5.650	3.731	.012
	Within Groups	53.000	35	1.514		
	Total	75.600	39			
NRS_Sesudah	Between Groups	16.150	4	4.038	2.906	.035
	Within Groups	48.625	35	1.389		
	Total	64.775	39			
NRS_Sebelum	Between Groups	6.500	4	1.625	1.076	.383
	Within Groups	52.875	35	1.511		
	Total	59.375	39			

Test Statistics^{a,b}

	NRS tanpa	NRS Sesudah	NRS Sebelum
Chi-Square	10.286	5.187	5.136
df	4	4	4
Asymp. Sig.	.036	.269	.274

a. Kruskal Wallis Test

b. Grouping Variable: Kelompok_5

HASIL OLAH DATA EFEK SAMPING**Crosstab**

			kelompok		Total
Mual	Tidak	Count	Tanpa	Sebelum	
		% within Mual	12.5%	87.5%	100.0%
Ya		% within kelompok	12.5%	87.5%	50.0%
		% of Total	6.3%	43.8%	50.0%
		Count	7	1	8
		% within Mual	87.5%	12.5%	100.0%
Total		% within kelompok	87.5%	12.5%	50.0%
		% of Total	43.8%	6.3%	50.0%
		Count	8	8	16
		% within Mual	50.0%	50.0%	100.0%
		% within kelompok	100.0%	100.0%	100.0%
		% of Total	50.0%	50.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	9.000 ^a	1	.003		
Continuity Correction ^b	6.250	1	.012		
Likelihood Ratio	10.124	1	.001		
Fisher's Exact Test				.010	.005
N of Valid Cases	16				

a. 4 cells (100.0%) have expected count less than 5. The minimum expected count is 4.00.

b. Computed only for a 2x2 table

Crosstab

			kelompok		Total
Muntah	Tidak	Count	Tanpa	Sebelum	
		% within Muntah	36.4%	63.6%	100.0%
Ya		% within kelompok	50.0%	87.5%	68.8%
		% of Total	25.0%	43.8%	68.8%
		Count	4	1	5
		% within Muntah	80.0%	20.0%	100.0%
		% within kelompok	50.0%	12.5%	31.3%
		% of Total	25.0%	6.3%	31.3%

Total	Count	8	8	16
	% within Muntah	50.0%	50.0%	100.0%
	% within kelompok	100.0%	100.0%	100.0%
	% of Total	50.0%	50.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.618 ^a	1	.106		
Continuity Correction ^b	1.164	1	.281		
Likelihood Ratio	2.756	1	.097		
Fisher's Exact Test				.282	.141
N of Valid Cases	16				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 2.50.

b. Computed only for a 2x2 table

Crosstab

		Kelompok2		Total
Mual2	Tidak	Tanpa	Sesudah	
Mual2	Count	1	6	7
	% within Mual2	14.3%	85.7%	100.0%
	% within Kelompok2	12.5%	75.0%	43.8%
	% of Total	6.3%	37.5%	43.8%
Ya	Count	7	2	9
	% within Mual2	77.8%	22.2%	100.0%
	% within Kelompok2	87.5%	25.0%	56.3%
	% of Total	43.8%	12.5%	56.3%
Total	Count	8	8	16
	% within Mual2	50.0%	50.0%	100.0%
	% within Kelompok2	100.0%	100.0%	100.0%
	% of Total	50.0%	50.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	6.349 ^a	1	.012		
Continuity Correction ^b	4.063	1	.044		
Likelihood Ratio	6.904	1	.009		
Fisher's Exact Test				.041	.020
N of Valid Cases	16				

a. 4 cells (100.0%) have expected count less than 5. The minimum expected count is 3.50.

b. Computed only for a 2x2 table

Crosstab

			Kelompok2		Total
			Tanpa	Sesudah	
Muntah2	Tidak	Count	4	6	10
		% within Muntah2	40.0%	60.0%	100.0%
		% within Kelompok2	50.0%	75.0%	62.5%
		% of Total	25.0%	37.5%	62.5%
	Ya	Count	4	2	6
		% within Muntah2	66.7%	33.3%	100.0%
		% within Kelompok2	50.0%	25.0%	37.5%
		% of Total	25.0%	12.5%	37.5%
Total		Count	8	8	16
		% within Muntah2	50.0%	50.0%	100.0%
		% within Kelompok2	100.0%	100.0%	100.0%
		% of Total	50.0%	50.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.067 ^a	1	.302		
Continuity Correction ^b	.267	1	.606		
Likelihood Ratio	1.082	1	.298		
Fisher's Exact Test				.608	.304
N of Valid Cases	16				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 3.00.

b. Computed only for a 2x2 table

Crosstab

			kelompok3		Total
			Sebelum	Setelah	
Mual3	Tidak	Count	7	6	13
		% within Mual3	53.8%	46.2%	100.0%
		% within kelompok3	87.5%	75.0%	81.3%
		% of Total	43.8%	37.5%	81.3%
	Ya	Count	1	2	3
		% within Mual3	33.3%	66.7%	100.0%
		% within kelompok3	12.5%	25.0%	18.8%
		% of Total	6.3%	12.5%	18.8%
Total		Count	8	8	16
		% within Mual3	50.0%	50.0%	100.0%
		% within kelompok3	100.0%	100.0%	100.0%
		% of Total	50.0%	50.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)

Pearson Chi-Square	.410 ^a	1	.522		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.417	1	.519		
Fisher's Exact Test				1.000	
N of Valid Cases	16				.500

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.50.

b. Computed only for a 2x2 table

Crosstab

		kelompok3		Total
		Sebelum	Setelah	
Muntah3	Tidak	Count	7	13
		% within Muntah3	53.8%	100.0%
		% within kelompok3	87.5%	81.3%
		% of Total	43.8%	81.3%
Ya		Count	1	3
		% within Muntah3	33.3%	100.0%
		% within kelompok3	12.5%	18.8%
		% of Total	6.3%	18.8%
Total		Count	8	16
		% within Muntah3	50.0%	100.0%
		% within kelompok3	100.0%	100.0%
		% of Total	50.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.410 ^a	1	.522		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.417	1	.519		
Fisher's Exact Test				1.000	
N of Valid Cases	16				.500

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.50.

b. Computed only for a 2x2 table

HASIL OLAH DATA NRS

Test of Homogeneity of Variances

	Levene Statistic	df1	df2	Sig.
NRS_2jam	3.847	2	21	.038
NRS_4jam	3.608	2	21	.045
NRS_8jam	6.943	2	21	.005
NRS_16jam	.248	2	21	.782
NRS_24jam	1.547	2	21	.236

ANOVA

	Sum of Squares	df	Mean Square	F	Sig.

NRS_2jam	Between Groups	.583	2	.292	1.400	.269
	Within Groups	4.375	21	.208		
	Total	4.958	23			
NRS_4jam	Between Groups	1.083	2	.542	.545	.588
	Within Groups	20.875	21	.994		
	Total	21.958	23			
NRS_8jam	Between Groups	16.333	2	8.167	6.441	.007
	Within Groups	26.625	21	1.268		
	Total	42.958	23			
NRS_16jam	Between Groups	4.083	2	2.042	.671	.522
	Within Groups	63.875	21	3.042		
	Total	67.958	23			
NRS_24jam	Between Groups	1.083	2	.542	.771	.475
	Within Groups	14.750	21	.702		
	Total	15.833	23			

Test Statistics^{a,b}

	NRS_2jam	NRS_4jam	NRS_8jam	NRS_16jam	NRS_24jam
Chi-Square	2.706	.093	8.752	1.360	1.364
df	2	2	2	2	2
Asymp. Sig.	.258	.954	.013	.507	.506

a. Kruskal Wallis Test

b. Grouping Variable: Kelompok

Uji Mann Whitney

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_2jam	Kontrol	8	9.50	76.00
	Anestesi infiltrasi setelah insisi	8	7.50	60.00
	Total	16		

Test Statistics^a

	NRS_2jam
Mann-Whitney U	24.000
Wilcoxon W	60.000
Z	-1.000
Asymp. Sig. (2-tailed)	.317
Exact Sig. [2*(1-tailed Sig.)]	.442 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_2jam	Kontrol	8	10.00	80.00
	Anestesi infiltrasi sebelum insisi	8	7.00	56.00
	Total	16		

Test Statistics^a

	NRS_2jam
Mann-Whitney U	20.000
Wilcoxon W	56.000
Z	-1.567
Asymp. Sig. (2-tailed)	.117
Exact Sig. [2*(1-tailed Sig.)]	.234 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_2jam	Anestesi infiltrasi setelah insisi	8	9.00	72.00
	Anestesi infiltrasi sebelum insisi	8	8.00	64.00
	Total	16		

Test Statistics^a

	NRS_2jam
Mann-Whitney U	28.000
Wilcoxon W	64.000
Z	-.620
Asymp. Sig. (2-tailed)	.535
Exact Sig. [2*(1-tailed Sig.)]	.721 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_4jam	Kontrol	8	8.75	70.00
	Anestesi infiltrasi setelah insisi	8	8.25	66.00
	Total	16		

Test Statistics^a

	NRS_4jam
Mann-Whitney U	30.000
Wilcoxon W	66.000
Z	-.276
Asymp. Sig. (2-tailed)	.782
Exact Sig. [2*(1-tailed Sig.)]	.878 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_4jam	Kontrol	8	8.69	69.50
	Anestesi infiltrasi sebelum insisi	8	8.31	66.50
	Total	16		

Test Statistics^a

	NRS_4jam
Mann-Whitney U	30.500
Wilcoxon W	66.500
Z	-.207
Asymp. Sig. (2-tailed)	.836
Exact Sig. [2*(1-tailed Sig.)]	.878 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_4jam	Anestesi infiltrasi setelah insisi	8	8.38	67.00
	Anestesi infiltrasi sebelum insisi	8	8.63	69.00
	Total	16		

Test Statistics^a

	NRS_4jam
Mann-Whitney U	31.000
Wilcoxon W	67.000
Z	-.139
Asymp. Sig. (2-tailed)	.890
Exact Sig. [2*(1-tailed Sig.)]	.959 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_8jam	Kontrol	8	11.25	90.00
	Anestesi infiltrasi setelah insisi	8	5.75	46.00
	Total	16		

Test Statistics^a

	NRS_8jam
Mann-Whitney U	10.000
Wilcoxon W	46.000
Z	-2.441
Asymp. Sig. (2-tailed)	.015
Exact Sig. [2*(1-tailed Sig.)]	.021 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_8jam	Kontrol	8	11.25	90.00
	Anestesi infiltrasi sebelum insisi	8	5.75	46.00
	Total	16		

Test Statistics^a

	NRS_8jam
Mann-Whitney U	10.000
Wilcoxon W	46.000
Z	-2.441
Asymp. Sig. (2-tailed)	.015
Exact Sig. [2*(1-tailed Sig.)]	.021 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_8jam	Anestesi infiltrasi setelah insisi	8	8.50	68.00
	Anestesi infiltrasi sebelum insisi	8	8.50	68.00
	Total	16		

Test Statistics^a

	NRS_8jam
Mann-Whitney U	32.000
Wilcoxon W	68.000
Z	.000
Asymp. Sig. (2-tailed)	1.000
Exact Sig. [2*(1-tailed Sig.)]	1.000 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_16jam	Kontrol	8	9.69	77.50
	Anestesi infiltrasi setelah insisi	8	7.31	58.50
	Total	16		

Test Statistics^a

	NRS_16jam
Mann-Whitney U	22.500
Wilcoxon W	58.500
Z	-1.025
Asymp. Sig. (2-tailed)	.305
Exact Sig. [2*(1-tailed Sig.)]	.328 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_16jam	Kontrol	8	9.63	77.00
	Anestesi infiltrasi sebelum insisi	8	7.38	59.00
	Total	16		

Test Statistics^a

	NRS_16jam
Mann-Whitney U	23.000
Wilcoxon W	59.000
Z	-.969
Asymp. Sig. (2-tailed)	.333
Exact Sig. [2*(1-tailed Sig.)]	.382 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_16jam	Anestesi infiltrasi setelah insisi	8	8.69	69.50
	Anestesi infiltrasi sebelum insisi	8	8.31	66.50
	Total	16		

Test Statistics^a

	NRS 16jam
Mann-Whitney U	30.500
Wilcoxon W	66.500
Z	-.163
Asymp. Sig. (2-tailed)	.871
Exact Sig. [2*(1-tailed Sig.)]	.878 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_24jam	Kontrol	8	9.63	77.00
	Anestesi infiltrasi setelah insisi	8	7.38	59.00
	Total	16		

Test Statistics^a

	NRS 24jam
Mann-Whitney U	23.000
Wilcoxon W	59.000
Z	-1.098
Asymp. Sig. (2-tailed)	.272
Exact Sig. [2*(1-tailed Sig.)]	.382 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_24jam	Kontrol	8	9.31	74.50
	Anestesi infiltrasi sebelum insisi	8	7.69	61.50
	Total	16		

Test Statistics^a

	NRS 24jam
Mann-Whitney U	25.500
Wilcoxon W	61.500
Z	-.762
Asymp. Sig. (2-tailed)	.446
Exact Sig. [2*(1-tailed Sig.)]	.505 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

Ranks

	Kelompok	N	Mean Rank	Sum of Ranks
NRS_24jam	Anestesi infiltrasi setelah insisi	8	8.06	64.50
	Anestesi infiltrasi sebelum insisi	8	8.94	71.50
	Total	16		

Test Statistics^a

	NRS_24jam
Mann-Whitney U	28.500
Wilcoxon W	64.500
Z	-.449
Asymp. Sig. (2-tailed)	.653
Exact Sig. [2*(1-tailed Sig.)]	.721 ^b

a. Grouping Variable: Kelompok

b. Not corrected for ties.

UJI WILCOXON NRS KELOMPOK KONTROL

Test Statistics^a

	NRS_kontrol_1_4 - NRS_kontrol_1_2	NRS_kontrol_1_8 - NRS_kontrol_1_2	NRS_kontrol_16 - NRS_kontrol_2	NRS_kontrol_24 - NRS_kontrol_2	NRS_kontrol_1_8 - NRS_kontrol_1_4	NRS_kontrol_16 - NRS_kontrol_4	NRS_kontrol_24 - NRS_kontrol_4	NRS_kontrol_16 - NRS_kontrol_8	NRS_kontrol_24 - NRS_kontrol_8	NRS_kontrol_24 - NRS_kontrol_16
Z	1.342 ^b	-1.897 ^c	-1.492 ^c	.000 ^d	-2.388 ^c	-2.060 ^c	-1.134 ^c	-.707 ^b	-1.848 ^b	-1.761 ^b
Asym.p. Si g. (2-tail ed)	.180	.058	.136	1.000	.017	.039	.257	.480	.065	.078

a. Wilcoxon Signed Ranks Test

b. Based on positive ranks.

c. Based on negative ranks.

d. The sum of negative ranks equals the sum of positive ranks.

UJI WILCOXON NRS KELOMPOK SEBELUM

Test Statistics^a

	NRS_sebelum_m_4 - NRS_sebelum_m_2	NRS_sebelum_m_8 - NRS_sebelum_m_2	NRS_sebelum_16 - NRS_sebelum_2	NRS_sebelum_24 - NRS_sebelum_2	NRS_sebelum_m_8 - NRS_sebelum_m_4	NRS_sebelum_16 - NRS_sebelum_4	NRS_sebelum_24 - NRS_sebelum_4	NRS_sebelum_16 - NRS_sebelum_8	NRS_sebelum_24 - NRS_sebelum_8	NRS_sebelum_24 - NRS_sebelum_16
Z	1.342 ^b	-1.897 ^c	-1.492 ^c	.000 ^d	-2.388 ^c	-2.060 ^c	-1.134 ^c	-.707 ^b	-1.848 ^b	-1.761 ^b

Z	1.069 ^b	-.577 ^c	-1.511 ^b	-1.069 ^b	-1.342 ^c	-.954 ^b	.000 ^d	-2.060 ^b	-1.342 ^b	-.954 ^c
Asymp.	.285	.564	.131	.285	.180	.340	1.000	.039	.180	.340

a. Wilcoxon Signed Ranks Test

b. Based on negative ranks.

c. Based on positive ranks.

d. The sum of negative ranks equals the sum of positive ranks.

UJI WILCOXON NRS KELOMPOK SESUDAH

Test Statistics^a

	NRS_s esudah _4 - NRS_s esudah 2	NRS_s esudah _8 - NRS_s esudah 2	NRS_s esudah _16 - NRS_s esudah 2	NRS_s esudah _24 - NRS_s esudah 2	NRS_s esudah _8 - NRS_s esudah 4	NRS_s esudah _16 - NRS_s esudah 4	NRS_s esudah _24 - NRS_s esudah 4	NRS_s esudah _16 - NRS_s esudah 8	NRS_s esudah _24 - NRS_s esudah 8	NRS_s esudah _24 - NRS_s esudah 16
Z	-.816 ^b	1.761 ^b	-1.000 ^b	-1.512 ^b	1.476 ^b	-.378 ^c	-.756 ^b	-1.633 ^c	-.957 ^c	-1.242 ^b
Asymp.	.414	.078	.317	.131	.140	.705	.450	.102	.339	.214

a. Wilcoxon Signed Ranks Test

b. Based on negative ranks.

c. Based on positive ranks.

HASIL OLAH DATA RESCUE FENTANYL

Crosstab

	Kelompok 5					Total
	2 jam	4 jam	8 jam	16 jam	24 jam	
Rescue_tanpa	Count	8	8	6	3	8
	% within					
Rescue_tanpa		24.2%	24.2%	18.2%	9.1%	24.2%
	% within					
Kelompok_5		100.0%	100.0%	75.0%	37.5%	100.0%
	% of Total	20.0%	20.0%	15.0%	7.5%	20.0%
						82.5%
						82.5%

Ya	Count	0	0	2	5	0	7
	% within						
	Rescue_tanpa	0.0%	0.0%	28.6%	71.4%	0.0%	100.0%
	% within						
	Kelompok_5	0.0%	0.0%	25.0%	62.5%	0.0%	17.5%
	% of Total	0.0%	0.0%	5.0%	12.5%	0.0%	17.5%
Total	Count	8	8	8	8	8	40
	% within						
	Rescue_tanpa	20.0%	20.0%	20.0%	20.0%	20.0%	100.0%
	% within						
	Kelompok_5	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	% of Total	20.0%	20.0%	20.0%	20.0%	20.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	16.623 ^a	4	.002
Likelihood Ratio	17.516	4	.002
N of Valid Cases	40		

a. 5 cells (50.0%) have expected count less than 5. The minimum expected count is 1.40.

Crosstab

			Kelompok_5					Total
			2 jam	4 jam	8 jam	16 jam	24 jam	
Rescue_sebelumnya	Tidak	Count	8	7	8	7	7	37
		% within						
		Rescue_sebelumnya	21.6%	18.9%	21.6%	18.9%	18.9%	100.0%
		% within						
		Kelompok_5	100.0%	87.5%	100.0%	87.5%	87.5%	92.5%
		% of Total	20.0%	17.5%	20.0%	17.5%	17.5%	92.5%
Ya	Count	0	1	0	1	1	3	
	% within							
	Rescue_sebelumnya	0.0%	33.3%	0.0%	33.3%	33.3%	100.0%	
	% within							
	Kelompok_5	0.0%	12.5%	0.0%	12.5%	12.5%	7.5%	
	% of Total	0.0%	2.5%	0.0%	2.5%	2.5%	7.5%	
Total	Count	8	8	8	8	8	40	
	% within							
	Rescue_sebelumnya	20.0%	20.0%	20.0%	20.0%	20.0%	100.0%	
	% within							
	Kelompok_5	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
	% of Total	20.0%	20.0%	20.0%	20.0%	20.0%	100.0%	

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	2.162 ^a	4	.706
Likelihood Ratio	3.226	4	.521
N of Valid Cases	40		

a. 5 cells (50.0%) have expected count less than 5. The minimum expected count is .60.

Crosstab

		Kelompok 5					Total
		2 jam	4 jam	8 jam	16 jam	24 jam	
Rescue_setelah	Tidak Count	8	8	6	8	8	38
	% within Rescue_setelah	21.1%	21.1%	15.8%	21.1%	21.1%	100.0%
	% within Kelompok_5	100.0%	100.0%	75.0%	100.0%	100.0%	95.0%
	% of Total	20.0%	20.0%	15.0%	20.0%	20.0%	95.0%
Ya	Count	0	0	2	0	0	2
	% within Rescue_setelah	0.0%	0.0%	100.0%	0.0%	0.0%	100.0%
	% within Kelompok_5	0.0%	0.0%	25.0%	0.0%	0.0%	5.0%
	% of Total	0.0%	0.0%	5.0%	0.0%	0.0%	5.0%
Total	Count	8	8	8	8	8	40
	% within Rescue_setelah	20.0%	20.0%	20.0%	20.0%	20.0%	100.0%
	% within Kelompok_5	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	% of Total	20.0%	20.0%	20.0%	20.0%	20.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	8.421 ^a	4	.077
Likelihood Ratio	6.884	4	.142
N of Valid Cases	40		

a. 5 cells (50.0%) have expected count less than 5. The minimum expected count is .40.

Crosstab

		Kelompok			Total
		Tanpa	Sebelum	Sesudah	
Rescue2jam	Tidak Count	8	8	8	24
	% within Rescue2jam	33.3%	33.3%	33.3%	100.0%
	% within Kelompok	100.0%	100.0%	100.0%	100.0%

		% of Total	33.3%	33.3%	33.3%	100.0%
Total	Count	8	8	8	24	
	% within Rescue2jam	33.3%	33.3%	33.3%	100.0%	
	% within Kelompok	100.0%	100.0%	100.0%	100.0%	
	% of Total	33.3%	33.3%	33.3%	100.0%	

Chi-Square Tests

	Value
Pearson Chi-Square	^a .
N of Valid Cases	24

a. No statistics are computed because Rescue2jam is a constant.

Crosstab

			Kelompok			Total
			Tanpa	Sebelum	Sesudah	
Rescue4jam	Tidak	Count	8	7	8	23
		% within Rescue4jam	34.8%	30.4%	34.8%	100.0%
		% within Kelompok	100.0%	87.5%	100.0%	95.8%
		% of Total	33.3%	29.2%	33.3%	95.8%
	Ya	Count	0	1	0	1
		% within Rescue4jam	0.0%	100.0%	0.0%	100.0%
		% within Kelompok	0.0%	12.5%	0.0%	4.2%
		% of Total	0.0%	4.2%	0.0%	4.2%
Total		Count	8	8	8	24
		% within Rescue4jam	33.3%	33.3%	33.3%	100.0%
		% within Kelompok	100.0%	100.0%	100.0%	100.0%
		% of Total	33.3%	33.3%	33.3%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	2.087 ^a	2	.352
Likelihood Ratio	2.286	2	.319
N of Valid Cases	24		

a. 3 cells (50.0%) have expected count less than 5. The minimum expected count is .33.

Crosstab

			Kelompok			Total
			Tanpa	Sebelum	Sesudah	
Rescue8jam	Tidak	Count	6	8	6	20
		% within Rescue8jam	30.0%	40.0%	30.0%	100.0%
		% within Kelompok	75.0%	100.0%	75.0%	83.3%
		% of Total	25.0%	33.3%	25.0%	83.3%

Ya	Count	2	0	2	4
	% within Rescue8jam	50.0%	0.0%	50.0%	100.0%
	% within Kelompok	25.0%	0.0%	25.0%	16.7%
	% of Total	8.3%	0.0%	8.3%	16.7%
Total	Count	8	8	8	24
	% within Rescue8jam	33.3%	33.3%	33.3%	100.0%
	% within Kelompok	100.0%	100.0%	100.0%	100.0%
	% of Total	33.3%	33.3%	33.3%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	2.400 ^a	2	.301
Likelihood Ratio	3.632	2	.163
N of Valid Cases	24		

a. 3 cells (50.0%) have expected count less than 5. The minimum expected count is 1.33.

Crosstab

			Kelompok			Total
		Tanpa	Sebelum	Sesudah		
Rescue16jam	Tidak	Count	3	7	8	18
		% within Rescue16jam	16.7%	38.9%	44.4%	100.0%
		% within Kelompok	37.5%	87.5%	100.0%	75.0%
		% of Total	12.5%	29.2%	33.3%	75.0%
Ya		Count	5	1	0	6
		% within Rescue16jam	83.3%	16.7%	0.0%	100.0%
		% within Kelompok	62.5%	12.5%	0.0%	25.0%
		% of Total	20.8%	4.2%	0.0%	25.0%
Total		Count	8	8	8	24
		% within Rescue16jam	33.3%	33.3%	33.3%	100.0%
		% within Kelompok	100.0%	100.0%	100.0%	100.0%
		% of Total	33.3%	33.3%	33.3%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	9.333 ^a	2	.009
Likelihood Ratio	10.379	2	.006
N of Valid Cases	24		

a. 3 cells (50.0%) have expected count less than 5. The minimum expected count is 2.00.

Crosstab

			Kelompok			Total
			Tanpa	Sebelum	Sesudah	
Rescue24jam	Tidak	Count	8	7	8	23
		% within Rescue24jam	34.8%	30.4%	34.8%	100.0%
		% within Kelompok	100.0%	87.5%	100.0%	95.8%
		% of Total	33.3%	29.2%	33.3%	95.8%
	Ya	Count	0	1	0	1
		% within Rescue24jam	0.0%	100.0%	0.0%	100.0%
		% within Kelompok	0.0%	12.5%	0.0%	4.2%
		% of Total	0.0%	4.2%	0.0%	4.2%
Total	Count	8	8	8	24	
	% within Rescue24jam	33.3%	33.3%	33.3%	100.0%	
	% within Kelompok	100.0%	100.0%	100.0%	100.0%	
	% of Total	33.3%	33.3%	33.3%	100.0%	

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	2.087 ^a	2	.352
Likelihood Ratio	2.286	2	.319
N of Valid Cases	24		

a. 3 cells (50.0%) have expected count less than 5. The minimum expected count is .33.

HASIL OLAH DATA KADAR IL-6

Berdasarkan kelompok

Tests of Normality

	Kelompok	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Pre	Kontrol	.207	8	.200*	.919	8	.419
	Anestesi infiltrasi setelah insisi	.102	8	.200*	.995	8	1.000
	Anestesi infiltrasi sebelum insisi	.208	8	.200*	.857	8	.112
Empat_jam	Kontrol	.217	8	.200*	.919	8	.423
	Anestesi infiltrasi setelah insisi	.236	8	.200*	.939	8	.597
	Anestesi infiltrasi sebelum insisi	.250	8	.150	.886	8	.212
Delapan_jam	Kontrol	.227	8	.200*	.887	8	.220
	Anestesi infiltrasi setelah insisi	.179	8	.200*	.915	8	.393
	Anestesi infiltrasi sebelum insisi	.250	8	.150	.867	8	.140

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Descriptives

		N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimu m	Maximu m
						Lower Bound	Upper Bound		
Pre	Kontrol	8	52.1250	17.19998	6.08111	37.7455	66.5045	29.00	78.00
	Anestesi infiltrasi setelah insisi	8	46.8750	13.51652	4.77881	35.5749	58.1751	26.00	69.00
	Anestesi infiltrasi sebelum insisi	8	28.1250	11.50699	4.06833	18.5049	37.7451	12.00	40.00
	Total	24	42.3750	17.22438	3.51591	35.1018	49.6482	12.00	78.00
Empat_jam	Kontrol	8	67.5000	18.17376	6.42540	52.3064	82.6936	44.00	98.00
	Anestesi infiltrasi setelah insisi	8	44.2500	12.27948	4.34145	33.9841	54.5159	24.00	60.00
	Anestesi infiltrasi sebelum insisi	8	17.0000	3.70328	1.30931	13.9040	20.0960	11.00	22.00
	Total	24	42.9167	24.39336	4.97927	32.6163	53.2171	11.00	98.00
Delapan_jam	Kontrol	8	40.0000	14.91883	5.27460	27.5275	52.4725	22.00	60.00
	Anestesi infiltrasi setelah insisi	8	24.8750	6.72814	2.37876	19.2501	30.4999	14.00	32.00
	Anestesi infiltrasi sebelum insisi	8	14.0000	5.34522	1.88982	9.5313	18.4687	4.00	20.00

Total	24	26.291	7	14.4507	5	2.9497	5	20.189	6	32.393	7	4.00		60.00
-------	----	--------	---	---------	---	--------	---	--------	---	--------	---	------	--	-------

Test of Homogeneity of Variances

	Levene Statistic	df1	df2	Sig.
Pre	.344	2	21	.713
Empat_jam	4.832	2	21	.019
Delapan_jam	8.991	2	21	.002

ANOVA

		Sum of Squares	df	Mean Square	F	Sig.
Pre	Between Groups	2547.000	2	1273.500	6.253	.007
	Within Groups	4276.625	21	203.649		
	Total	6823.625	23			
Empat_jam	Between Groups	10222.333	2	5111.167	30.990	.000
	Within Groups	3463.500	21	164.929		
	Total	13685.833	23			
Delapan_jam	Between Groups	2728.083	2	1364.042	13.806	.000
	Within Groups	2074.875	21	98.804		
	Total	4802.958	23			

Multiple Comparisons

Tukey HSD

Dependent Variable	(I) Kelompok	(J) Kelompok	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
						Lower Bound	Upper Bound
Pre	Kontrol	Anestesi infiltrasi setelah insisi	5.25000	7.13528	.745	-12.7350	23.2350
		Anestesi infiltrasi sebelum insisi	24.00000*	7.13528	.008	6.0150	41.9850
		Kontrol	-5.25000	7.13528	.745	-23.2350	12.7350
	Anestesi infiltrasi setelah insisi	Anestesi infiltrasi sebelum insisi	18.75000*	7.13528	.040	.7650	36.7350
		Kontrol	24.00000*	7.13528	.008	-41.9850	-6.0150
		Anestesi infiltrasi setelah insisi	18.75000*	7.13528	.040	-36.7350	-.7650
Empat_jam	Kontrol	Anestesi infiltrasi setelah insisi	23.25000*	6.42123	.004	7.0648	39.4352

	Anestesi infiltrasi sebelum insisi	50.50000*	6.42123	.000	34.3148	66.6852
Anestesi infiltrasi setelah insisi	Kontrol	23.25000*	6.42123	.004	39.4352	-7.0648
	Anestesi infiltrasi sebelum insisi	27.25000*	6.42123	.001	11.0648	43.4352
Anestesi infiltrasi sebelum insisi	Kontrol	50.50000*	6.42123	.000	66.6852	34.3148
	Anestesi infiltrasi setelah insisi	27.25000*	6.42123	.001	43.4352	11.0648
Delapan_jam	Kontrol	Anestesi infiltrasi setelah insisi	15.12500*	4.97000	.016	2.5978
		Anestesi infiltrasi sebelum insisi	26.00000*	4.97000	.000	13.4728
	Anestesi infiltrasi setelah insisi	Kontrol	15.12500*	4.97000	.016	27.6522
		Anestesi infiltrasi sebelum insisi	10.87500	4.97000	.097	-1.6522
	Anestesi infiltrasi sebelum insisi	Kontrol	26.00000*	4.97000	.000	38.5272
		Anestesi infiltrasi setelah insisi	-10.87500	4.97000	.097	23.4022

*. The mean difference is significant at the 0.05 level.

Berdasarkan waktu pengukuran

Tests of Normality

	Kelompok	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Tanpa	Pre SC	.207	8	.200*	.919	8	.419
	4 jam pasca SC	.217	8	.200*	.919	8	.423
	8 jam pasca SC	.227	8	.200*	.887	8	.220
Sesudah	Pre SC	.102	8	.200*	.995	8	1.000
	4 jam pasca SC	.236	8	.200*	.939	8	.597
	8 jam pasca SC	.179	8	.200*	.915	8	.393
Sebelum	Pre SC	.208	8	.200*	.857	8	.112
	4 jam pasca SC	.250	8	.150	.886	8	.212
	8 jam pasca SC	.250	8	.150	.867	8	.140

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Descriptives

		N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
						Lower Bound	Upper Bound		
Tanpa	Pre SC	8	52.1250	17.19998	6.08111	37.7455	66.5045	29.00	78.00
	4 jam pasca SC	8	67.5000	18.17376	6.42540	52.3064	82.6936	44.00	98.00
	8 jam pasca SC	8	40.0000	14.91883	5.27460	27.5275	52.4725	22.00	60.00
	Total	24	53.2083	19.75938	4.03337	44.8647	61.5520	22.00	98.00
	Sesudah	Pre SC	8	46.8750	13.51652	4.77881	35.5749	58.1751	26.00
Sebelum	4 jam pasca SC	8	44.2500	12.27948	4.34145	33.9841	54.5159	24.00	60.00
	8 jam pasca SC	8	24.8750	6.72814	2.37876	19.2501	30.4999	14.00	32.00
	Total	24	38.6667	14.68707	2.99799	32.4649	44.8685	14.00	69.00
	Pre SC	8	28.1250	11.50699	4.06833	18.5049	37.7451	12.00	40.00
	4 jam pasca SC	8	17.0000	3.70328	1.30931	13.9040	20.0960	11.00	22.00
Pair 3	8 jam pasca SC	8	14.0000	5.34522	1.88982	9.5313	18.4687	4.00	20.00
	Total	24	19.7083	9.57569	1.95463	15.6649	23.7518	4.00	40.00

UJI PAIRED T TEST IL-6 KELOMPOK KONTROL**Paired Samples Statistics**

		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	Kontrol_pre	52.1250	8	17.19998	6.08111
	Kontrol_4	67.5000	8	18.17376	6.42540
Pair 2	Kontrol_pre	52.1250	8	17.19998	6.08111
	Kontrol_8	40.0000	8	14.91883	5.27460
Pair 3	Kontrol_4	67.5000	8	18.17376	6.42540
	Kontrol_8	40.0000	8	14.91883	5.27460

Paired Samples Correlations

		N	Correlation	Sig.
Pair 1	Kontrol_pre & Kontrol_4	8	.069	.871
Pair 2	Kontrol_pre & Kontrol_8	8	-.473	.237
Pair 3	Kontrol_4 & Kontrol_8	8	.393	.336

Paired Samples Test

	Paired Differences					t	df	Sig. (2-tailed)			
	Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference							
				Lower	Upper						
Pair 1 Kontrol_pre - 1 Kontrol_4	15.37500	24.14206	8.53551	35.55827	4.80827	1.801	7	.115			
Pair 2 Kontrol_pre - 2 Kontrol_8	12.12500	27.58590	9.75309	10.93739	35.18739	1.243	7	.254			
Pair 3 Kontrol_4 - 3 Kontrol_8	27.50000	18.43909	6.51920	12.08454	42.91546	4.218	7	.004			

UJI PAIRED T TEST IL-6 KELOMPOK SEBELUM INSISI

Paired Samples Statistics

		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	Sebelum_pre	28.1250	8	11.50699	4.06833
	Sebelum_4	17.0000	8	3.70328	1.30931
Pair 2	Sebelum_pre	28.1250	8	11.50699	4.06833
	Sebelum_8	14.0000	8	5.34522	1.88982
Pair 3	Sebelum_4	17.0000	8	3.70328	1.30931
	Sebelum_8	14.0000	8	5.34522	1.88982

Paired Samples Correlations

		N	Correlation	Sig.
Pair 1	Sebelum_pre & Sebelum_4	8	.288	.489
Pair 2	Sebelum_pre & Sebelum_8	8	.323	.435
Pair 3	Sebelum_4 & Sebelum_8	8	.909	.002

Paired Samples Test

	Paired Differences					t	df	Sig. (2-tailed)			
	Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference							
				Lower	Upper						
Pair 1 Sebelum_pre - 1 Sebelum_4	11.12500	11.02513	3.89797	1.90776	20.34224	2.854	7	.025			
Pair 2 Sebelum_pre - 2 Sebelum_8	14.12500	11.01217	3.89339	4.91860	23.33140	3.628	7	.008			
Pair 3 Sebelum_4 - 3 Sebelum_8	3.00000	2.50713	.88641	.90398	5.09602	3.384	7	.012			

UJI PAIRED T TEST IL-6 KELOMPOK SETELAH INSISI

Paired Samples Statistics

		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	Sesudah_pre	46.8750	8	13.51652	4.77881
	Sesudah_4	44.2500	8	12.27948	4.34145
Pair 2	Sesudah_pre	46.8750	8	13.51652	4.77881
	Sesudah_8	24.8750	8	6.72814	2.37876

Pair 3	Sesudah_4	44.2500	8	12.27948	4.34145
	Sesudah_8	24.8750	8	6.72814	2.37876

Paired Samples Correlations

		N	Correlation	Sig.
Pair 1	Sesudah_pre & Sesudah_4	8	.385	.346
Pair 2	Sesudah_pre & Sesudah_8	8	.217	.606
Pair 3	Sesudah_4 & Sesudah_8	8	-.435	.281

Paired Samples Test

	Paired Differences					t	df	Sig. (2-tailed)			
	Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference							
				Lower	Upper						
Pair 1 Sesudah_pre - 1 Sesudah_4	2.62500	14.34212	5.07071	-9.36531	14.61531	.518	7	.621			
Pair 2 Sesudah_pre - 2 Sesudah_8	22.00000	13.73213	4.85504	10.51965	33.48035	4.531	7	.003			
Pair 3 Sesudah_4 - 3 Sesudah_8	19.37500	16.37016	5.78773	5.68920	33.06080	3.348	7	.012			