

Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-Cov-2) A Symptomatic Infection in Pediatric Population: A Literature Review

Harun Achmad¹, Nur Afni Puspita²

¹Pediatric Dentistry Department, Faculty of Dentistry, Hasanuddin University, Makassar, Indonesia

²Dental Intern, Faculty of Dentistry, Hasanuddin University, Makassar, Indonesia
Corresponding e-mail:harunachmader@gmail.com

ABSTRACT

Introduction: *Severe Acute Respiratory Syndrome Coronavirus-2* (SARS-Cov-2) is a zoonotic pathogen that causes COVID-19 disease. This disease manifests with nonspecific symptoms similar to flu, the most common clinical complication is acute respiratory distress, which is the main cause of death. SARS-Cov-2 rarely causes severe illness in children. The database obtained comes from the Centers for Disease Control and Prevention, SARS-Cov-2 only occurs in 1.7% of people younger than 4 years and 7.7% of those aged 5 to 17 years. **Objective:** This article aims to review the basic issues regarding asymptomatic SARS-Cov-2 infection in children. **Method:** The search was carried out using databases from Google Scholar and Pubmed. The included articles were articles published from 2020 to 2021. The keywords and terms used for the article search are as follows: -SARS-Cov-2l, -SARS-Cov-2 Asymptomatic Infectionl, SARS-Cov-2 Asymptomatic Infection in Pediatricl. **Results:** There were 13 articles that discussed things related to asymptomatic SARS-Cov-2 infection in children. **Conclusion:** All age groups of children are susceptible to infection with SARS-CoV-2. Infection in children usually has milder clinical signs and symptoms than in adults and is usually asymptomatic.

Keywords: SARS-Cov-2 asymptomatic infection, Pediatric.

INTRODUCTION

Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-Cov-2) is a zoonotic pathogen that causes COVID-19 disease. This disease manifests with nonspecific symptoms similar to flu, the most common clinical complication is acute respiratory distress, which is the main cause of death.¹SARS-Cov-2 rarely causes severe illness in children, the severity of SARS-Cov-2 infection is more common in adults.^{2,3}The database obtained comes from the Centers for Disease. Control and Prevention, SARS-Cov-2 occurred in only 1.7% of people younger than 4 years and 7.7% of those aged 5 to 17 years.⁴ Patients who require medical care may have comorbid conditions that can affect their SARS-Cov-2 infection.⁵ The asymptomatic group reported an increase of 0.6% to 13.7%. 6–10 Most of these studies included adults only. Three studies examining asymptomatic pediatric surgical oncological patients through Polymerase Chain Reaction (PCR) reported positive results for SARS-Cov-2, namely 0.6% to 2.5%.⁶⁻⁸ In addition a research involving 33.041 children tested in various hospitals in the

United States showed asymptomatic results with a prevalence of 0.7%.¹¹ Although children are classified as more benign or remain asymptomatic when compared to adults, they have the potential to carry the virus and become a source of infection.¹²

SARS-Cov-2 is present in high concentrations in the oral cavity and pharynx.¹³ In addition, procedures in dentistry can produce aerosols that increase the risk of transmission.¹⁴ Thus, dental and oral health care providers are at high risk of infection if they handle SARS-Cov positive patients. -2. The Center for Disease Control (CDC) and the American Dental Association (ADA) have developed infection control guidelines in dental practice during the COVID-19 pandemic.¹⁵ These include screening each patient for signs and symptoms of COVID-19, use of universal personal protective equipment and sources of control strategies such as wearing face masks at all times, hand hygiene and implementing technical approaches to reduce disease transmission. The CDC recommends asymptomatic SARS-Cov-2 testing for dental patients undergoing dental and oral health care as a way to identify operators and reduce risks in dental and oral health care facilities. This new strategy, however, has not been used universally. Identification of patients with SARS-Cov-2 is important in infection control strategies and reduces the risk of infection in oral health care.

METHOD

Search Strategy

The search was carried out using online databases from Google Scholar and Pubmed. The included articles were articles published from 2020 to 2021. The keywords and terms used for the article search are as follows: -SARS-Cov-2 -SARS-Cov-2 A symptomatic Infectional -SARS-Cov-2 Asymptomatic Infection in Pediatricl. In each article found according to keywords, the title and abstract of each article were checked and articles that met the criteria were downloaded.

The flow chart in figure 1 identifies the articles included and excluded at each stage. A total of 74 articles were found using the above keywords, after removing duplicates, 74 articles were found. A total of 74 articles were screened, and 61 articles were excluded, 13 full-text articles were examined for their eligibility and 13 full-text articles were included as articles to be reviewed

Inclusion and Exclusion Criteria


1. Inclusion Criteria

- a. Articles published in the last 2 years (2020-2021)
- b. Articles in English
- c. Articles that focus on the definition, prevalence, classification, etiology, signs and symptoms, effects, and management of asymptomatic SARS-Cov-2 infection in children.

2. Exclusion Criteria

- a. Articles that only have abstracts written in English
- b. Full-text articles unavailable for open access
- c. The article does not focus on asymptomatic SARS-Cov-2 infection in children, only touches slightly on asymptomatic SARS-Cov-2 infection in children.

Figure1.PRISMA flow chart of the systemic literature review and article identification process.



RESULTS

Of the articles found, 13 were analyzed and included in this systematic review. Information regarding the article can be found in table 1.

No.	Writer	Year	Title	Conclusion
1.	A Isohime F, Temsah M, Al-Nemri AM, Somily AM, Al- Subaie S. ¹⁶	2020	COVID-19 Infection prevalence in pediatric population: Etiology, clinical presentation, and outcome.	Currently, the COVID-19 pandemic is developing rapidly. As of April 14, 2020, cases reported in the literature showed that, most of the pediatric COVID-19 patients have a good prognosis, and in mild cases, recovery takes 7 to 14 days after onset. 17 In an extensive analysis of 72,314 cases in China, there were 549 cases of single death, in the 10–19-year age group.18 In Spain, 60% of COVID-19 infections occurred in children, 10% were admitted to pediatric intensive care units.19 In the United States, 5.7% - 20 % of children were hospitalized with a percentage of 0.58% - 2% needing care in the pediatric intensive care unit. ²⁰
2.	Kronbichler A, Kresse D, Yoon S, Lee KH, Effenberger M, ShinJI.	2020	Asymptomatic patients as a source of COVID-19 Infections: A systematic review and meta- analysis.	More than half of the asymptomatic patients presented with abnormalities on CT scans. Asymptomatic patients are contagious and a potential source of infection. In this systematic review and meta-analysis, it can be proved that a CT scan is not necessary for young patients who present asymptotically because most of them will not show any relevant abnormalities.

				In this analysis it was also found that asymptomatic cases can cause more than 20% of the total COVID-19 patients
3.	Yoon S, Li H, Lee KH, Hong SH, Ki M D, Im H, Rah W, Kim E, Cha S, Yang J, Kronbicher A, Kresse D, Koyanagi A, Jacob L, Ghayda RA, Shin Ji, Smith L. ²¹	2020	Clinical characteristic of asymptomatic and symptomatic Pediatric coronavirus Disease 2019 (COVID-19): A systematic review	CT scan results of children suffering from COVID-19 did not show any relevant results compared to clinical findings, nor was there a significant difference in clinical findings between asymptomatic and symptomatic children. Further studies evaluating COVID-19 in pediatric could contribute to potential therapeutic interventions and prevention strategies to limit its spread.
4.	Consiglio CR, Nicola C, Sardh F, Landegren N, Palma P, Brodin P. ²²	2020	The Immunology of Multi system Inflammatory Syndrome in Children with COVID-19	Immune system, blood, cytokines and autoantibodies in healthy children, children with a history of Kawasaki disease, children infected with SARS-CoV-2 and children presenting with MIS-C. It was found that the inflammatory response in MIS-C is different from that of the cytokines in severe acute COVID-19.
5.	Cui X, Zhao Z, Zhang T, Guo W, Guo W, Zheng J, Zhang J, Dong C, Na R, Zheng L, Li W, Li Z, Ma J, Wang J, He S, Xu Y, Si P, Shen Y, Cai C. ²³	2021	A Systematic review and meta-analysis of children with coronavirus disease 2019 (COVID-19).	The COVID-19 pandemic can strike all age groups of children with milder symptoms. Pediatric patients suspected of having COVID-19 have non-specific signs and symptoms, which can include fever and cough. Pediatric patients with COVID-19 may experience milder signs and symptoms compared to atypical clinical manifestations and rare lymphopenia. The high incidence of vomiting and symptoms of vomiting gives more attention to children under 1 year of age. The characteristics of COVID-19 in children and adults are different, therefore further research is needed.

6.	Ho CLT, Oligbu P, Ojubolamo O, Pervaiz M, Oligbu G. ²⁴	2020	Clinical characteristic of children with COVID-19.	The data currently available suggests that children who catch COVID-19 are more likely to experience milder symptoms than adults. This study identifies the international standard of COVID-19 cases in children in order to better understand the development and possible complications associated with this virus. In addition, given the absence of adequate treatment, prospective randomized controlled trials would be useful to provide strong evidence for the development of treatment strategies, hopefully reducing morbidity in children.
7.	De Luca CD, Esposito E, Cristiani L, Mancino E, Nenna R, Cortis E, Midulla F. ²⁵	2020	Covid-19 in children: A brief overview after three months experience.	COVID-19 shows a milder clinical history in children than in adults. A much lower percentage of children suffering from severe or critical illness and death is an exception. Children may present with non-specific viral infection symptoms which demonstrate the importance of accurate differential diagnosis with typical pediatric clinical conditions such as upper respiratory tract infections, fever of unknown cause, viral or bacterial pneumonia, bronchiolitis, gastroenteritis and asthma attacks. ²⁶ Same as in adults, in children cardiovascular disease also appears to be the comorbid disease that most often causes the severity of COVID-19.

8.	Ludvigsson, JF. ²⁷	2020	Children are unlikely to be the main drivers of the COVID-19 pandemic—a systematic review.	The conclusion of this systematic review is that children are unlikely to be the main drivers of the pandemic. Opening schools and kindergartens is unlikely to affect the death rate for COVID-19 in parents.
9.	Rabinowicz S, Leshem A, Pessach IM. ²⁸	2020	COVID-19 in The Pediatric Population – Review and current evidence.	Children in any age group can be infected with SARS-CoV-2, with a lower frequency and severity than adults, although further epidemiological data are needed. Data on antiviral care, vaccination safety and immunogenicity and better specification of high-risk patients in the pediatric population are also needed. As the pandemic continues to evolve, it is still difficult to assess the long-term effects that significant changes will have on society, the economy and human behavior on the health and well-being of children in the future.
10	Ciuca IM. ²⁹	2020	COVID-19 in Children: An ample review.	The study emphasizes that although COVID-19 is rare in children, asymptomatic SARS-CoV-2 infection is more common in children than in adults. Collecting high-quality evidence is important to better understand COVID-19 in children, as well as a way to determine the most efficient case management strategy.
11	Paglia L. ³⁰	2020	COVID-19 and Pediatric Dentistry after lock down	A common clinical observation is that COVID-19 is less severe in children, and in this group infection usually occurs asymptotically.

				<p>Further clinical studies can clarify infection and transmission dynamics; therefore, it is important to apply to children all the preventive measures not hygiene measures recommended by health authorities during dental care. Dentists should avoid aerosol-generating procedures as much as possible, minimizing the use of syringes. If possible, it is recommended to use minimally invasive procedures and a traumatic restorative treatment.</p>
12	<p>Serrano CO, Alonso E, Andres M, Buitrago NM, Vigara AP, Pajares MP, LopezA C, Moll GG, Espin IM, BarriocanalMB.³¹</p>	2020	<p>Pediatric chest x-ray in covid-19 infection.</p>	<p>This study proves that children with symptomatic COVID-19 symptoms show abnormalities in CXR results. The findings are not specific and therefore the CXR cannot be used as the main diagnostic tool. However, radiographs should be considered for use in these patients.</p>
13	<p>Badal S, Bajgain KT, Badal S, Thapa R, Bajgain BB, Santana MJ.³²</p>	2020	<p>Prevalence, clinical characteristic, and outcomes of pediatric COVID-19: A Systematic review and meta-analysis</p>	<p>This study shows that all age groups of children are susceptible to infection with COVID-19, usually having mild clinical signs and symptoms. Critical circumstances or death are very rare. Cough fever and headache are the most common symptoms, whereas laboratory or radiological results do not show consistency, making it impossible to determine a diagnosis. Overall, the prognosis for COVID-19 in the pediatric age group is</p>

				<p>good. However, the differences in the characteristics of clinical findings in children suspected of having COVID-19 have implications for uncontrolled transmission and control of viral infection.</p> <p>Therefore, proper guidelines for testing and quarantine in children are needed. Effective strategies to ensure the prevention and maintenance of contacts are highly recommended for proper management of children during the time of this pandemic.</p>
--	--	--	--	--

DISCUSSION

The rapid and continuous spread of SARS-CoV-2 infection indirectly forces practitioners and health care providers to have a clear picture of the mode of transmission and clinical characteristics of the infection. The pediatric population has recently been examined by various systematic reviews,³³⁻³⁵ this is done in order to find out more about the clinical characteristics of infection in children and to evaluate the extent to which children with asymptomatic infections can act as carriers of symptomatic disease that are mild, despite mild symptoms, SARS-CoV-2 infection can still be transmitted rapidly during the incubation period.

Diagnostic markers are more accurate in determining symptom onset in the population of children under 10 years of age. In this group, abnormal laboratory characteristics, in particular high lymphocyte levels, were associated with symptomatic infections, and radiographic findings that were abnormal, low white blood cell count, low neutrophil count and low creatinine level.^{36,37,38} Previous studies have shown that pediatric patients both boys and girls have the same likelihood of becoming infected, but children who are male sex are more likely to experience asymptomatic infections.

All age groups of children are susceptible to infection with COVID-19. Acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in children is usually very mild and asymptomatic. Critical circumstances or death are very rare. The data currently available suggests that children who catch COVID-19 are more likely to experience milder symptoms than adults. Cough fever and headache are the most common symptoms, while laboratory or radiological results are not consistent, so it is not easy to determine the diagnosis. Overall, the prognosis for COVID-19 in the pediatric age group is good. However, the differences in the characteristics of clinical findings in children suspected of having COVID-19 have implications for uncontrolled transmission and control of viral infection.

Most of the patients with asymptomatic symptoms show normal laboratory results. The predominant laboratory findings were leukopenia, lymphopenia, LDH and elevated CRP. Existing studies of symptomatic patients showed laboratory findings with leukocytosis rather than leukopenia^{39,40} whereas CT scan results of 135 asymptomatic patients showed abnormal results. Asymptomatic patients especially in children showed significantly normal CT scan results. Several studies have objected to the use of CT scanning in patients in the age group under 20 years with normal chest radiographs.^{41,42,43,44,45,46} CT scans need not be performed on young patients who present asymptotically because most of them will not show any

relevant abnormalities. CT scan results in children suffering from COVID-19 did not show any relevant results compared to clinical findings, nor was there a significant difference in clinical findings between asymptomatic and symptomatic children.

CONCLUSION

All age groups of children are susceptible to infection with SARS-CoV-2. Infection in children usually has milder clinical signs and symptoms than in adults and is usually asymptomatic. Critical circumstances or death are very rare. Overall, the prognosis for COVID-19 in the pediatric age group is good. However, the differences in the characteristics of clinical findings in children suspected of having COVID-19 have implications for uncontrolled transmission and control of viral infection. Therefore, proper guidelines for testing and quarantine in children are needed. Effective strategies to ensure the prevention and maintenance of contact are highly recommended for the proper management of children in this time of the pandemic.

REFERENCES

1. Vincent J-L, Taccone FS. Understanding pathways to death in patients with COVID-19. *Lancet Respir Med.* 2020;8(5):430–2.
2. Milani GP, Bottino I, Rocchi A. Frequency of Children Adults Carrying Severe Acute Respiratory Syndrome Coronavirus 2 Asymptotically. *JAMA Pediatr.* 2020;175(2).
3. Prevention. C for D Cand. COVID View: Key updates for week 39 ending. 2020;
4. COVID-19 cases, deaths and trends in the US. Centers for Disease Control and Prevention. 2020.
5. Shekerdemian LS, Mahmood NR, Wolfe KK. Characteristics and Outcomes of Children with Coronavirus Disease 2019 (COVID-19) Infection Admitted to US and Canadian Pediatric Intensive Care Units. *JAMA Pediatr.* 2020; 174(9):868–73.
6. Boulad F, Kamboj Mi, Bouvier N. COVID-19 in Children with Cancer in New York City. *JAMA on col.* 2020; 6(9):1459–60.
7. Lin EE, Blumberg TJ, Adler AC. Incidence of COVID-19 in Pediatric Surgical Patients among 3 US Children’s Hospital. *JAMA Surg.* 2020; 155(8):775–7.
8. Shah MA, Mayer S, Emlen F. Clinical Screening for COVID-19 in Asymptomatic Patients with Cancer. *JAMA Netw Open.* 2020; 3(9).
9. Blumberg TJ, Adler AC, Lin EE, Fazal FZ. Universal Screening for COVID-19 Children Undergoing Orthopedic Surgery: A Multicenter Report. *J Pediatr Orthop.* 2020; 40(10):990–3.
10. Sutton D, Fuchs K, D’Alton M, Goffman D. Universal Screening for SARs-Cov-2 in Women Admitted for Delivery. *N Engl JMed.* 2020; 382:2163–4.
11. Sola A, David A, Roshe K, Baha A, Ramirez-Avila L, Chan D. Prevalence of SARS-Cov-2 infection in children without symptoms of coronavirus disease 2019. *JAMA Pediatr.* 2020; 227:45-52.e5.

12. Yonker L, Neilan AM, Bartsch Y, Patel AB. Pediatric Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2): Clinical Presentation, Infectivity, and Immune Responses. *JPediatr.* 2020; 227:45–52.
13. Centers for Diseases Control and Prevention COVID-19: guidance for dental settings.
14. Harrel SK, Molinari J. Aerosols and splatter in dentistry: A brief review of the literature and infection control implications. *J Am Dent Assoc.*2004; 135(4):429–37.
15. ADA Interim Guidance for Minimizing Risk of COVID-19 Transmission.2020.
16. Alsohime F, Temsah M, Al-Nemri A, Somily A, Al-Subaie S. COVID-19 infection prevalence in pediatric population: Etiology, clinical presentation, and outcome. *J Infect Public Health.* 2020; 13(2020):1791–6.
17. Feldstein L, Rose E, Horwitz S, Collins J. Overcoming COVID-19 Investigators and the CDC COVID-19 Response Team. Multisystem inflammatory syndrome in U.S Children and adolescents. *N Engl JMed.*2020; 383:334–46.
18. Prevention CC for DC and. The Novel Corona Virus Pneumonia Emergency Response Epidemiology Team. Vital Surveillances: The Epidemiological Characteristics of an Outbreak of 2019 Novel Corona virus Diseases (COVID-19). *China CDC Wkly.* 2020; 2(8):113–22.
19. Tagarro A, Epalza C, Santos M, Sanz-Santaefemia F, Otheo E, Moraleda C. Screening and Severity of Coronavirus Disease 2019 (COVID-19) in Children in Madrid, Spain. *JAMA Pediatr.* 2021; 175(3):316–7.
20. Prevention. C for DC and. Coronavirus Disease 2019 in Children-United States. *Morb Mortal Wkly Rep.* 2020; 69(14):422–6.
21. YoonS, LiH, Lee K, Hong, SH Kim D, Im H, Rah W, et al. Clinical characteristic of asymptomatic and symptomatic pediatric coronavirus disease 2019 (COVID-19): Asystematic review. *Medicina (B Aires).* 2020; 56:474.
22. Consiglio C, Cotugno N, Sardh F, Landegren N, Palma P, Brodin P. The Immunology of Multisystem Inflammatory Syndrome in Children with COVID-19. *Cell Press.*2020; 183:968–81.
23. Cui X, Zhao Z, Zhang T, Guo W, Guo W, Zheng J, et al. A systematic review and meta- analysis of children with coronavirus disease 2019 (COVID-19) [Internet]. Vol.93, *Journal of Medical Virology.* John Wiley & Sons, Ltd; 2021. 1057–1069 p. Available from: <http://dx.doi.org/10.1002/jmv.26398>
24. Lok Tung Ho C, Oligbu P, Ojubolamo O, Pervaiz M, Oligbu G. Clinical Characteristics of Children with COVID-19. *AIMS Public Heal.*2020; 7(2):258–73.
25. Dolores C, Luca D, Esposito E, Cristiani L, Mancino E, Nenna R, et al. Covid-19 in Children: A brief overview after three months experience. *Pediatr Respir Rev.*2020; 35(2020):9–14.
26. Alfredo T, Cristina E, Mar S, Jose S, Enrique O, Cinta M. Screening and severity of corona virus disease 2019 (COVID-19) in children in Madrid, Spain. *JAMA Pediatr.*2020; 9–10.
27. Ludvigsson JF. Children are unlikely to be the main drivers of the COVID-19pandemic—Asystematicreview. *ActaPaediatrIntJPediatr.*2020; 109(8):1525–30.
28. Rabinowicz S, Leshem E, Pessach IM. COVID-19 in the Pediatric Population—Review and Current Evidence. *Curr InfectDisRep.*2020; 22(11).
29. Ciuca IM. COVID-19 in children: An ample review. *Risk Manag Healthc Policy.*2020; 13:661–9.
30. Paglia L. COVID- 19 and pediatric dentistry after the lock down. *Eur J Pediatr Dent.* 2020; 21(2):89.
31. Serrano C, Alonso E, Andres M, Buitrago N, Vigara A, Pajares M, et al. Pediatric chest x-ray incovid-19 infection. *Eur J Radiol.* 2020; 131.
32. Badal S, Thapa K,Badal S, ThapaR, Bajgain BB. Prevalence, clinical characteristics, <http://annalsofrscb.ro>

- and outcomes of pediatric COVID-19: A systematic review and meta-analysis. *J Clin Virol.* 2020; 135(2021):104715.
33. Castagnoli R, Votto M, Licari A, Brambilla I, Bruno R, Perlini S, et al. Severe Acute Respiratory Syndrome Coronavirus2 (SARS-CoV-2) Infection in Children and Adolescents A Systematic Review. *JAMA Pediatr.*2020; 174(9):882–9.
 34. Mustafa N, Selim L. Characterisation of COVID-19 Pandemic in Paediatric Age Group: A Systematic Review and Meta-Analysis. *J Clin Virol.*2020; 128(104395).
 35. Ludvigsson J. Systematic review of COVID-19 in children shows milder cases and a better prognosis than adults. *J Acta Paediatr.* 2020; 109(6):1088–95.
 36. Huang L, Zhang X, Zhang X, Xu Y, Zhang C, Xu A. Rapid asymptomatic transmission of COVID-19 during the incubation period demonstrating strong infectivity in a cluster of youngsters aged 16-23 years outside Wuhan and characteristics of young patients with COVID-19: A prospective contact-tracing study. *J Infect.* 2020; 80(6): e1–3.
 37. Tian W, Jiang W, Yao J, Nicholson C, Li R, Sigurslid H, et al. Predictors of mortality in hospitalized COVID-19 patients: A systematic review and meta-analysis. *J Med Virol.* 2020; 92(10):1875–83.
 38. Shi S, Qin M, Shen B, Cai Y, Liu T, Yang F, et al. Association of Cardiac Injury with Mortality in Hospitalized Patients With COVID-19 in Wuhan, China. *JAMA Cardiol.*2020; 5(7):802–10.
 39. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet.* 2020; 395(10223):507–13.
 40. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y. Clinical features of patients infected with 2019 novel corona virus in Wuhan, China. *Lancet.*2020; 395(10223):497–506.
 41. Inui S, Fujikawa A, Jitsu M, Kunishima N, Watanabe S, Suzuki Y, et al. Chest CT Findings in Cases from the Cruise Ship Diamond Princess with Corona virus Disease (COVID-19). *Radiol Cardiothoracic Imaging.*2020; 2(2).
 42. Achmad H, Gani A, Djais AI, Hatta LI, Rieuwpassa IE, Monry A. Effectiveness of Edible Film Chitosan from Waste White Shrimp (*Litopenaeus vannamei*) in Reducing colonization of *Porphyromonas Gingivalis* Bacteria: In Vitro Research. *Annals of R.S.C.B., ISSN: 1583-6258, Vol. 25, Issue 2, 2021, Pages. 1673 - 1681.*
 43. Achmad H, Auliya N. Management of Malocclusion in Children Using Myobrace Appliance: A Systematic Review. *Annals of R.S.C.B., ISSN: 1583-6258, Vol. 25, Issue 2, 2021, Pages.2120 - 2136.*
 44. Djais AI, Mappangara S, Gani A, Achmad H, Endang S, Tjokro J, Raja N. The Effectiveness of Milkfish (*ChanosChanos*) Scales Chitosan on Soft and Hard Tissue Regeneration In tooth Extraction Socket: A Literature Review. *Annals of R.S.C.B., ISSN: 1583-6258, Vol. 25, Issue 3, 2021, Pages. 8729 - 8752.*
 45. Utama MD, Chotimah C, Achmad H, Arifin NF, Furqani AW. Effect of Solvent Temperature in Effervescent Granule Denture Cleanser with Cacao Pod (*Theobroma cacao* l) 6,5% toward the Growth of *Streptococcus mutans* and *Candida albican*. *Annals of R.S.C.B., ISSN: 1583-6258, Vol. 25, Issue 4, 2021, Pages. 10858 - 10864.*
 46. Yustisia P, Utama MD, Rachmi B, Achmad H, Ilmianti, Yusrini S, Novawati E, Rusdi FJ. The Influence of Fixed Orthodontic Treatment on Tooth Discoloration among Dental Students in Makassar, Indonesia. *Annals of R.S.C.B., ISSN: 1583-6258, Vol. 25, Issue 4, 2021, Pages. 1072010728.11.*