

DAFTAR PUSTAKA

- Adane, T. et al. (2022) "Hematological Abnormalities in Culture Positive Neonatal Sepsis," *Pediatric Health, Medicine and Therapeutics*, Volume 13(May), hal. 217–225. Tersedia pada: <https://doi.org/10.2147/phmt.s361188>.
- Angele, M.K. et al. (2014) "Gender differences in sepsis: Cardiovascular and immunological aspects," *Virulence*, 5(1), hal. 12–19. Tersedia pada: <https://doi.org/10.4161/viru.26982>.
- Alkas Matsas. et al. (2023). "Chronic stress in pregnancy is associated with low birth weight: A Mata Analysis", *Journal of Clinical Medicine*, hal 1-10. Tersedia pada: <https://doi.org/10.3390/jcm12247686>
- Anita Juniatiningsih, Asril Aminullah dan Agus Firmansyah (2008) "Profil Mikroorganisme Penyebab Sepsis Neonatorum di Departemen Ilmu Kesehatan Anak Rumah Sakit Cipto Mangunkusumo Jakarta," *Sari Pediatri*, 10(1), hal. 60–65.
- Bartáková, E. et al. (2019) "Calprotectin and calgranulin C serum levels in bacterial sepsis," *Diagnostic Microbiology and Infectious Disease*, 93(3), hal. 219–226. Tersedia pada: <https://doi.org/10.1016/j.diagmicrobio.2018.10.006>.
- Belachew, A. dan Tewabe, T. (2020) "Neonatal sepsis and its association with birth weight and gestational age among admitted neonates in Ethiopia: systematic review and meta-analysis," *BMC Pediatrics*, 20(1), hal. 1–75. Tersedia pada: [https://morelos.orex.es/bib/47419](https://doi.org/10.1186/s12887-020-02419-w).
- Berzeni, I. (2004) "Diagnostic markers of infection in neonates," *Archives of Disease in Childhood: Fetal and Neonatal Edition*, 89(3), hal. 229–236. Tersedia pada: <https://doi.org/10.1136/adc.2002.023838>.
- Buchori dan Prihatini (2018) "Diagnosis Sepsis Menggunakan Procalcitonin," *Indonesian Journal of Clinical Pathology and Medical Laboratory*, 12(3), hal. 127. Tersedia pada: <https://doi.org/10.24293/ijcpml.v12i3.873>.
- de Camargo, J.F., de Siqueira Caldas, J.P. dan Marba, S.T.M. (2022) "Early neonatal sepsis: Prevalence, complications and outcomes in newborns with 35 weeks of gestational age or more," *Revista Paulista de Pediatria*, 40, hal. 1–8. Tersedia pada: <https://doi.org/10.1590/1984-0462/2022/40/2020388>.
- Campeotto, F. et al. (2004) "High faecal calprotectin concentrations in newborn infants," *Archives of Disease in Childhood: Fetal and*

- Neonatal Edition*, 89(4), hal. 353–356. Tersedia pada: <https://doi.org/10.1136/adc.2002.022368>.
- Castanheira, C.I.G., Corley, K.T.T. dan Lamas, L.R.G.. (2018) “The Influence of Sex in the Incidence and Outcome of Neonatal Sepsis in Horses : a Retrospective Study , 2008 – 2016,” *Equine Veterinary Journal*, 52(2014), hal. 2018. Tersedia pada: <https://doi.org/10.1111/evj.38>.
- Costa, S. et al. (2020) “Effect of Different Milk Diet on the Level of Fecal Calprotectin in Very Preterm Infants,” *Frontiers in Pediatrics*, 8(September), hal. 1–5. Tersedia pada: <https://doi.org/10.3389/fped.2020.00552>.
- Decembrino, L. et al. (2015) “Serum Calprotectin: A Potential Biomarker for Neonatal Sepsis,” *Journal of Immunology Research*, 2015. Tersedia pada: <https://doi.org/10.1155/2015/147973>.
- Downey, L.C., Smith, P.B. dan Benjamin, D.K. (2010) “Risk factors and prevention of late-onset sepsis in premature infants,” *Early Human Development*, 86(SUPPL. 1), hal. 7–12. Tersedia pada: <https://doi.org/10.1016/j.earlhumdev.2010.01.012>.
- Gao, R.Y. et al. (2022) “Calprotectin as a diagnostic marker for sepsis: A meta-analysis,” *Frontiers in Cellular and Infection Microbiology*, 12(November), hal. 1–10. Tersedia pada: <https://doi.org/10.3389/fcimb.2022.1045636>.
- Gerdes, J.S. (2004) “Diagnosis and management of bacterial infections in the neonate,” *Pediatric Clinics of North America*, 51(4), hal. 939–959. Tersedia pada: <https://doi.org/10.1016/j.pcl.2004.03.009>.
- Goold, E., Pearson, L. dan Johnson, L.M. (2020) “Can fecal calprotectin serve as a screen for necrotizing enterocolitis in infants?,” *Clinical Biochemistry*, 84(June), hal. 51–54. Tersedia pada: <https://doi.org/10.1016/j.clinbiochem.2020.06.015>.
- Hadinegoro, S.R.S. et al. (2016) “Diagnosis dan tatalaksana sepsis pada anak,” *Pedoman nasional pelayanan kedokteran Ikatan Dokter Anak Indonesia*, hal. 1–47.
- Hamid, N.F. et al. (2021) “Neonatal sepsis in association with birth weight, gestational age, and mode of delivery in Saudi Arabia: A systematic review and meta-analysis Nazim,” *Medical Science*, 25(111).
- Hapsari, T. (2009) *Kadar seng serum sebagai indikator prognosis pada keluaran sepsis neonatorum*.

- Hoskin, T.S. et al. (2019) "Oxidative cross-linking of calprotectin occurs in vivo, altering its structure and susceptibility to proteolysis," *Redox Biology*, 24(March), hal. 101202. Tersedia pada: <https://doi.org/10.1016/j.redox.2019.101202>.
- Jena, A., Sharma, V. dan Dutta, U. (2023) "Fecal calprotectin in cirrhosis— Does 'ESR' of the intestine measure the liver dysfunction too?," *Indian Journal of Gastroenterology*, 42(6), hal. 754–756. Tersedia pada: <https://doi.org/10.1007/s12664-023-01481-2>.
- Jung Min Yoon, Ju Yi Park (2014)" Fecal Calprotectin concentration in neonatal necrotizing enterocolitis," *Korean Journal of Pediatrics*, 57(8), hal 354. Tersedia pada: <https://doi:10.3345/kjp.2014.57.8.351>
- Kapel N, Campeotto F, Kalach N et al. (2010) Fecal calprotectin in term and preterm neonates. *Journal of Gastroenterology Hepatology and Nutrition* 51 (5): 542 –547. doi: 10.1097/MPG.0b013e3181e2ad7
- Kido, J.I. et al. (2003) "Calprotectin release from human neutrophils is induced by *Porphyromonas gingivalis* lipopolysaccharide via the CD-14-Toll-like receptor-nuclear factor κB pathway," *Journal of Periodontal Research*, 38(6), hal. 557–563. Tersedia pada: <https://doi.org/10.1034/j.1600-0765.2003.00691.x>.
- Larsson, A. et al. (2020) "Calprotectin is superior to procalcitonin as a sepsis marker and predictor of 30-day mortality in intensive care patients," *Scandinavian Journal of Clinical and Laboratory Investigation*, 80(2), hal. 156–161. Tersedia pada: <https://doi.org/10.1080/00365513.2019.1703216>.
- Mahmoud, H.A.H. et al. (2023) "Insight Into Neonatal Sepsis: An Overview," *Cureus*, 15(9), hal. 1–12. Tersedia pada: <https://doi.org/10.7759/cureus.45530>.
- El Manouni El Hassani, S. et al. (2019) "Risk factors for late-onset sepsis in preterm infants: A multicenter case-control study," *Neonatology*, 116(1), hal. 42–51. Tersedia pada: <https://doi.org/10.1159/000497781>.
- Moussa, R. et al. (2016) "Fecal calprotectin levels in preterm infants with and without feeding intolerance," *Jornal de Pediatria*, 92(5), hal. 486–492. Tersedia pada: <https://doi.org/10.1016/j.jped.2015.11.007>.
- Nasution, D.A. (2008) "Faktor Risiko Dan Kesaamaan Jenis Bakteri Jalan Lahir Ibu Dengan Kultur Darah Pada Sepsis Neonatal Awitan Dini."

- Niemelä, O. (2020) "Serum Calprotectin , a Marker of Neutrophil Activation , and Other Mediators of Inflammation in Response to Various Types of Extreme Physical Exertion in Healthy Volunteers," hal. 223–231.
- Odabasi, I.O. dan Bulbul, A. (2020) "Review Neonatal Sepsis," *Sisli Etfal Hastan Tip Bul.*, 54(2), hal. 142–158. Tersedia pada: <https://doi.org/10.14744/SEMB.2020.00236>.
- Pathirana, W.P.N.G.W. et al. (2018) "Faecal calprotectin," *Clinical Biochemist Reviews*, 39(3), hal. 77–90. Tersedia pada: <https://doi.org/10.1097/mcg.0b013e318250e34e>.
- Pietrasanta, C. et al. (2019) "Vascular Endothelium in Neonatal Sepsis: Basic Mechanisms and Translational Opportunities," *Frontiers in Pediatrics*, 7(August), hal. 1–16. Tersedia pada: <https://doi.org/10.3389/fped.2019.00340>.
- Pirr, S. et al. (2017) "High amounts of S100-alarmins confer antimicrobial activity on human breast milk Targeting pathogens relevant in neonatal sepsis," *Frontiers in Immunology*, 8(DEC), hal. 1–10. Tersedia pada: <https://doi.org/10.3389/fimmu.2017.01822>.
- Polin, R.A. et al. (2012) "Management of Neonates with Suspected or Proven Early-Onset Bacterial Sepsis," *Pediatrics*, 129(5), hal. 1006–1015. Tersedia pada: <https://doi.org/10.1542/peds.2012-0541>.
- Pusponegoro, T.S. (2016) "Sepsis pada Neonatus (Sepsis Neonatal)," *Sari Pediatri*, 2(2), hal. 96. Tersedia pada: <https://doi.org/10.14238/sp2.2.2000.96-102>.
- Rosti L, et al. "Formula milk feeding does not increase the release of the inflammatory marker calprotectin, compared to human milk". *Pediatr Med Chir*. 2011 Jul-Aug;33(4):178-81. PMID: 22423477.
- Rougé, C. et al. (2010) "Fecal calprotectin excretion in preterm infants during the neonatal period," *PLoS ONE*, 5(6), hal. 16–19. Tersedia pada: <https://doi.org/10.1371/journal.pone.0011083>.
- Santos, R.P. (2015) "A Practical Guide to the Diagnosis, treatment, and Prevention of Neonatal Infections"62, hal. 491-493
- Simm, M. et al. (2016) "Performance of plasma calprotectin as a biomarker of early sepsis: A pilot study," *Biomarkers in Medicine*, 10(8), hal. 811–818. Tersedia pada: <https://doi.org/10.2217/bmm-2016-0032>.
- Simonsen, K.A. et al. (2014) "Early-onset neonatal sepsis," *Clinical Microbiology Reviews*, 27(1), hal. 21–47. Tersedia pada:

[https://doi.org/10.1128/CMR.00031-13.](https://doi.org/10.1128/CMR.00031-13)

Stefanovic, Iva M. (2011) "Inflammation Neo natalis sepsis," *Biochimia Medica*, 21(3), hal. 276–281.

Suryani, Y.D., Prasetyo, D. dan Hilmanto, D. (2018) "Fecal Calprotectin in Preterm Infants Sepsis with and without Necrotizing Enterocolitis Symptoms," *Global Medical & Health Communication (GMHC)*, 6(3), hal. 188–192. Tersedia pada: <https://doi.org/10.29313/gmhc.v6i3.3337>.

Tita, A.T.N. dan Andrews, W.W. (2010) "Diagnosis and management of clinical chorioamnionitis," *Clinics in Perinatology*, 37(2), hal. 339–354. Tersedia pada: <https://doi.org/10.1016/j.clp.2010.02.003>.

UNICEF (2015) "Maternal and newborn health: Indonesia," *UNICEF for every child*, (000), hal. 142–143.

Utomo, T.T. et al. (2022) "The Association between Birth Route and Early/Late-onset Neonatal Sepsis in Term Infants: A Case-control Study in the NICU of a Tertiary Hospital in East Java," *Indonesia. Iranian Journal of Neonatology*, 13(4), hal. 13. Tersedia pada: <https://doi.org/10.22038/ijn.2022.63955.2237>.

Widger, J., O'Connell, N.H. dan Stack, T. (2010) "Breast milk causing neonatal sepsis and death," *Clinical Microbiology and Infection*, 16(12), hal. 1796–1798. Tersedia pada: <https://doi.org/10.1111/j.1469-0691.2010.03071.x>.

Wu, M. et al. (2024) "Development of risk prediction nomogram for neonatal sepsis in Group B Streptococcus-colonized mothers: a retrospective study," *Scientific Reports*, 14(1), hal. 1–15. Tersedia pada: <https://doi.org/10.1038/s41598-024-55783-2>.

Wulandari, A., Martuti, S. dan Kaswadi, P. (2018) "Perkembangan diagnosis sepsis pada anak," *Sari Pediatri*, 19(4), hal. 237. Tersedia pada: <https://doi.org/10.14238/sp19.4.2017.237-44>.

Wynn, J.L. dan Wong, H.R. (2017) "Pathophysiology of Neonatal Sepsis," *Fetal and Neonatal Physiology*, (January), hal. 1536-1552.e10. Tersedia pada: <https://doi.org/10.1016/b978-0-323-35214-7.00152-9>.